

University of Louisville

ThinkIR: The University of Louisville's Institutional Repository

Faculty Scholarship

1-2013

Assessing depression in cancer patients : a longitudinal comparison of three questionnaires.

Patrick Possel
University of Louisville

Kerstin Knopf

Follow this and additional works at: <https://ir.library.louisville.edu/faculty>



Part of the [Health Psychology Commons](#), and the [Palliative Care Commons](#)

Original Publication Information

Possel, Patrick and Kerstin Knopf. 2013. "Assessing Depression in Cancer Patients -- A Longitudinal Comparison of Three Questionnaires." *Annals of Palliative Medicine*, 2(1): 45-47.

This Article is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Faculty Scholarship by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. For more information, please contact thinkir@louisville.edu.

Assessing depression in cancer patients - a longitudinal comparison of three questionnaires

Patrick Pössel¹, Kerstin Knopf²

¹Department of Educational and Counseling Psychology, University of Louisville, Louisville, KY 40292, USA; ²Palliative Care Team, Univ Hospital, 530 S Jackson St, Louisville, KY 40202, USA

Corresponding to: Patrick Pössel, Dr. rer. soc., Lic Psych. Department of Educational and Counseling Psychology, University of Louisville, 2301 S. Third Street, Louisville, KY 40292, USA. Email: patrick.possel@louisville.edu.



Submitted Dec 26, 2012. Accepted for publication Jan 25, 2013.

doi: 10.3978/j.issn.2224-5820.2013.01.07

Scan to your mobile device or view this article at: <http://www.amepc.org/apm/article/view/1432/2053>

The article by Johns *et al.* (1) compares the Hopkins Symptom Checklist 20-item depression scale (HSCL-20) (2), the Short-Form 36 Mental Health Inventory five-item distress scale (MHI-5) (3), and the Patient Health Questionnaire nine-item depression scale (PHQ-9) (4) in 309 adults with cancer in a longitudinal study. The authors compare internal consistencies and construct validity. Moreover, they address responsiveness of these three scales to results of a systematic treatment of depression over three-months by comparing these measures to patients' self-reports of their depressive symptoms (improved, no change, or worsening). These comparisons provide clinically relevant information considering the fact that studies find comorbidity of cancer with depression ranging from 3-77% with especially high numbers in patients receiving palliative care (for a review see) (5). As Vodermaier *et al.* (6) point out, the wide range of prevalence rates can be partially explained by differences in diagnostic criteria and instruments used to measure depression. Taking this into account and considering the significant negative consequences of depression (including but not limited to functional impairment, reduced quality of life, poor treatment adherence, and even increased mortality; for reviews see) (6-8), Johns *et al.*'s (1) focus on patient-reports of depression with good psychometrics address an important topic.

Johns *et al.* (1) report good internal consistencies and construct validity at baseline for all three scales. Looking into responsiveness, John *et al.* (1) confirm that all three instruments differentiate between patients reporting their depression as improved, unchanged, or worse after three months of systematic treatment and each of these measures

is able to depict a treatment effect when compared to a control group. Finally, using receiver operating curve analysis the scales distinguish improved from non-improved patients. In summarizing, this study is indicating the HSCL-20 (2), the MHI-5 (3), and the PHQ-9 (4) to be appropriate measures to assess depressive symptoms in adult cancer patients based on their psychometric values. From a clinical point of view, this is an encouraging result, as all three measures are short and therefore easily utilized in a patient population oftentimes challenged by limited attention span and fatigue (9). Moreover, the straightforward scoring options for each of the scales increase the likelihood of their application in busy clinical settings.

One other interesting finding that Johns *et al.* (1) report is that two abbreviated versions of the studied instruments (MHI-d and PHQ-2) show less responsiveness to changes in depression when compared to the full length instruments (HSCL-20, MHI-5, PHQ-9). Johns *et al.* (1) use this pattern of findings to engage in the debate whether depression instruments that contain somatic items overestimate the associations between depression and cancer [e.g., (7,10-14)]. This discussion is fueled by the fact that the DSM-IV relies heavily on somatic symptoms for the diagnosis of a depressive disorder and these symptoms are seen commonly in cancer patients as result of their illness progression or as adverse effects of their treatments (e.g., fatigue, poor appetite, sleep problems). Therefore, the interpretation of somatic symptoms has been labeled as one of the most common problems in diagnosing depression in cancer patients (15). However, Johns *et al.* (1) argue their findings that instruments including somatic items show

more responsiveness to demonstrate that somatic items do not lead to an overestimation of depression in cancer patients. Nevertheless, such interpretations need to be made cautiously as change in depression is only assessed by patient-responses to the retrospective question, "Overall, since starting the study, would you say that your depression is worse, about the same, or better?". It might be that patients include their overall well-being and physical symptoms into their own depression evaluation as they are not given any criteria on how to assess their depression. This may be particularly true as patients have been asked this question after repeatedly filling in the questionnaires (including somatic symptoms) that might shape their impression of how the researchers define depression. Therefore, the patient report may at least in part be based on symptoms related to the patient's cancer or treatment side effects rather than their depression. In line with this topic, a recent meta-analysis about depression as predictor of the development of breast cancer found that studies using depression instruments including somatic items found stronger associations between cancer and depression than studies that do not assess these symptoms (16). This underscores the need for future research to clarify this issue further.

In conclusion, Johns *et al.*'s (1) study is set apart from other attempts to evaluate depression scales by taking the extra effort to utilize a longitudinal design. Moreover, including three measures in their research design allows for comprehensive discussions. Given these advantages, it would have been desirable to include instruments with widespread clinical utilization like the Hospital Anxiety and Depression Scale (HADS) (17), even if they have been researched in cross-sectional designs before. Considering that the HADS is one of the most often utilized instruments to measure depression in cancer patients (8) and the fact that recent reviews of depression instruments come to the conclusion that the HADS is one of the best instrument for assessing depression in cancer patients (6-8), the inclusion of the HADS would allow to compare the findings regarding the HSCL, MHI, and PHQ to this "gold standard" of patient-reports in cancer patients.

Depression is one of the most common mental health problems in palliative care (18) but current literature still reports concerning underdiagnosing of depressive disorders in this patient population (19,20). Efforts to provide psychometrically sound tools to assess for depressive symptoms are one important step to address this challenge. As many settings still do not have mental health

professionals readily available to routinely assess all cancer patients (9), future research is needed to focus on how these assessment tools can be used by other professionals (e.g., nurses) to screen for emotional distress in general and depressive symptoms in particular to trigger referrals to specialized professionals who are qualified to diagnose depressive disorders.

Acknowledgements

Disclosure: The authors declare no conflict of interest.

References

1. Johns SA, Kroenke K, Krebs EE, et al. Longitudinal comparison of three depression measures in adult cancer patients. *J Pain Symptom Manage* 2013;45:71-82.
2. Derogatis LR, Rickels K, Uhlenhuth EH, et al. The Hopkins Symptom Checklist: a measure of primary symptom dimensions. In: Pihot P. eds. *Psychological measurement in psychopharmacology: Problems in pharmacopsychiatry*. Basel, Switzerland: Kargerman, 1974:79-110.
3. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473-83.
4. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire*. *JAMA* 1999;282:1737-44.
5. Solano JP, Gomes B, Higginson IJ. A comparison of symptom prevalence in far advanced cancer, AIDS, heart disease, chronic obstructive pulmonary disease and renal disease. *J Pain Symptom Manage* 2006;31:58-69.
6. Vodermaier A, Linden W, Siu C. Screening for emotional distress in cancer patients: a systematic review of assessment instruments. *J Natl Cancer Inst* 2009;101:1464-88.
7. Luckett T, Butow PN, King MT, et al. A review and recommendations for optimal outcome measures of anxiety, depression and general distress in studies evaluating psychosocial interventions for English-speaking adults with heterogeneous cancer diagnoses. *Support Care Cancer* 2010;18:1241-62.
8. Ziegler L, Hill K, Neilly L, et al. Identifying psychological distress at key stages of the cancer illness trajectory: a systematic review of validated self-report measures. *J Pain Symptom Manage* 2011;41:619-36.
9. Knopf K, Head B. As if the cancer wasn't enough... A case

- study of depression in terminal illness. *J Hosp Palliat Nurs* 2012;14:319-29.
10. Bukberg J, Penman D, Holland JC. Depression in hospitalized cancer patients. *Psychosom Med* 1984;46:199-212.
 11. Raison CL, Miller AH. Depression in cancer: new developments regarding diagnosis and treatment. *Biol Psychiatry* 2003;54:283-94.
 12. Dugan W, McDonald MV, Passik SD, et al. Use of the Zung Self-Rating Depression Scale in cancer patients: feasibility as a screening tool. *Psychooncology* 1998;7:483-93.
 13. Kathol RG, Mutgi A, Williams J, et al. Diagnosis of major depression in cancer patients according to four sets of criteria. *Am J Psychiatry* 1990;147:1021-4.
 14. van Wilgen CP, Dijkstra PU, Stewart RE, et al. Measuring somatic symptoms with the CES-D to assess depression in cancer patients after treatment: comparison among patients with oral/oropharyngeal, gynecological, colorectal, and breast cancer. *Psychosomatics* 2006;47:465-70.
 15. Pessin H, Rosenfeld B, Breithart W. Assessing psychological distress near the end of life. *Am J Behav Sci* 2002;46:357-72.
 16. Pössel P, Adams E, Valentine JC. Depression as a risk factor for breast cancer: investigating methodological limitations in the literature. *Cancer Causes Control* 2012;23:1223-9.
 17. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70.
 18. Wilson KG, Chochinov HM, Skirko MG, et al. Depression and anxiety disorders in palliative cancer care. *J Pain Symptom Manage* 2007;33:118-29.
 19. Hallford DJ, McCabe MP, Mellor D, et al. Depression in palliative care settings: the need for training for nurses and other health professionals to improve patients' pathways to care. *Nurse Educ Today*. 2011. Published online September 8, 2011. Available online: <http://dx.doi.org/10.1016/j.nedt.2011.07.011>
 20. Irwin SA, Rao S, Bower K, et al. Psychiatric issues in palliative care: recognition of depression in patients enrolled in hospice care. *J Palliat Med* 2008;11:158-63.

Cite this article as: Pössel P, Knopf K. Assessing depression in cancer patients - a longitudinal comparison of three questionnaires. *Ann Palliat Med* 2013;2(1):45-47. doi: 10.3978/j.issn.2224-5820.2013.01.07