University of Louisville

ThinkIR: The University of Louisville's Institutional Repository

Electronic Theses and Dissertations

5-2012

Exploration of the chemistry of alkynes and Selectfluor : search for cytotoxic agents from the Amazonian rainforest.

Zhuang Jin
University of Louisville

Follow this and additional works at: <https://ir.library.louisville.edu/etd>

Part of the [Chemistry Commons](#)

Recommended Citation

Jin, Zhuang, "Exploration of the chemistry of alkynes and Selectfluor : search for cytotoxic agents from the Amazonian rainforest." (2012). *Electronic Theses and Dissertations*. Paper 691.
<https://doi.org/10.18297/etd/691>

This Doctoral Dissertation is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. This title appears here courtesy of the author, who has retained all other copyrights. For more information, please contact thinkir@louisville.edu.

EXPLORATION OF THE CHEMISTRY OF ALKYNES AND
SELECTFLUOR. SEARCH FOR CYTOTOXIC AGENTS FROM THE
AMAZONIAN RAINFOREST

By

Zhuang Jin

A Dissertation
Submitted to the Faculty of the
College of Arts and Sciences of the University of Louisville
in Partial Fulfillment of the Requirements
for the Degree of

Doctor of Philosophy

Department of Chemistry
University of Louisville
Louisville, Kentucky

May 2012

EXPLORATION OF THE CHEMISTRY OF ALKYNES AND
SELECTFLUOR. SEARCH FOR CYTOTOXIC AGENTS FROM THE
AMAZONIAN RAINFOREST

By

Zhuang Jin

A Dissertation Approved on

March 5, 2012

by the following Dissertation Committee:

Dr. Gerald B. Hammond
Dissertation Director

Dr. Paula J. Bates

Dr. Craig A. Grapperhaus

Dr. Frederick A. Luzzio

Dr. Michael H. Nantz

DEDICATION

This dissertation is dedicated to three ladies:

My wonderful mother Xiulan Hai

My beautiful wife Lanlan Bao

My gorgeous daughter Solonga Julia Jin

ACKNOWLEDGEMENTS

I am very grateful to my mentor, Dr. Gerald B. (GB) Hammond for giving me an opportunity to find beauty in discovery, for always keeping his trust in me, and for showing me that the real targets in life are not always chemical structures. I also greatly appreciate Dr. Bo Xu who has been my day-to-day mentor in all aspects of my research, providing me with guidance and ideas for my research, and helping me to find out ‘why my reaction didn’t work...’

I want to express my deep appreciation to our collaborators: Dr. Paula J. Bates (James Graham Brown Cancer Center, University of Louisville) and Dr. Abraham J. Vaisberg (Universidad Peruana Cayetano Heredia, Lima, Perú) for their support on the biological aspects of my natural product research; Dr. Stephen G. DiMagno (University of Nebraska) for his suggestions and help on the fluorine chemistry aspects of my work; Dr. Walter H. Lewis (Washington University St Louis) for his taxonomical work on the Amazonian plants; and, last but not least, Dr. Jose C. Aponte (former member of the Hammond’s group, now at Brown University) for introducing me to the study of Peruvian medicinal plants.

I want to thank the faculty members serving in my thesis committee members Dr. Gerald B. Hammond, Dr. Paula J. Bates, Dr. Craig A. Grapperhaus, Dr. Frederick A. Luzzio, and Dr. Michael H. Nantz. I am also grateful to our HRMS expert Dr. Richard M. Higashi, to

our X-ray manager Dr. Mark S. Mashuta, to the instrumental specialist Dr. William N. Richmond and to our NMR manager Dr. Neal J. Stolowich for their technical help. I also want to acknowledge the help and friendship from all faculty and staff in the Department of Chemistry of the University of Louisville.

I am lucky to have joined Dr. Hammond's group and to have worked side by side with the many talented chemists in his group, both, past and present. I have learned very much from all of them. I especially want to single out Dr. Leping Liu, Dr. Weibo Wang, Dr. Satoru Arimitsu, Junbin Han, Deepika Malhotra, Manish Kumar, Rachel S. Hidinger, Rebecca Bottom, Majeid Ali and Han Yang.

I am also thankful to my former mentor Dr. Gereltu Borjihan, of the Institute of Macromolecular Chemistry and Mongolian Medicine, Inner Mongolia University, Inner Mongolia, China. He provided me with a solid background in bioassay-directed fractionation techniques, which were helpful to my graduate studies in Louisville. All past and present members of his group are remembered, but most importantly, Lanlan Bao, whom I first met in his lab, and who would become my lovely wife and mother of our beautiful daughter Solonga Julia Jin.

It would have been impossible to finish my graduate studies in the United State without the love, support, patience and encouragement from my family. Especially, I want to thank my wonderful mother Xiulan Hai, who taught me the virtues of perseverance and character.

Finally, my work would not have been possible if it wasn't for the financial and material support from the various funding agencies, especially the National Science Foundation,

the Department of Defense, the Institute of Molecular Diversity and Drug Design of the University of Louisville (IMD3), and The American Society of Pharmacognosy (ASP).

ABSTRACT

EXPLORATION OF THE CHEMISTRY OF ALKYNES AND SELECTFLUOR. SEARCH FOR CYTOTOXIC AGENTS FROM THE AMAZONIAN RAINFOREST

Zhuang Jin

March 5, 2012

Vicinal dithioethers and alkenyl thioethers were synthesized under environmentally friendly conditions using alkyne and thiol in water. Alkynes were also used to develop a multibond-forming reaction that formed cyclic ketones or ketoesters through a gold-catalyzed intramolecular oxygen transfer isomerization of 2-alkynyl-1,5-diketones or 2-alkynyl-5-ketoesters.

The investigation of Selectfluor chemistry yielded a highly stereoselective synthesis of fluoroalkyl (*E*)- α,β -unsaturated ketones from allenyl esters, through a gold-catalyzed rearrangement that produced an intermediate dienyl ester. When Selectfluor was combined with copper, it produced two oxidative systems, F-TEDA-BF₄ and F-TEDA-PF₆, both of which efficiently converted amides into imides at room temperature in short time, but whereas the former employed stoichiometric amounts of copper(I), the latter only needed catalytic amounts.

The bioassay-directed fractionation of *Physalis angulata* L. afforded three new antiproliferative withanolides: physangulidines A, B and C. Each has shown significant

cytotoxic activity (GI_{50} is less than $4\mu\text{g/mL}$) on DU145 and RWPE-1 *in vitro*. In addition, compared to positive drug 5-fluorouracil, physangulidine A had significant cytotoxic activity on different cells. The bioassay-guided fractionation of *Crematosperma microcarpum* led to the isolation and identification of dehydrodiisoeugenol as its main cytotoxic agent. Lastly, selected fractions of *Hyptis lantanaefolia* have shown promising cytotoxic activities; with one of semipurified chromatographic fraction (LHL15) exhibiting very high bioactivity on various cell lines (IC_{50} is lower than 50 ng/mL).

TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	iv
ABSTRACT.....	vii
TABLE OF CONTENTS.....	ix
LIST OF FIGURES	xi
LIST OF TABLES.....	xii
LIST OF SCHEMES	xiii
1. OVERVIEW OF THIS DISSERTATION.....	1
1.1. Alkyne and Fluorine Chemistry	1
1.2. Natural Products Research	3
2. EXPLORATION OF ALKYNE CHEMISTRY.....	5
2.1. Environmentally Friendly Addition of Thiols to Alkynes	5
2.1.1 Background	5
2.1.2 Scope of the Aqueous Dihydrothiolation of Alkynes	6
2.1.3 Regio- and Stereo-selective Monohydrothiolation of Non-Terminal Propargyl Alcohols	10
2.1.4 Summary	15
2.1.5 Experimental Section.....	16
2.2. Gold-Catalyzed Intramolecular Oxygen Transfer Reaction of 2-Alkynyl-1,5- diketones and 2-Alkynyl-5-Ketoesters	30
2.2.1 Background and Introduction	30
2.2.2 Result and Discussion	31
2.2.3 Experimental Section.....	33
3. EXPLORATION OF THE CHEMISTRY OF SELECTFLUOR.....	41
3.1 Selectfluor TM	41
3.2 Stereoselective Synthesis of Fluoroalkyl α,β -Unsaturated Ketones.....	43

3.2.1	Background	43
3.2.2	Our Efficient Strategy to Prepare Ketone 3-2-1	45
3.2.3	Summary	49
3.2.4	Experimental Section	50
3.3.	Combination of Copper and Selectfluor: a Strong yet Selective Oxidant System 57	
3.3.1	Hydroxyl Group Directed C-H Amination	57
3.3.2	Copper-Mediated Oxidation of Amides into Imides by Selectfluor (F-TEDA- BF ₄)	62
3.3.3	Copper Catalyzed Oxidation of Amide into Imide by F-TEDA-PF ₆	68
4.	SEARCH FOR CYTOTOXIC AGENTS FROM THE AMAZONIAN RAINFOREST	81
4.1.	Background	81
4.2.	<i>Physalis angulata</i> L. (Solanaceae)	83
4.3.	<i>Crematosperma microcarpum</i> (Annonaceae)	94
4.4.	<i>Hyptis lantanaefolia</i> (Lamiaceae)	95
4.5	<i>Bocconia intengrifolia</i> (Papaveraceae)	96
4.6	Other Plant Species Studied	97
4.7.	Summary	97
4.8.	Experimental Section	98
	REFERENCES	112
	APPENDIX A: LIST OF ABBREVIATIONS	128
	APPENDIX B: COPYRIGHT PERMISSION	130
	APPENDIX C: NMR SPECTRA FOR NEW COMPOUNDS	135
	CURRICULUM VITAE	243

LIST OF FIGURES

Figure 1. Outline of Natural Products Chemistry	4
Figure 2. NOSEY Spectra of 2-1-4c	12
Figure 3. Natural Product Containing the Cyclopentenone Core Structure.....	30
Figure 4. Monofluoroalkyl (<i>E</i>)- α,β -Unsaturated Ketones A , B and C are Important Building Blocks for Bioactive Agents	43
Figure 5. Therapeutic and Agrochemical Agents Containing the Imide Moiety.....	62
Figure 6. HRMS of Imide 3-3-2-2b and O ¹⁸ labeled 3-3-2-2b	80
Figure 7. ¹³ C NMR of Carbonyl Groups in Imide 3-3-2-2b and O ¹⁸ labeled 3-3-2-2b	80
Figure 8. Drugs and Drug Candidates Isolated from Natural Source	81
Figure 9. Structures of Physangulidines A (1), B (2) and C (3)	84
Figure 10. Carbon Numbering and Crucial ROESY Interactions of Physangulidines A (1), B (2) and C (3)	85
Figure 11. An <i>ORTEP</i> -3 Diagram of 1 Showing 40% Ellipsoids.....	85
Figure 12. Main structures of Withanolides Have Been Isolated	86
Figure 13. An <i>ORTEP</i> -3 Diagram of 2 Showing 40% Ellipsoids.....	87
Figure 14. Selected HMBC Interactions in Physangulidines A , B and C	93
Figure 15. Selected ROESY Interactions in Physangulidines A , B and C	93
Figure 16. Structure of Trans-dehydrodiisoeugenol	94
Figure 17. Cytotoxic Activity of Fractions LHL-14 and LHL-15	96
Figure 18. Cytotoxic Activity of SPA, SPA1 and SPA2	104
Figure 19. Cytotoxic Activity of SPA11, SPA12, SPA13 and SPA14.....	105
Figure 20. Cytotoxic Activity of SPA14-3-2, SPA14-3-3 and SPA14-3-4	106

LIST OF TABLES

Table 1. Effect of Solvents on Dihydrothiolation of Alkyne 2-1-1a .	7
Table 2. Dihydrothiolation of Alkyne 2-1-1 .	8
Table 3. Regio and Stereoselective Monohydrothiolation of Propargyl Alcohols.	11
Table 4. Isomerization of 2-Alkynyl-1,5-Diketones and 2-Alkynyl-5-Ketoesters	31
Table 5. Preparation of α,β -Unsaturated Ketones 3-2-1 from <i>in Situ</i> Generated Dienes 3-2-2 .	46
Table 6. Screening for a Suitable Transition Metal Catalyst	58
Table 7. Substrate Scope of Hydroxyl Group Directed C-H Amination	59
Table 8. Screening for Optimal Oxidation Conditions	64
Table 9. Comparison of the Effectiveness of Selectfluor (F-TEDA- BF ₄) (Condition A) and F-TEDA- PF ₆ (Condition B) in Amide Oxidation	66
Table 10. List of Targeted Plants	82
Table 11. All NMR Data of Physangulidine A (1)	89
Table 12. All NMR Data of Physangulidine B (2)	90
Table 13. All NMR Data of Physangulidine C (3)	91
Table 14. GI ₅₀ Values (μ M) of Physangulidine A and 5-Fluororacil on Different Cells.	92
Table 15. GI ₅₀ Values (μ M) of ($\pm$)- <i>Trans</i> -dehydrodiisoeugenol and Doxorubicin on Different Cells	94
Table 16. NMR data for ($\pm$)- <i>Trans</i> -dehydrodiisoeugenol and Literature Data	95

LIST OF SCHEMES

Scheme 1. Gold Catalyzed Functionalized Hydration of Alkynes	1
Scheme 2. Outline of Alkyne and Selectfluor Chemistry	2
Scheme 3. Synthesis of Vicinal Dithioethers and Alkenyl Thioethers from Alkynes and Thiols	6
Scheme 4. Nucleophilic and Radical Additions of Thiol to Propargyl Alcohols.....	13
Scheme 5. Regioselectivity and Stereoselectivity Considerations	14
Scheme 6. Synthetic Transformations of Alkenyl Thioethers	15
Scheme 7. ¹⁸ O Isotopic Labeling Experiment for Mechanistic Studies	32
Scheme 8. Commercially Synthetic pathway for Selectfluor	41
Scheme 9. Literature Construction of Monofluoroalkyl α,β -Unsaturated Ketones by Fluorine Containing Building Blocks (Top) and Fluorination of Specific Ketones (Bottom)	44
Scheme 10. Lengthy Preparation Pathway for a Fluorine-Containing Wittig Reagent....	44
Scheme 11. Gold-catalyzed Isomerization of Alkyne and Allene and Following Fluorination Strategy	45
Scheme 12. Limitations of Our Protocol	48
Scheme 13. Proposed Mechanism for Fluorinated Cation Enabled Isomerization	49
Scheme 14. Hydroxyl Group Directed C-H amination by Selectfluor in Acetonitrile	57
Scheme 15. Evidence of Trace H ₂ O as Ultimate Oxygen Source	67
Scheme 16. Tertiary Amide not Suitable for This Oxidation	67
Scheme 17. Limitation of The Copper Catalyzed/Mediated oxidations.....	69
Scheme 18. Nicolaou's Proposal of Amide Oxidation by DMP	69
Scheme 19. Attempt to Capture Possible Imine Intermediate	70
Scheme 20. Proposed Rationale for Different Effect of BF ₄ ⁻ and PF ₆ ⁻	72
Scheme 21. Bioassay Directed Fractionation of <i>P. angulata</i> L	103
Scheme 22. Bioassay Directed Fractionation of <i>C. microcarpum</i>	107
Scheme 23. Bioassay Directed Fractionation of <i>H. lantanaefolia</i>	108

Scheme 24. Bioassay Directed Fractionation of <i>B. intengrifolia</i>	110
---	-----

1. OVERVIEW OF THIS DISSERTATION

1.1. Alkyne and Fluorine Chemistry

Alkyne is an important synthon in organic chemistry, as demonstrated by the many examples of elegant synthetic work centered on alkynes.¹ The Hammond group has dedicated much of their efforts to explore the chemistry of alkynes.² Of more relevance to the work presented herein, members of the Hammond group have recently reported a cationic gold-catalyzed functionalized hydration of alkynes (Scheme 1).³

Scheme 1. Gold Catalyzed Functionalized Hydration of Alkynes

Various types of alkynes were hydrated to give α -substituted α -fluorinated ketones using boronic acid, Selectfluor and trace amount of water in the presence of a cationic gold catalyst. The synthetic work contained in this dissertation is centered around alkynes and Selectfluor. This work (Scheme 2) includes the synthesis of vicinal dithioesters and alkenyl thioesters via a new environmentally friendly use of alkynes and thiols in water; the gold-catalyzed intramolecular oxygen transfer reaction of 2-alkynyl-1,5-diketones or 2-alkynyl-5-diketoesters; and the stereoselective synthesis of fluoroalkyl *E*- α,β -

unsaturated ketones via gold-catalyzed rearrangement of allenyl carbinol ester followed by fluorination. Finally, two new oxidative systems containing Selectfluor have been investigated, namely the combination of CuBr (stoichiometric amounts) with excess Selectfluor (F-TEDA-BF₄), and the combination of catalytic CuBr with excess F-TEDA-PF₆--the anionic exchange product of Selectfluor. Both combinations can efficiently oxidize amides into imides at room temperature.

A. Alkyne Chemistry

Chapter 2. *Eur. J. Org. Chem.* **2010** 168
J. Organomet. Chem. **2011** 696, 269

Chapter 2. *Chem. Eur. J.* **2011** 17, 10690.

B. Selectfluor Chemistry

Chapter 3. *Tetrahedron Lett.*, **2011** 52, 1956.

Chapter 3. Manuscript in preparation

Scheme 2. Outline of Alkyne and Selectfluor Chemistry

1.2. Natural Products Research

Nature as source for drugs and inspiration for the development of front-line-drugs is indisputable if one considers that more than 75% of the world's population, especially in developing countries, rely on medicinal plants as primary source of medicinal care,⁴ and that around 50% of the total new chemical entities launched onto the market in the last 25 years are natural products or natural products analogs.⁵

In order to demonstrate the medicinal value of natural products, a multidisciplinary and international group of scientists led by Professors Abraham J. Vaisberg (Universidad Peruana Cayetano Heredia), Walter H. Lewis (Washington University, St. Louis), and Gerardo Lamas (Universidad Nacional Mayor de San Marcos), under the auspices of the International Cooperative Biodiversity Group (ICBG) project--funded by a consortium of the following agencies, NIH, US AID, and NSF--embarked on an ethnobotanical study of Peruvian medicinal plants, guided by members of the Aguaruna community.⁶ The Aguaruna are one of the four tribes that constitute the Jivaro linguistic family, living in the upper Amazon basin and adjacent foothills of the eastern Andes Mountains of northern Perú and nearby Ecuador. Northern Perú, particularly the eastern slopes of the Andes Mountains and adjacent upper Amazon basin, is exceedingly rich in diverse woody plants. Only a small percentage (~2%) of these Peruvian species have been investigated chemically and/or biologically, and it is estimated that over 2000 plants in the Amazon region are used in traditional medicine.⁶ Since the early 1990's this group has collected 3591 plants, with, 1600 of them being rare or/and medicinal plants. Our group has isolated numerous bioactive agents from targeted plant through a bioassay-

guided fractionation,⁶⁻⁷ then built library of drug candidates around them by synthesizing derivatives of the bioactive agents.⁸

This dissertation includes our results on 13 plant species collected in the Amazonian rain forest. Trans-dehydrodiisoeugenol was isolated as the main cytotoxic agent present in *Crematosperma microcarpum* (Annonaceae) by bioassay-directed isolation. Similarly, three new antiproliferative withanolides with an unusual carbon framework, namely, physangulidines A, B and C were isolated and characterized from *Physalis angulata* (Solanaceae) (Figure 1). We have also found that selected fractions of *Hyptis lantanaefolia* (Lamiaceae) have promising cytotoxic activities; one of its semipurified chromatographic fraction (LHL15) has shown extremely high cytotoxic activity on different cell lines (IC_{50} is lower than 50 ng/mL). *Hyptis lantanaefolia*, together with *Bocconia intengrifolia* (Papaveraceae), are currently under investigation.

Figure 1. Outline of Natural Products Chemistry

2. EXPLORATION OF ALKYNE CHEMISTRY

2.1. Environmentally Friendly Addition of Thiols to Alkynes

2.1.1 Background

Organosulfur compounds have become increasingly important as the role of sulfur is probed deeper in biological processes, new materials, and chemical synthesis.⁹ As a result, the synthesis of organosulfur compounds has attracted much attention.¹⁰ Specifically, vicinal dithioethers and alkenyl thioethers have been widely used as target or intermediates. For example, vicinal dithioethers have been used as ligands for zirconium or titanium complexes for alkene polymerization and hydroamination.¹¹

Vicinal dithioethers can be made by nucleophilic substitutions,^{11b, 11d} or nucleophilic ring-opening reaction of thiolate.^{11a, 11c, 11e} Vicinal dithioethers can also be prepared from an alkene and a disulfide under acid or metals catalysis.¹² Hydroelementation is a versatile and atom-efficient method for installing heteroelements to unsaturated carbon-carbon bonds.¹³ Therefore, one of the most straightforward methods to make vicinal dithioethers is the dihydrothiolation of an alkyne. Under controlled conditions, monohydrothiolation of alkynes could yield alkenyl thioethers (Scheme 3). There are reports on the preparation of vicinal dithioethers from alkynes and thiols in organic solvents¹⁴ using various radical initiators, and/or heating or UV light. The use of water as solvent has also been reported

by Oshima and coworkers, who isolated the vicinal dithioether as a side product, during their investigation of thiol-yne radical reactions in water assisted by a water-soluble radical initiator.¹⁵ In all these literature syntheses, there are detracting experimental limitations, including the use of metal catalysts, high temperatures or radical initiators, and organic solvent. The addition of thiols to alkenes has been used in the synthesis of dendrimers (so called ‘thiol-ene click’ chemistry).¹⁶ The spectrum of application of these dendrimers, ranging from medicine to nanoengineering¹⁷ should spur the development of novel and efficient synthesis of macromolecules. In this chapter we report an atom-economical and ‘green’ synthesis of vicinal dithioethers from the reaction of alkyne and thiol without the use of metal catalysts or radical initiators, and using water as the sole solvent. Furthermore, this method can also be used for regio- and stereo-selective monohydrothiolation of propargyl alcohols, which leads to an effective synthesis of alkenyl thioethers with exclusive *E*-selectivity.

Scheme 3. Synthesis of Vicinal Dithioethers and Alkenyl Thioethers from Alkynes and Thiols

2.1.2 Scope of the Aqueous Dihydrothiolation of Alkynes

We used the reaction of 1-hexyne and 4-methylbenzenethiol as our model reaction (Table 1). While THF, MeOH, H₂O, individually or as a mixture, or non-solvent conditions gave

good results, traditional organic solvents, such as CH₃CN, CH₂Cl₂ and toluene, were not satisfactory.

Table 1. Effect of Solvents on Dihydrothiolation of Alkyne **2-1-1a**.^a

$n\text{-C}_4\text{H}_9\text{-C}\equiv\text{CH}$ + $\text{C}_6\text{H}_4\text{-SH}$ $\rightarrow$ $\text{C}_6\text{H}_4\text{-S-CH}_2\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-S-C}_6\text{H}_4$

2-1-1a **2-1-2a** **2-1-3a**

Entry	Solvent	Yield ^b (%)
1	CH ₃ CN	31
2	CH ₂ Cl ₂	24
3	Toluene	34
4	THF	72
5	MeOH	65
6	H ₂ O:MeOH (4:1)	76
7	H ₂ O	80
8	No solvent	72

^a **2-1-1a** (1 mmol), **2-1-2a** (2.4 mmol) and solvent (0.5 mL) reacted for 24 h at r.t.

^b Determined by ¹HNMR, benzyl bromide (0.4 mmol) as internal standard.

Considering both, chemical yields and the increasing environmental consciousness of the scientific community, we chose water as the ideal reaction media. It is surprising that water was found to be the best solvent for the reaction considering that both starting material and product are virtually insoluble in water. According to Sharpless and co-workers,¹⁸ reaction rates can be accelerated when insoluble reactants were stirred in aqueous suspension, denoted as “on water” conditions. Thus, this reaction could be considered as an ‘on water’ reaction. With optimized conditions in hand, we examined the scope of this dihydrothiolation reaction; the results are summarized in Table 2.

Table 2. Dihydrothiolation of Alkyne **2-1-1**.^a

$$\text{R}^1\text{—}\text{C}\equiv\text{C—R}^2 + \text{R}^3\text{—SH} \xrightarrow{\text{H}_2\text{O}} \text{R}^1\text{—CH(R}^2\text{)—CH(R}^3\text{)—SR}^3$$

2-1-1 **2-1-2** **2-1-3**

Entry	Alkyne 2-1-1	Thiol 2-1-2	Temp	Product 2-1-3 (%) ^b
1			r.t.	2-1-3a , 73
2	2-1-1a		r.t.	2-1-3b , 74
3	2-1-1a		r.t.	2-1-3c , 75
4		2-1-2a	r.t.	2-1-3d , 79
5		2-1-2b	r.t.	2-1-3e , 96
6		2-1-2c	60 °C	2-1-3f , 44
7		2-1-2c	60 °C	2-1-3g , 55, d.r.=1:1
8	2-1-1a		60 °C	2-1-3h , 31
9			60 °C	2-1-3i , 81
10	2-1-1c	2-1-2e	60 °C	2-1-3j , 70

^a Alkyne **2-1-1** (1 mmol), thiol **2-1-2** (2.4 mmol) and deionized water (0.5 mL), reacted at indicated temperature for 24 h. ^b Isolated yields.

1-Hexyne and aryl thiols reacted smoothly (entries 1, 2 and 3). The reaction of functionalized terminal alkynes, such as methyl propargyl ether and 4-pentyn-1-ol, with

an aryl thiol also gave good to excellent yields (entries 4 and 5). However, propargyl alcohol and 2-hexyne reacted with 4-chlorothiophenol to produce moderate yields of **2-1-3** (entries 6 and 7). The low yield in entry 8 is probably due to the volatility of the substrate, as higher molecular weight alkynes reacted effectively with aliphatic thiols under mild heating (entries 9 and 10).

Unfortunately, under our conditions, phenylacetylene and other internal alkynes reacted with thiols to give only the monohydrothiolation product.^{12c, 19} Upon further investigation, we found that aryl thiols are more reactive than aliphatic thiols; electron withdrawing chlorine accelerates the dihydrothiolation, and weak electron donating groups slow the reaction. The observed reaction rates follow this order: 4-chlorothiophenol > thiophenol > 4-methylbenzenthio. Strong electron withdrawing groups, such as nitro, methyl ester and carboxylic acid, on the *para* position of thiophenol hinder the reaction. This may be related to their ease of emulsification, as 4-nitrothiophenol, 4-mercaptobenzoic acid and 4-mercaptobenzoic acid methyl ester do not form an emulsion with alkynes, whereas the thiol in Table 2 readily formed an emulsion with alkyne in water upon stirring.

Our mechanistic studies hinted that the reaction probably proceeded through a radical mechanism, because no reaction occurred in the presence of galvinoxyl free radical (1.1 equiv to thiol).²⁰ The possibility of a nucleophilic addition could be ruled out since nucleophilic dihydrothiolation of alkynes normally gives thioacetals.²¹ Furthermore, small amounts of disulfide (less than 5% yield based on thiol), formed from the homocoupling of thiolate radical, were observed in the course of this study, which is consistent with the proposed radical mechanism. Furthermore, no hydration product was

found in the reaction mixture. The radical initiator could be dioxygen in the air or light since reaction, done in dark area, gave low yield (entry 1 in Table 2, gave 30%). The specific role of the solvent is not clear at this time, it seems water has some ability to stabilize the radical intermediate and therefore facilitates the radical-mediated reaction. A literature report²² speculated that a hydrogen bond between thiol and water could enable a nucleophilic addition to the alkene.

2.1.3 Regio- and Stereo-selective Monohydrothiolation of Non-Terminal Propargyl Alcohols

While investigating dihydrothiolation conditions for non-terminal propargyl alcohols (e.g. **2-1-1g**), we found, to our surprise, that only the monohydrothiolation product was obtained, and this reaction proceeded in a regio- and stereo-selective manner. The thiol only attacked the carbon next to the alcohol and only the *E*-isomer was isolated. The reaction scope is shown in Table 3. But-2-yn-1-ol and 4-chlorothiophenol gave a moderate yield of product with high stereoselectivity (entry 1). Similarly, **2-1-1h** reacted with three different thiophenols in moderate to high yields with high *E*-selectivity (entries 2-4). Entry 5 also showed satisfactory yield and stereoselectivity. Reaction of secondary alcohols (**2-1-1j** and **2-1-1k**) with thiophenols gave satisfactory yields and excellent stereoselectivity (entries 6-9), while tertiary propargyl alcohol **2-1-1l** gave very high stereoselectivity (100% *E*), albeit in low yield (entry 10). Reaction of **2-1-1m** and 4-chlorothiophenol **2-1-2c** was also highly stereoselective (entry 11).

Table 3. Regio and Stereoselective Monohydrothiolation of Propargyl Alcohols.^a

Entry	Alkyne 2-1-1	Thiol 2-1-2	Product 2-1-4	Yield ^b
1				53 (<i>E/Z</i> =9:1)
2				76 (<i>E/Z</i> =9:1)
3				66 (<i>E/Z</i> =12.5:1)
4				58 (<i>E/Z</i> =12.5:1)
5				77 (<i>E/Z</i> =12.5:1)
6				64 (<i>E/Z</i> =17:1)
7				63 (<i>E/Z</i> =17:1)
8				78 (<i>E/Z</i> =17:1)
9				70 (<i>E/Z</i> =20:1)
10				17 (<i>E/Z</i> =100:0)
11				65 (<i>E/Z</i> =100:0)
12				42 (<i>E/Z</i> =100:0)
13				44 (<i>E/Z</i> =50:1)

^a Alkyne **2-1-1** (1 mmol), thiol **2-1-2** (1.2 mmol) and deionized water (0.25 mL) reacted at r.t. for 12 h.

^b Isolated yield, *E* and *Z* were determined by NOESY study.

In entries 12 and 13, we investigated substrates having a phenyl group; both of them, without exception, reacted with aryl thiols to give products with very high stereoselectivity. Stereochemistry of alkenyl thioethers (**2-1-4**) was determined by NOSEY study, and selected NOSEY spectra of **2-1-4c** is shown in Figure 2, which showing that Stereochemistry of **2-1-4c** is mainly *E*.

Figure 2. NOSEY Spectra of **2-1-4c**

During this study, we also found that alkyl thiols are not effective under these conditions, and terminal alkynes, such as propargyl alcohol, overreacted with thiophenol to give dihydrothiolation products (Table 2). Regio- and stereo-selective monohydrothiolation was observed only with non-terminal propargyl alcohols, even with excess thiol in water. While the monohydrothiolation of non-terminal propargyl alcohol with thiol has been

studied before,²³ our mild reaction conditions showed distinctive regio- and stereoselectivity for a wider range of substrates. A case in point is the base mediated nucleophilic addition of thiol to propargyl alcohol (Scheme 4, top) that yields the *trans* adduct product, *Z*-alkenyl thioether^{23a-g} or a mixture of *Z* and *E*-alkenyl thioethers.^{23h, i}

Scheme 4. Nucleophilic and Radical Additions of Thiol to Propargyl Alcohols

And, addition of a thiol, using a radical initiator, usually affords a mixture of *Z* (predominant) and *E*-alkenyl thioethers (Scheme 4, bottom).^{23j, k}

The reason for the high regio and stereoselectivity of **2-1-4** can be explained by the relative stability of vinyl radical intermediates (Scheme 5, top). The vinyl radical intermediates **C** and **D** are higher in energy than **A** and **B** (Gaussian 03, UB3LYP/6-311+G(d)). This can explain why the thiol always attacks the carbon next to hydroxymethyl group. The stereoselectivity outcome of **2-1-4** may be explained by steric effects and the isomerization of vinyl radical **A** to **B**. It is widely accepted that a hydrogen donor can approach from the less hindered side of the vinyl radical.²⁴

Scheme 5. Regioselectivity and Stereoselectivity Considerations

The bulkiness of hydroxymethyl group in radical **A** may hinder the approach of the hydrogen donor (thiol) to form a *Z*-isomer (Scheme 5, bottom) (Note: hydroxyl methyl group is bulkier than thiol ester group, because bond length between sp^2 and sp^3 carbons is 150 pm²⁵ and bond length between sp^2 carbon and sulfur is 175 pm²⁶). However, the thiol can easily approach the less hindered side of the vinyl radical **B** to produce the *E*-isomer. As a result, radical **A** may slowly isomerize to radical **B**. This is consistent with the fact that the bulkier the hydroxyalkyl, the higher the stereoselectivity observed (Table 3). And, the mild conditions (room temperature/water as solvent) minimized the

isomerization of the final product. This is in contrast with literature reports in which isomerization may occur at high temperature (Scheme 5, bottom).^{23j, k}

Alkenyl thioesters have already demonstrated their utility.²⁷ Compound **2-1-4** can be used in further transformations; for example, the free hydroxyl group in **2-1-4** can be protected to give **2-1-5**, and **2-1-4** can be easily oxidized to sulfone **2-1-6** with *m*-CPBA (Scheme 6). Both **2-1-5** and **2-1-6** have been used in nickel catalyzed cross-coupling reactions with Grignard reagents²⁸ or organozinc reagents²⁹ to give functionalized allylic alcohols, which are important intermediates in total synthesis³⁰ and methodology.³¹

Scheme 6. Synthetic Transformations of Alkenyl Thioethers

2.1.4 Summary

We have found that a wide range of alkynes react with excess thiols to give vicinal dithioethers under mild condition using only water as solvent, without radical initiator or UV light sources. Alternatively, non-terminal propargyl alcohols reacted with phenyl thiols in water to produce *E*-alkenyl thioethers in highly regio- and stereo-selective fashion. Considering the mild conditions of our reaction, it could have great potential in the preparation of highly branched dendrimers, and as linkers incorporating fluorescence tags in biological applications.

2.1.5 Experimental Section

General

Substrates **2-1-1m** and **2-1-1o** were prepared using a literature method.³² Other substrates and reagents were commercially obtained from Alfa or Adrich, and were used without further purification. Structures were identified by NMR spectra, assisted by elemental analysis or/and IR spectra. ¹H and ¹³C NMR spectra were recorded at 500 and 125 MHz, respectively, using CDCl₃ as solvent. The chemical shifts were reported in δ (ppm) value relative to CDCl₃ (7.26 ppm for ¹H NMR and 77.0 for ¹³C NMR), and multiplicities are indicated by s (for singlet), d (doublet), dd (double doublet), m (multiplet), and br (broad). Coupling constant, *J*, was reported in Hz. Elemental analysis was performed at Atlantic Microlabs Inc., Norcross, Georgia 30091. Melting points were measured using a DigiMelt PMA 160 melting point apparatus.

Typical procedure for preparation of **2-1-1m** and **2-1-1o**

To oven dried 10 mL reaction flask containing 1-Hexyne (340 μ L, 3.0 mmol), anhydrous THF (5.0 mL) and stir bar at -78 °C, n-Butyl lithium (3.0 mmol) was added over 5 min by syringe, and slowly warm up to room temperature and stirred for another 2 h. Then benzaldehyde (306 μ L, 3.0 mmol) was added into the above reaction mixture by syringe, and stirred at room temperature for 2 h. The reaction mixture was quenched by saturated ammonium chloride (20 mL) and followed by diethyl ether extraction (25 mL $\times$ 4). The ether layers were combined and dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to yield amide **2-1-1o** (526.0 mg, 93 %).

General procedure for dihydrothiolation (of monohydrothiolation) of alkynes

To a reaction flask containing a stir bar, alkyne (1 mmol), thiol (2.4 mmol, or 1.2 mmol for monohydrothiolation), deionized water (0.5 mL, 0.25 mL for monohydrothiolation) was added, and the resulting mixture was sealed and stirred for 24 h (12 h for monohydrothiolation) at the indicated temperature. Then the mixture was subjected to silica gel column chromatography using gradient elution from hexane to a mixture of hexane: ethyl acetate (10:1) to get final pure product **2-1-3** (or **2-1-4**).

1,2-Bis(p-tolylsulfanyl)hexane (**2-1-3a**)

$C_{20}H_{26}S_2$ (330.2): Calcd. C 72.67, H 7.93; Found C 72.80, H 8.25.

FTIR (neat) / cm^{-1} : 2955, 2925, 1491 and 804;

1H NMR (500 MHz; $CDCl_3$): δ 0.93 (t, $J=7.0$ Hz, 3H), 1.29-1.39 (m, 2H), 1.43-1.46 (m, 1H), 1.53-1.59 (m, 2H), 1.95-1.98 (m, 1H), 2.34 (s, 3H), 2.36 (s, 3H), 2.86 (dd, $J=13.25$, 10Hz, 1H), 3.05-3.10 (m, 1H), 3.23 (dd, $J=13.75$, 4.0 Hz, 1H), 7.04 (d, $J=8.0$ Hz, 2H), 7.09 (d, $J=7.5$ Hz, 2H), 7.13 (d, $J=7.5$ Hz, 2H), 7.25 (d, $J=8.0$ Hz, 2H).

^{13}C NMR ($CDCl_3$, 125 MHz): δ 14.29, 21.31, 21.41, 22.79, 29.22, 32.44, 40.11, 48.80, 129.92, 129.94, 130.52, 130.68, 132.45, 133.39, 136.51, 137.60.

1,2-Bis(phenylsulfanyl)hexane (**2-1-3b**)^{12b}

$C_{18}H_{22}S_2$ (302.1): Calcd. C 71.47, H 7.33; Found C 71.87, H 7.62.

FTIR (neat) / cm^{-1} : 2955, 2928, 1478, 1437, 738 and 690.

1H NMR ($CDCl_3$, 500 MHz): δ 0.83 (t, $J=7.5Hz$, 3H), 1.84-1.25 (m, 2H), 1.31-1.38 (m, 1H), 1.43-1.50 (m, 2H), 1.87-1.93 (m, 1H), 2.81 (dd, $J=13.5, 9.5Hz$, 1H), 3.02-3.15 (m, 1H), 3.17 (dd, $J=13.5, 4.0Hz$, 1H), 7.09-7.25 (m, 10H).

^{13}C NMR ($CDCl_3$, 125 MHz): δ 14.27, 22.77, 29.21, 32.51, 39.59, 48.54, 126.49, 127.47, 129.18, 129.23, 129.98, 132.78, 134.59, 136.10.

2-Bis(4-chlorophenylsulfanyl)hexane (**2-1-3c**)

$C_{18}H_{20}Cl_2S_2$ (370.0): Calcd. C 58.21, H, 5.43; Found C 58.50, H 5.36.

FTIR (neat) / cm^{-1} : 2956, 2928, 1474, 1095, 1011, 817.

1H NMR ($CDCl_3$, 500 MHz): δ 0.92(t, $J=7.0Hz$, 3H), 1.27-1.35 (m, 2H), 1.39-1.42 (m, 1H), 1.53-1.55 (m, 2H), 1.96-2.00(m, 1H), 2.89 (dd, $J=13.0, 9.0Hz$, 1H), 3.01-3.05 (m, 1H), 3.15 (dd, $J=13.0, 4.0Hz$, 1H), 7.12 (d, $J=8.5Hz$, 2H), 7.20 (d, $J=8.0Hz$, 2H), 7.25-7.27 (m, 4H)

^{13}C NMR ($CDCl_3$, 125 MHz): δ 13.90, 22.41, 28.90, 32.34, 39.79, 48.82, 129.05, 129.10, 131.27, 132.49, 132.74, 133.57, 133.96, 134.28.

1-Methoxy-2,3-bis(p-tolylsulfanyl)propane (**2-1-3d**)

C₁₈H₂₂OS₂ (318.1): Calcd. C 67.88, H, 6.96; Found C 68.29, H 7.16.

FTIR (neat) / cm⁻¹: 2920, 2886, 1491, 1119 and 805.

¹H NMR (CDCl₃, 500 MHz): δ 2.33 (s, 3H), 2.36 (s, 3H), 3.12 (dd, *J*= 13.75, 5.0Hz, 1H), 3.19(dd, *J*= 13.5, 8.5Hz, 1H), 3.24-3.28(m, 1H), 3.34 (s, 3H), 3.62 (dd, *J*= 10.0, 4.5Hz, 1H), 3.70 (dd, *J*= 9.5, 5.0Hz, 1H), 7.05 (d, *J*=8.0Hz, 2H), 7.09 (d, *J*=8.0Hz, 2H), 7.16 (d, *J*=8.0Hz, 2H), 7.29 (d, *J*=8.0Hz, 2H).

¹³C NMR (CDCl₃, 125 MHz): δ 20.96, 21.07, 35.85, 48.44, 58.96, 72.54, 129.65, 129.71, 129.94, 130.07, 132.01, 133.16, 136.25, 137.59.

4,5-Bis(phenylsulfanyl)pentan-1-ol (**2-1-3e**)

C₁₇H₂₀OS₂ (304.1): Calcd. C 67.06, H, 6.62; Found C 67.25, H 6.67.

FTIR (neat) / cm⁻¹: 3364, 2937, 1478, 1437, 1064, 1024, 739 and 691.

¹H NMR (CDCl₃, 500 MHz): δ 1.64-1.67 (m, 1H), 1.71-1.77 (m, 1H), 1.87-1.93 (m, 1H), 2.08-2.14 (m, 1H), 2.91 (dd, *J*= 13.75, 9.5 Hz, 1H), 3.14-3.19 (m, 1H), 3.30 (dd, *J*= 13.5, 4.0Hz, 1H), 3.67 (t, *J*=6.0Hz, 2H), 7.19-7.27 (m, 6H), 7.27-7.29 (m, 2H), 7.34-7.36 (m, 2H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 28.83, 29.87, 39.35, 48.07, 62.43, 126.29, 127.32, 128.91, 128.95, 129.72, 132.58, 133.92, 135.53.

2,3-Bis(4-chlorophenylsulfanyl)propan-1-ol (**2-1-3f**)

$\text{C}_{15}\text{H}_{14}\text{Cl}_2\text{OS}_2$ (343.9): Calcd. C 52.17, H 4.09; Found C 52.07, H 4.09.

FTIR (neat) / cm^{-1} : 3408, 2930, 2873, 1474, 1388, 1093 and 814.

^1H NMR (CDCl_3 , 500 MHz): δ 3.08-3.14 (m, 2H), 3.16-3.23 (m, 1H), 3.76-3.84 (m, 1H), 3.84-3.86 (m, 1H), 7.19-7.32 (m, 8H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 35.70, 51.51, 62.44, 129.17, 129.30, 131.07, 131.16, 132.69, 133.67, 134.24, 134.33.

2,3-Bis(4-chlorophenylsulfanyl)hexane (**2-1-3g**) Diastereomers **1**

$\text{C}_{18}\text{H}_{20}\text{Cl}_2\text{S}_2$ (370.0) Calcd. C 58.21, H 5.43; Found C 58.46, H 5.47.

FTIR (neat) / cm^{-1} : 2958, 2929, 1474, 1094, 1012 and 819.

^1H NMR (CDCl_3 , 500 MHz): δ 0.94 (t, $J=7.5\text{Hz}$, 3H), 1.33 (d, $J=7.0\text{Hz}$, 3H), 1.36-1.45 (m, 2H), 1.68-1.72 (m, 1H), 1.95-1.98(m, 1H), 3.02 (d, $J=10.5\text{Hz}$, 1H), 3.31 (m, 1H), 7.08 (d, $J=7.5\text{Hz}$, 2H), 7.10(d, $J=7.0\text{Hz}$, 2H), 7.19 (d, $J=8.5\text{Hz}$, 4H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 13.82, 14.11, 21.11, 29.74, 47.41, 53.05, 129.04, 133.09, 133.38, 133.51, 133.98.

2,3-Bis(4-chlorophenylsulfanyl)hexane (**2-1-3g**) Diastereomers **2**

C₁₈H₂₀Cl₂S₂ (370.0) Calcd. C 58.21, H 5.43; Found C 58.74, H 5.54.

FTIR (neat) / cm⁻¹: 2958, 2929, 1474, 1094, 1011 and 819.

¹H NMR (CDCl₃, 500 MHz): δ 0.89 (t, *J*=7.5Hz, 3H), 1.36 (d, *J*=7.0Hz, 3H), 1.40-1.44 (m, 2H), 1.58-1.62 (m, 1H), 1.67-1.74 (m, 1H), 3.20 (dt, *J*=9.0, 4.5Hz, 1H), 3.45 (dt, *J*=11.0, 4.5Hz, 1H), 7.23-7.27(m, 6H), 7.32 (d, *J*=8.0Hz, 2H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.94, 18.03, 20.87, 35.79, 49.16, 56.12, 129.00, 129.04, 132.93, 133.10, 133.28, 133.87, 134.79.

1,2-Bis(butylsulfanyl)hexane (**2-1-3h**)

C₁₄H₃₀S₂ (262.1) Calcd. C 64.05, H 11.52; Found C 64.66, H 11.68.

FTIR (neat) / cm⁻¹: 2956, 2928, 2859 and 1457.

¹H NMR (CDCl₃, 500 MHz): δ 0.90-0.93 (m, 9H), 1.26-1.49 (m, 9H), 1.54-1.59 (m, 4H), 1.78-1.82 (m, 1H), 2.52-2.55 (m, 4H), 2.65 (dd, *J*=12.25, 9.0Hz, 1H), 2.69-2.74 (m, 1H), 2.84 (dd, *J*=12.5, 4.5Hz, 1H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.67, 13.99, 21.97, 22.08, 22.58, 28.88, 30.51, 31.85, 31.92, 32.74, 33.19, 38.39, 45.71.

1,2-Bis(pentylsulfanyl)decane (**2-1-3i**)

C₂₀H₄₂S₂ (346.2) Calcd. C 69.29, H 12.21; Found C 69.44, H 12.38.

FTIR (neat) / cm⁻¹: 2954, 2934, 2854 and 1457. ¹H NMR (CDCl₃, 500 MHz): δ 0.87-0.92 (m, 9H), 1.29-1.38 (m, 19H), 1.45-1.51 (m, 2H), 1.57-1.59 (m, 4H), 1.78-1.82 (m, 1H), 2.51-2.54 (m, 4H), 2.65 (dd, *J*=12.75, 8.5Hz, 1H), 2.70-2.73 (m, 1H), 2.84 (dd, *J*=12.5, 4.5Hz, 1H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.97, 14.10, 22.30, 22.65, 26.69, 29.27, 29.47, 29.53, 30.81, 31.05, 31.16, 31.85, 33.04, 33.46, 38.37, 45.71.

4,5-Bis(pentylsulfanyl)pentan-1-ol (**2-1-3j**)

C₁₅H₃₂OS₂ (292.1) Calcd. C 61.58, H 11.03; Found C 61.61, H 11.13.

FTIR (neat)/ cm⁻¹: 3364, 2953, 2926, 2857, 1456 and 1058.

¹H NMR (CDCl₃, 500 MHz): δ 0.90 (t, *J*=7.0Hz, 6H), 1.29-1.40 (m, 8H), 1.53-1.62 (m, 5H), 1.66-1.74 (m, 1H), 1.76-1.82 (m, 1H), 1.84-1.96 (m, 1H), 2.52-2.55 (m, 4H), 2.65 (dd, *J*=12.75, 9.5Hz, 1H), 2.72-2.77 (m, 1H), 2.87(dd, *J*=12.75, 4.5Hz, 1H), 3.68 (brs, 2H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.92, 22.26, 24.07, 29.42, 29.49, 29.64, 29.86, 30.87, 31.01, 31.11, 33.01, 38.43, 45.44, 62.61.

(*E*)-2-(4-Chlorophenylsulfanyl)but-2-en-1-ol (**2-1-4a**)

C₁₀H₁₁ClOS (214.0): Calcd. C 56.00, H 5.12; Found C 55.94, H 5.16.

FTIR (neat) / cm⁻¹: 3355, 2912, 2854, 1474, 1388, 1079, 1011 and 816.

^1H NMR (CDCl_3 , 500 MHz): δ 1.94 (d, $J=7.0\text{Hz}$, 3H), 4.10 (s, 2H), 6.38 (q, $J=6.5\text{ Hz}$, 1H), 7.20-7.27 (m, 4H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 15.31, 65.99, 129.11, 130.04, 132.09, 132.96, 134.02, 135.91.

(*E*)-2-(4-Chloro-phenylsulfanyl)hex-2-en-1-ol (**2-1-4b**)

$\text{C}_{12}\text{H}_{15}\text{ClOS}$ (242.0) Calcd. C 59.37, H 6.23; Found C 59.50, H 6.25.

FTIR (neat) / cm^{-1} : 3354, 2958, 2929, 2869, 1474, 1092, 1011 and 815.

^1H NMR (CDCl_3 , 500 MHz): δ 0.95 (t, $J=7.5\text{Hz}$, 3H), 1.44-1.49 (m, 2H), 2.36 (q, $J=7.0\text{Hz}$, 2H), 4.08 (d, $J=5.0\text{Hz}$, 2H), 6.29 (t, $J=7.0\text{Hz}$, 1H), 7.20-7.26 (m, 4H). ^{13}C NMR (CDCl_3 , 125 MHz): δ 13.81, 22.26, 31.55, 65.78, 129.09, 130.12, 132.02, 133.29, 139.17, 141.55.

(*E*)-2-(Phenylsulfanyl)hex-2-en-1-ol (**2-1-4c**)

$\text{C}_{12}\text{H}_{16}\text{OS}$ (208.0) Calcd. C 69.19, H 7.74; Found C 69.22, H 7.80.

FTIR (neat) / cm^{-1} : 3351, 2957, 2929, 2869, 1582, 1476, 1438, 1086, 997 and 740.

^1H NMR (CDCl_3 , 500 MHz): δ 0.96 (t, $J=7.5\text{Hz}$, 3H), 1.46-1.52 (m, 2H), 2.39 (q, $J=7.5\text{Hz}$, 2H), 4.09 (d, $J=5.5\text{Hz}$, 2H), 6.29 (t, $J=7.0\text{Hz}$, 1H), 7.19-7.21 (m, 1H), 7.28-7.31 (m, 4H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 13.83, 22.32, 31.55, 65.85, 126.19, 128.99, 132.42, 134.56, 138.57, 140.97.

(*E*)-2-(*p*-tolylsulfanyl)hex-2-en-1-ol (**2-1-4d**)

C₁₃H₁₈OS (222.1) Calcd. C 70.22, H 8.16; Found C 70.18, H 8.10.

FTIR (neat) / cm⁻¹: 3380, 2957, 2927, 2869, 1491, 1455, 1086, 997 and 746.

¹H NMR (CDCl₃, 500 MHz): δ 0.96 (t, *J*=7.5Hz, 3H), 1.46-1.50 (m, 2H), 2.33 (s, 3H), 2.39 (q, *J*=7.0Hz, 2H), 4.06 (s, 2H), 6.20 (t, *J*=7.0Hz, 1H), 7.10 (d, *J*=8.0Hz, 2H), 7.21 (d, *J*=8.5Hz, 2H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.83, 20.96, 22.36, 31.49, 65.73, 129.79, 129.84, 130.54, 133.10, 136.47, 137.38.

(*E*)-2-(phenylsulfanyl)hept-2-en-1-ol (**2-1-4e**)

C₁₃H₁₈OS (222.1) Found: C, 70.22, H, 8.16. Calcd. for: C, 70.06, H, 8.30.

FTIR (neat)/ cm⁻¹: 3348, 2955, 2926, 2857, 1476, 1438, 1087, 1008 and 739.

¹H NMR (CDCl₃, 500 MHz): δ 0.92 (t, *J*=7.5Hz, 3H), 1.34-1.45 (m, 4H), 2.40 (q, *J*=7.0Hz, 2H), 4.09 (d, *J*=5.0Hz, 2H), 6.28 (t, *J*=7.0Hz, 1H), 7.19-7.21 (m, 1H), 7.27-7.31 (m, 4H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.86, 22.31, 29.21, 31.18, 65.82, 126.17, 128.97, 129.02, 132.21, 134.58, 138.73.

(*E*)-4-(4-chlorophenylsulfanyl)hex-4-en-3-ol (**2-1-4f**)

C₁₂H₁₅ClOS(242.0).

FTIR (neat) / cm⁻¹: 3388, 2963, 2931, 2874, 1473, 1388, 1091, 1011 and 816.

¹H NMR (CDCl₃, 500 MHz): δ 0.90 (t, *J*=7.5Hz, 3H), 1.58-1.69 (m, 2H), 1.86 (d, *J*=6.5Hz, 3H), 4.07(q, *J*=6.5Hz, 1H), 6.41 (q, *J*=7.0Hz, 1H), 7.18-7.23 (m, 4H).

¹³C NMR (CDCl₃, 125 MHz): δ 10.01, 15.51, 29.09, 77.38, 128.78, 128.99, 131.34, 134.63, 134.83, 135.88.

(*E*)-4-(Phenylsulfanyl)hex-4-en-3-ol (**2-1-4g**)

C₁₂H₁₆OS (208.0) Calcd. C 69.19, H 7.74; Found C 69.36, H 8.02.

FTIR (neat) / cm⁻¹: 3405, 2963, 2932, 2874, 1477, 1438, 1024, and 739.

¹H NMR (CDCl₃, 500 MHz): δ 0.91 (t, *J*=7.5Hz, 3H), 1.61-1.72 (m, 2H), 1.89(d, *J*=6.5Hz, 3H), 1.94 (d, *J*=6.5Hz, 1H), 4.08 (q, *J*=6.5Hz, 1H), 6.41 (q, *J*=6.5Hz, 1H), 7.31-7.17 (m, 1H), 7.24-7.29 (m, 4H).

¹³C NMR (CDCl₃, 125 MHz): δ 10.03, 15.51, 29.07, 77.35, 125.51, 127.61, 128.91, 134.26, 135.92, 136.06.

(*E*)-3-(4-chlorophenylsulfanyl)hex-3-en-2-ol (**2-1-4h**)

C₁₂H₁₅ClOS (242.0) Calcd. C 59.37, H 6.23; Found C 60.05, H 6.66.

FTIR (neat) / cm⁻¹: 3365, 2967, 2929, 2872, 1474, 1092, 1011 and 815.

¹H NMR (CDCl₃, 500 MHz): δ 1.01 (t, *J*=8.0Hz, 3H), 1.35 (d, *J*=6.0Hz, 3H), 1.98 (s, 1H), 2.28-2.33 (m, 2H), 4.29-4.32 (m, 1H), 6.35 (t, *J*=7.5Hz, 1H), 7.18-7.24 (m, 4H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 13.39, 22.61, 23.17, 71.32, 128.91, 129.02, 131.47, 134.70, 135.53, 140.66.

(*E*)-3-(*p*-tolylsulfanyl)hex-3-en-2-ol (**2-1-4i**)

$\text{C}_{13}\text{H}_{18}\text{OS}$ (222.1) Calcd. C 70.22, H 8.16; Found C 70.35, H, 8.30.

FTIR (neat) / cm^{-1} : 3378, 2966, 2928, 2871, 1491, 1118, 1074 and 805.

^1H NMR (CDCl_3 , 500 MHz): δ 1.02 (t, $J=7.5\text{Hz}$, 3H), 1.34 (d, $J=6.5\text{Hz}$, 3H), 2.03 (d, $J=5.5\text{Hz}$, 1H) 2.31 (s, 3H), 2.33-2.37 (m, 2H), 4.28-4.30 (m, 1H), 6.28 (t, $J=7.5\text{Hz}$, 1H), 7.08 (d, $J=8.0\text{Hz}$, 2H), 7.18 (d, $J=8.0\text{Hz}$, 2H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 13.51, 20.91, 22.48, 23.11, 70.98, 128.38, 129.73, 132.08, 135.72, 136.35, 139.09.

(*E*)-1-(1-phenylsulfanylpropenyl)cyclohexanol (**2-1-4j**)

$\text{C}_{15}\text{H}_{20}\text{OS}$ (248.1) Calcd. C 72.53, H 8.12; Found C 72.77, H, 8.33.

FTIR (neat) / cm^{-1} : 3402, 2931, 2853, 1477, 1438 and 737.

^1H NMR (CDCl_3 , 500 MHz): δ 1.56-1.75 (m, 10H), 1.81 (d, $J=6.5\text{Hz}$, 3H), 2.00 (s, 1H), 6.56 (q, $J=7.0\text{Hz}$, 1H), 7.09-7.11 (m, 1H), 7.20-7.25 (m, 4H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 16.21, 21.97, 25.51, 36.32, 74.92, 124.84, 126.19, 128.81, 132.90, 136.90, 140.91.

(*E*)-3-(4-chlorophenylsulfanyl)2-methyl-oct-3-en-2-ol (**2-1-4k**)

White crystal, mp 44-46 °C (from hexane and ethyl acetate). C₁₅H₂₁ClOS (284.1)

Calcd. C 63.25, H 7.43; Found C 63.38, H, 7.56.

FTIR (neat) / cm⁻¹: 3419, 2958, 2927, 2858, 1474, 1091 and 814.

¹H NMR (CDCl₃, 500 MHz): δ 0.86 (t, *J*=7.5Hz, 3H), 1.26-1.29 (m, 2H), 1.32-1.36 (m, 2H), 1.43 (s, 6H), 2.20-2.23 (m, 3H), 6.46 (t, *J*=7.0Hz, 1H), 7.14 (d, *J*=9.0Hz, 2H), 7.20 (d, *J*=9.0Hz, 2H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.85, 22.34, 29.21, 30.13, 30.84, 74.31, 127.59, 128.87, 130.75, 135.78, 138.83, 138.87.

(*E*)-3-(phenyl-2-*p*-tolylsulfanyl)prop-2-en-1-ol (**2-1-4l**)

C₁₆H₁₆OS (256.0) Calcd. C 74.96, H 6.29; Found C 74.96, H, 6.39.

FTIR (neat) / cm⁻¹: 3399, 2921, 2869, 1491, 1016, 808 and 759.

¹H NMR (CDCl₃, 500 MHz): δ 2.37 (s, 3H), 4.34(d, *J*=5.5Hz, 1H), 6.84 (s, 1H), 7.18 (d, *J*=7.5Hz, 2H), 7.29 (t, *J*=7.0Hz, 1H), 7.35-7.41 (m, 6H).

¹³C NMR (CDCl₃, 125 MHz): δ 21.11, 60.25, 127.53, 128.43, 128.61, 129.33, 130.13, 132.32, 133.71, 136.02, 137.61, 138.01.

(*E*)-1-(phenyl-2-phenylsulfanyl)hept-2-en-1-ol (**2-1-4m**)

C₁₉H₂₂OS (298.1) Calcd. C 76.46, H 7.43; Found C 76.70, H 7.64.

FTIR (neat) / cm⁻¹: 3365, 2956, 2931, 2870, 1454, 1002 and 759.

¹H NMR (CDCl₃, 500 MHz): δ 0.89 (t, *J*=7.5 Hz, 3H), 1.30-1.36 (m, 2H), 1.37-1.42 (m, 2H), 2.33-2.39 (m, 2H), 2.43 (d, *J*=5.0Hz, 1H), 5.22 (d, *J*=5.5 Hz, 1H), 6.39 (t, *J*=7.5Hz, 1H), 7.15-7.18 (m, 1H), 7.25-7.35 (m, 9H).

¹³C NMR (CDCl₃, 125 MHz): δ 13.89, 22.39, 29.48, 31.06, 76.75, 125.85, 126.68, 127.75, 128.29, 128.32, 128.94, 134.99, 135.57, 139.99, 141.67.

(*E*)-tert-butyl-dimethyl-(2-phenylsulfanyl-hex-2-enyloxy)silane (**2-1-5**)

¹H NMR (CDCl₃, 500 MHz): δ 0.02 (s, 6H), 0.90 (s, 9H), 0.94 (t, *J*=7.5Hz, 3H), 1.44-1.49 (m, 2H), 2.38 (q, *J*=7.5Hz, 2H), 4.13 (d, *J*=1.0Hz, 2H), 6.39 (t, *J*=7.5Hz, 1H), 7.15 (t, *J*=4.5Hz, 1H), 7.25 (d, *J*=4.0Hz, 4H).

¹³C NMR (CDCl₃, 125 MHz): δ -5.39, 13.86, 18.37, 22.50, 25.87, 31.31, 65.87, 125.58, 128.14, 128.83, 130.81, 135.77, 136.98.

(*E*)-2-benzenesulfonyl-hex-2-en-1-ol (**2-1-6**)

^1H NMR (CDCl_3 , 500 MHz): δ 0.85 (t, $J=7.0\text{Hz}$, 3H), 1.34-1.38 (m, 2H), 2.50 (q, $J=7.5\text{Hz}$, 2H), 4.33 (s, 2H), 6.34 (t, $J=7.5\text{Hz}$, 1H), 7.54 (t, $J=7.0\text{Hz}$, 2H), 7.62 (t, $J=7.5\text{Hz}$, 1H), 7.94 (d, $J=7.5\text{Hz}$, 2H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 13.56, 21.81, 30.29, 63.89, 127.20, 129.13, 133.39, 140.27, 141.34, 146.84.

2.2. Gold-Catalyzed Intramolecular Oxygen Transfer Reaction of 2-Alkynyl-1,5-diketones and 2-Alkynyl-5-Ketoesters

2.2.1 Background and Introduction

The Hammond group has been particularly interested in the interactions of allene and gold.^{2d, 33} This led to the finding that 2-alkynyl-1,5-diketones or 2-alkynyl-5-ketoesters can easily isomerized into 1,5-diketo cyclopentene in the presence of catalytic amounts of gold (I) chloride. This methodology makes available substituted cyclopentenones, which are core structures or precursors of important natural products (Figure 3).³⁴

Figure 3. Natural Product Containing the Cyclopentenone Core Structure

Isotopic experiments and theoretical calculations performed by members of the Hammond group supported the novel proposed intramolecular [4+2] cycloaddition of a gold-containing furanium intermediate to a carbonyl group, instead of the previous well-accepted [2+2] pathway.

2.2.2 Result and Discussion

From entry 1 to 3, aromatic rings at R¹ were examined, giving excellent yield of 2-2-**2a**, 2-2-**2b**, 2-2-**2c** at room temperature (Table 4).

Table 4. Isomerization of 2-Alkynyl-1,5-Diketones and 2-Alkynyl-5-Ketoesters

Reaction scheme showing the isomerization of 2-alkynyl-1,5-diketones (2-2-1) to 2-alkynyl-5-ketoesters (2-2-2) using AuCl (5-10 mol %) in CDCl₃ at room temperature.

Entry	Substrate	Product	(Yield %)
1			91%
2			93%
3			100%
4			91%
5			92 %
6			92 %
7			90 %
8 ^a			50 %
9			92 %

^a Yield on conversion.

[Note: My participation in this collective work from the Hammond's group was restricted to the isomerization of 2-alkynyl-1,5-diketones and 2-alkynyl-5-ketoesters].

Similarly, the same aromatic rings at R¹ worked equally well with the 2-alkynyl-5-ketoester system (entries 4 to 6). In entries 7 and 8, the benzyl group at R² was tested on a 2-alkynyl-5-ketoester system. While substrate **2-2-7a** gave excellent yield, **2-2-8a** was not suitable, affording moderate yields of the desired product, together with an unidentified side product. In the last entry, substrate **2-2-9a**, having two methyl groups at R³ and R⁴, still isomerized well using this methodology.

We proposed a very unique and unprecedented intramolecular [4+2] cycloaddition (Scheme 7) for this transformation, rather than the well-accepted [2+2] cycloaddition.

Scheme 7. ¹⁸O Isotopic Labeling Experiment for Mechanistic Studies

Mechanistic investigations on the cycloisomerization were carried out by means of both ^{18}O isotopic experiments and quantum chemical calculations. The results from both, the designed isotopic experiments and theoretical calculations (omitted in this thesis because they were done by Dr. Leping Liu and Professor Kenneth Houk at UCLA), satisfactorily supported the novel proposed intramolecular [4+2] cycloaddition of a gold-containing furanium intermediate to a carbonyl group, instead of the previously accepted [2+2] pathway.³⁵

2.2.3 Experimental Section

2-Alkynyl-1,5-diketones or 2-alkynyl-5-ketoesters were prepared according to our previously published paper.^{2d} Gold(I) chloride was purchased from Aldrich. 2-Alkynyl-1,5-diketone (or 2-alkynyl-5-ketoester) (0.02-0.15 mmol), gold(I) chloride (5-10 mol %) were dissolved in CDCl_3 (0.6 mL) in NMR tube and monitored constantly until the reaction was completed. The reaction mixture was then subjected to a short chromatography separation, eluted by a hexane: ethyl acetate system, to give the cyclized product. The products were characterized by ^1H , ^{13}C NMR, HRMS and IR spectra.

Typical procedure for preparation of 2-((4-methoxyphenyl)ethynyl)-2-methyl-1,5-diphenylpentane-1,5-dione (2-2-1a)

4-(4-methoxyphenyl)-2-methyl-1-phenylbuta-2,3-dien-1-one (528.0 mg, 2.0 mmol), 1-phenylprop-2-en-1-one (396 mg, 3.0 mmol) and TBAF (0.2 mmol) were dissolved in THF (3 mL) and stirred at room temperature for 5 h. The reaction mixture was quenched by saturated ammonium chloride (15 mL) and followed by diethyl ether extraction (25 mL $\times$ 3). The ether layers were combined and dried over anhydrous Na_2SO_4 , filtered and

evaporated under reduced pressure, the resulting residue was subjected to silica gel chromatography eluted by hexanes and ethyl acetate system to yield 2-((4-methoxyphenyl)ethynyl)-2-methyl-1,5-diphenylpentane-1,5-dione (**2-2-1a**) (475.0 mg, 60 %).

(5-Benzoyl-5-methyl-2-phenyl-cyclopent-1-enyl)-(4-methoxy-phenyl)-methanone (**2-2-2a**)

^1H NMR (CDCl_3 , 400MHz) δ 7.90 (d, J = 8.0Hz, 2H), 7.67 (d, J = 8.0Hz, 2H), 7.42 (t, J = 6.8Hz, 1H), 7.35 (t, J = 6.8Hz, 2H), 7.15 (brs, 2H), 7.09 (brs, 3H), 6.62 (d, J = 8.4 Hz, 2H), 3.70 (s, 3H), 3.02-3.29 (m, 1H), 2.95-3.01 (m, 1H), 2.61-2.69 (m, 1H), 2.17-2.23 (m, 1H), 1.66 (s, 3H).

^{13}C NMR (CDCl_3 , 100MHz) δ 203.4, 195.1, 162.9, 146.7, 141.1, 137.7, 135.7, 131.8, 131.3, 130.1, 128.6, 128.3, 128.2, 128.1, 113.1, 65.9, 55.2, 36.1, 35.7, 24.2.

HRMS m/z (ES^+) calcd for $\text{C}_{27}\text{H}_{25}\text{O}_3$ ($[\text{M}+\text{H}]^+$) 397.1798, found 397.1790, m/z (ES^+) calcd for $\text{C}_{27}\text{H}_{24}\text{O}_3\text{Na}$ ($[\text{M}+\text{Na}]^+$) 419.1618, found 419.1608.

FTIR (neat)/ cm^{-1} : 1675, 1637, 1596, 1508, 1334, 1255, 1162, 763, 698.

(5-Benzoyl-5-methyl-2-phenyl-cyclopent-1-enyl)-phenylmethanone (**2-2-2b**)

^1H NMR (CDCl_3 , 400MHz) δ 7.90 (d, $J = 7.2\text{Hz}$, 2H), 7.65 (d, $J = 7.6\text{Hz}$, 2H), 7.44 (t, $J = 7.2\text{Hz}$, 1H), 7.37 (t, $J = 7.6\text{Hz}$, 2H), 7.24 (t, $J = 7.6\text{Hz}$, 1H), 7.05-7.13 (m, 7H), 3.22-3.29 (m, 1H), 3.00-3.09 (m, 1H), 2.64-2.72 (m, 1H), 2.22-2.28 (m, 1H), 1.73 (s, 3H).

^{13}C NMR (CDCl_3 , 100MHz) δ 203.3, 196.5, 148.6, 141.2, 137.4, 137.3, 135.6, 132.1, 131.5, 129.5, 128.6, 128.4, 128.3, 128.1, 127.9, 127.7, 66.2, 36.4, 35.8, 24.3.

HRMS m/z (ES^+) calcd for $\text{C}_{26}\text{H}_{23}\text{O}_2$ ($[\text{M}+\text{H}]^+$) 367.1693, found 367.1687, m/z (ES^+) calcd for $\text{C}_{26}\text{H}_{22}\text{O}_2\text{Na}$ ($[\text{M}+\text{Na}]^+$) 389.1512, found 389.1505.

FTIR (neat)/ cm^{-1} : 1674, 1637, 1447, 1277, 1251, 968, 725, 693.

(5-Benzoyl-5-methyl-2-phenyl-cyclopent-1-enyl)-(4-chloro-phenyl)-methanone (**2-2-2c**)

^1H NMR (CDCl_3 , 400MHz) δ 7.88 (d, $J = 8.0\text{Hz}$, 2H), 7.60 (d, $J = 8.0\text{Hz}$, 2H), 7.46 (t, $J = 7.2\text{Hz}$, 1H), 7.38 (t, $J = 7.6\text{Hz}$, 2H), 7.10 (brs, 7H), 3.17-3.25 (m, 1H), 3.04-3.12 (m, 1H), 2.64-2.72 (m, 1H), 2.26-2.72 (m, 1H), 1.77 (s, 3H).

^{13}C NMR (CDCl_3 , 100MHz) δ 203.4, 195.4, 148.5, 140.8, 138.3, 136.8, 135.8, 135.4, 131.7, 130.8, 128.7, 128.6, 128.3, 128.2, 128.1, 128.0, 66.8, 36.3, 36.0, 24.6.

HRMS m/z (ES^+) calcd for $C_{26}H_{22}O_2Cl$ ($[M+H]^+$) 401.1303, found 401.1297, m/z (ES^+) calcd for $C_{26}H_{21}O_2NaCl$ ($[M+Na]^+$) 423.1122, found 423.1113.

FTIR (neat)/ cm^{-1} : 1674, 1637, 1576, 1275, 1251, 1089, 968, 747, 697.

2-Benzoyl-1-methyl-3-phenyl-cyclopent-2-enecarboxylic acid ethyl ester (**2-2-2d**)

1H NMR ($CDCl_3$, 400MHz) δ 7.65 (d, J = 8.0Hz, 2H), 7.26 (t, J = 7.6Hz, 1H), 7.09-7.13 (m, 4H), 7.03-7.06 (m, 3H), 3.93-4.05 (m, 2H), 3.10-3.18 (m, 1H), 2.92-2.98 (m, 1H), 2.54-2.59 (m, 1H), 2.00-2.06 (m, 1H), 1.62 (s, 3H), 1.00 (t, J = 7.2Hz, 3H).

^{13}C NMR ($CDCl_3$, 100MHz) δ 197.0, 175.7, 148.7, 139.4, 137.3, 135.6, 132.3, 129.4, 128.2, 128.2, 127.9, 127.8, 60.8, 59.9, 36.4, 35.7, 23.1, 13.8.

HRMS m/z (ES^+) calcd for $C_{22}H_{23}O_3$ ($[M+H]^+$) 335.1642, found 335.1637, m/z (ES^+) calcd for $C_{22}H_{22}O_3Na$ ($[M+Na]^+$) 357.1461, found 357.1456, m/z (ES^+) calcd for $C_{19}H_{17}O$ ($[M-COOEt]^+$) 261.1279, found 261.1272.

FTIR (neat)/ cm^{-1} : 1731, 1646, 1447, 1279, 1254, 1173, 1093, 696.

2-(4-Methoxy-benzoyl)-1-methyl-3-phenyl-cyclopent-2-enecarboxylic acid ethyl ester (**2-2e**)

^1H NMR (CDCl_3 , 400MHz) δ 7.67 (d, $J = 8.8\text{Hz}$, 2H), 7.14 (brs, 2H), 7.07 (brs, 3H), 6.64 (d, $J = 8.4\text{Hz}$, 2H), 3.91-4.05 (m, 2H), 3.72 (s, 3H), 3.10-3.18 (m, 1H), 2.87-2.94 (m, 1H), 2.54-2.61 (m, 1H), 1.97-2.04 (m, 1H), 1.58 (s, 3H), 1.00 (t, $J = 7.2\text{Hz}$, 3H).

^{13}C NMR (CDCl_3 , 100MHz) δ 195.7, 175.7, 163.0, 146.8, 139.2, 135.7, 131.8, 130.3, 128.2, 128.1, 127.9, 113.1, 60.8, 60.0, 55.3, 36.3, 35.4, 23.2, 13.8.

HRMS m/z (ES^+) calcd for $\text{C}_{23}\text{H}_{25}\text{O}_4$ ($[\text{M}+\text{H}]^+$) 365.1747, found 365.1741, m/z (ES^+) calcd for $\text{C}_{23}\text{H}_{24}\text{O}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$) 387.1567, found 387.1560, m/z (ES^+) calcd for $\text{C}_{20}\text{H}_{19}\text{O}_2$ ($[\text{M}-\text{COOEt}]^+$) 291.1385, found 291.1377.

FTIR (neat)/ cm^{-1} : 1729, 1637, 1597, 1507, 1256, 1162, 1093, 1027, 760, 697.

2-(4-Chloro-benzoyl)-1-methyl-3-phenyl-cyclopent-2-enecarboxylic acid ethyl ester (**2-2-f**)

^1H NMR (CDCl_3 , 400MHz) δ 7.59 (d, $J = 7.6\text{Hz}$, 2H), 7.08 (brs, 7H), 3.95-4.07 (m, 2H), 3.10-3.17 (m, 1H), 2.88-3.10 (m, 1H), 2.51-2.58 (m, 1H), 2.00-2.06 (m, 1H), 1.62 (s, 3H), 1.02 (t, $J = 6.8\text{Hz}$, 3H).

^{13}C NMR (CDCl_3 , 125 MHz) δ 195.7, 175.6, 149.2, 139.1, 138.5, 135.8, 135.4, 130.8, 128.6, 128.2, 128.1, 128.1, 60.8, 60.1, 36.5, 35.7, 23.2, 13.9.

HRMS m/z (ES^+) calcd for $C_{22}H_{22}O_3Cl$ ($[M+H]^+$) 369.1252, found 369.1248, m/z (ES^+) calcd for $C_{22}H_{21}O_3ClNa$ ($[M+Na]^+$) 391.1071, found 391.1064, m/z (ES^+) calcd for $C_{19}H_{16}OCl$ ($[M-COOEt]^+$) 295.0890, found 295.0881.

FTIR (neat)/ cm^{-1} : 1732, 1646, 1278, 1253, 1166, 1091, 860, 755, 697.

Ethyl 2-benzoyl-1-benzyl-3-phenylcyclopent-2-enecarboxylate (**2-2-2g**)

1H NMR ($CDCl_3$, 400MHz) δ 7.68 (d, $J = 7.2Hz$, 2H), 7.38 (d, $J = 6.8Hz$, 2H), 7.28 (t, $J = 7.6Hz$, 1H), 7.13-7.25 (m, 6H), 7.01 (brs, 4H), 3.88-4.03 (m, 2H), 3.70 (d, $J = 13.6Hz$, 1H), 3.33 (d, $J = 13.6Hz$, 1H), 2.75-2.83 (m, 1H), 2.46-2.54 (m, 1H), 2.27-2.32 (m, 1H), 1.77-1.86 (m, 1H), 0.94 (t, $J = 6.8Hz$, 3H).

^{13}C NMR ($CDCl_3$, 100MHz) δ 197.8, 175.1, 151.1, 137.5, 136.2, 135.6, 132.2, 130.6, 129.5, 128.2, 128.0, 127.8, 127.8, 127.7, 126.5, 65.3, 61.0, 42.3, 35.7, 32.7, 13.7.

HRMS m/z (ES^+) calcd for $C_{28}H_{27}O_3$ ($[M+H]^+$) 411.1955, found 411.1949, m/z (ES^+) calcd for $C_{28}H_{26}O_3Na$ ($[M+Na]^+$) 433.1774, found 433.1762, (ES^+) calcd for $C_{25}H_{21}O$ ($[M-COOEt]^+$) 337.1592, found 337.1583.

FTIR (neat)/ cm^{-1} : 1727, 1642, 1447, 1278, 1247, 1175, 697.

2-Benzoyl-1-benzyl-3-methyl-cyclopent-2-enecarboxylic acid ethyl ester (**2-2-2h**)

^1H NMR (CDCl_3 , 500 MHz) δ 7.85 (d, $J = 7.5\text{Hz}$, 2H), 7.55 (t, $J = 8.0\text{Hz}$, 1H), 7.45 (t, $J = 8.0\text{Hz}$, 2H), 7.25 (brs, 2H), 7.20 (brs, 3H), 4.00-4.08 (m, 2H), 3.64 (d, $J = 13.0\text{Hz}$, 1H), 3.24 (d, $J = 13.0\text{Hz}$, 1H), 2.18-2.27 (m, 1H), 2.11-2.16 (m, 2H), 1.36-1.40 (m, 4H), 1.04 (t, $J = 7.0\text{Hz}$, 3H).

^{13}C NMR (CDCl_3 , 125 MHz) δ 197.3, 175.7, 151.5, 139.3, 137.8, 135.7, 132.5, 130.6, 129.3, 128.3, 127.6, 126.2, 64.2, 60.8, 41.5, 37.7, 32.6, 16.8, 13.8.

HRMS m/z (ES^+) calcd for $\text{C}_{23}\text{H}_{25}\text{O}_3$ ($[\text{M}+\text{H}]^+$) 349.1798, found 349.1799, m/z (ES^+) calcd for $\text{C}_{23}\text{H}_{24}\text{O}_3\text{Na}$ ($[\text{M}+\text{Na}]^+$) 371.1618, found 371.1614, m/z (ES^+) calcd for $\text{C}_{20}\text{H}_{19}\text{O}$ ($[\text{M}-\text{COOEt}]^+$) 275.1436, found 275.1431.

FTIR (neat)/ cm^{-1} : 1728, 1643, 1447, 1280, 1238, 1172, 1074, 703.

1-(2-Benzoyl-1,3-dimethyl-cyclopent-2-enyl)-ethanone (**2-2-2i**)

^1H NMR (CDCl_3 , 400MHz) δ 7.75 (d, $J = 7.6\text{Hz}$, 2H), 7.52 (t, $J = 7.2\text{Hz}$, 1H), 7.42 (t, $J = 7.6\text{Hz}$, 2H), 2.61-2.66 (m, 2H), 2.09-2.17 (m, 4H), 1.78-1.83 (m, 1H), 1.56 (s, 3H), 1.38 (s, 3H).

^{13}C NMR (CDCl_3 , 100MHz) δ 210.7, 196.1, 150.1, 140.4, 139.2, 132.6, 129.0, 128.5, 64.8, 38.7, 34.9, 25.7, 22.0, 17.3.

HRMS m/z (ES^+) calcd for $C_{16}H_{19}O_2$ ($[M+H]^+$) 243.1380, found 243.1381, m/z (ES^+)
calcd for $C_{16}H_{18}O_2Na$ ($[M+Na]^+$) 265.1199, found 265.1199.

FTIR (neat)/ cm^{-1} : 1706, 1640, 1447, 1346, 1281, 739, 696.

3. EXPLORATION OF THE CHEMISTRY OF SELECTFLUOR

3.1 SelectfluorTM

Recently, a number of N-F fluorinating agents have been emerged as electrophilic fluorinating agents, which are easy to handle, safe and stable. They are either quaternary ammonium salt $R_3N^+F^-$ or R_2NF ,³⁶ in which SelectfluorTM (F-TEDA- BF_4) (Scheme 8) is one of cheapest and most reactive ones.³⁷ Selectfluor is commercially prepared by Air Products by following process shown in Scheme 8.³⁸

Scheme 8. Commercially Synthetic pathway for Selectfluor

There are many nice examples of fluorination by Selectfluor in literature.³⁹ Selectfluor can fluorinate electron rich aromatic systems⁴⁰, aliphatic hydrocarbon,⁴¹ aliphatic amines⁴², alkenes and their derivatives⁴³ and asymmetric fluorination.⁴⁴ In addition to fluorination, Selectfluor can be oxidant for functional groups transformation,⁴⁵ halogens (iodine),⁴⁶ C-H functionalization,⁴⁷ gold catalyzed oxidative C-C and C-heteroatom bond formation,⁴⁸ palladium catalyzed C-C and C-heteroatom bond formation.⁴⁹

In this section, Selectfluor will be explored as fluorinating agent to prepare fluoroalkyl (*E*) α,β -unsaturated ketones, which are very important intermediates for synthesis of bioactive agents. Then, combination of copper and selectfluor was found as a powerful oxidant, which readily oxidizes amide into imide at room temperature.

3.2. Stereoselective Synthesis of Fluoroalkyl α,β -Unsaturated Ketones

3.2.1 Background

Monofluoroalkyl α,β -unsaturated ketones **3-2-1**, such as **A**,⁵⁰ **B**⁵¹ and **C**,⁵² are important molecular synthons, as showcased by their use in the synthesis of numerous bioactive agents (Figure 4).⁵⁰⁻⁵³ Fluoroketone **3-2-1** is most commonly synthesized either by using a fluorine-containing building block or through the fluorination of an unsaturated ketone. But the above literature methodologies have limitations.

Figure 4. Monofluoroalkyl (*E*)- α,β -Unsaturated Ketones **A**, **B** and **C** are Important Building Blocks for Bioactive Agents

The building block approach relies on a shallow pool of fluorine containing moiety (Scheme 9, top),⁵⁰⁻⁵⁴ whose preparation is not always trivial; for example, the synthesis of fluorine-containing Wittig reagents is lengthy and inefficient (Scheme 10).^{54a, 54d, 55}

Scheme 9. Literature Construction of Monofluoroalkyl α,β -Unsaturated Ketones by Fluorine Containing Building Blocks (Top) and Fluorination of Specific Ketones (Bottom)

Scheme 10. Lengthy Preparation Pathway for a Fluorine-Containing Wittig Reagent

On the other hand, the fluorination approach⁵⁶ is hampered by the fact that regioselective fluorination of the starting unsaturated ketone is case specific (Scheme 9, Bottom). Owing to these limitations, medicinal chemists find it difficult to build libraries of fluoroalkyl α,β -unsaturated ketone. An efficient preparation of the said ketones is highly desirable.

3.2.2 Our Efficient Strategy to Prepare Ketone 3-2-1

Inspired by the works of Gouverneur⁵⁷ and Nevado⁵⁸ (Scheme 11, top), we envisioned an environmentally friendly and efficient fluorination process to overcome aforementioned drawbacks (Scheme 11, bottom). We posited that an electron rich 1,3-butadien-2-ol ester **3-2-2** could be a suitable substrate for electrophilic fluorination. The diene **3-2-2** is readily made by the gold-catalyzed isomerization of allenyl carbinol ester **3-2-3** using Gagosz's methodology.⁵⁹ Gagosz and co-workers had reported that terminal allenyl carbinol ester **3-2-3** isomerize to 1,3-butadien-2-ol ester **3-2-2** with moderate to good stereoselectivity in presence of gold catalyst **3-2-4**.

Scheme 11. Gold-catalyzed Isomerization of Alkyne and Allene and Following Fluorination Strategy

In this chapter we report that a stereomixture of **3-2-2** can be fluorinated at room temperature by Selectfluor to give fluoroalkyl α,β -unsaturated ketone **3-2-1** in high yields and with exclusive *E* stereoselectivity. All results are shown in Table 5.

Table 5. Preparation of α,β -Unsaturated Ketones **3-2-1** from *in Situ* Generated Dienes **3-2-2**.^a

Entry	Substrate (1)	Dienes (2)	Ketone (3), Yield ^b
1	 3-2-3a	 3-2-2a (<i>E:Z</i> =95:5)	 3-2-1a , 90% (<i>E:Z</i> =99:1)
2	 3-2-3b	 3-2-2b (<i>E:Z</i> =95:5)	 3-2-1a , 95% (<i>E:Z</i> =99:1)
3	 3-2-3c	 3-2-2c (<i>E:Z</i> =95:5)	 3-2-1a , 96% (<i>E:Z</i> =99:1)
4	 3-2-3d	 3-2-2d (<i>E:Z</i> =90:10)	 3-2-1a , 95% (<i>E:Z</i> =99:1)
5	 3-2-3e	 3-2-2e (<i>E:Z</i> =95:5)	 3-2-1e , 95% (<i>E:Z</i> =99:1) ^c
6	 3-2-3f	 3-2-2f (<i>E:Z</i> =87:13)	 3-2-1e , 80% (<i>E:Z</i> =99:1) ^c
7	 3-2-3g	 3-2-2g (<i>E:Z</i> =95:5)	 3-2-1g , 83% (<i>E:Z</i> =99:1) ^d
8	 3-2-3h	 3-2-2h (1:0.41:0.68:0.87)	 3-2-1h , 72% (<i>E:Z</i> =99:1) ^d
9	 3-2-3i	 3-2-2i (<i>E:Z</i> =87:13)	 3-2-1i , 79% (<i>E:Z</i> =99:1) ^d
10	 3-2-3j	 3-2-2j (<i>E:Z</i> =87:13)	 3-2-1i , 70% (<i>E:Z</i> =97:3) ^d

^a Reaction condition: allenyl carbinol ester (**3-2-3**) (0.25 M) and the gold catalyst **3-2-4** (1 mol %) in CDCl₃ and monitored by ¹H NMR until no progress. Then CDCl₃ was dried in vacuum, Selectfluor (1.2 equiv.) and acetonitrile (1.5 mL) were added and stirred for 3 h. ^b Yield was calculated on starting allenyl carbinol ester (**3-2-3**). ^c ¹⁹F NMR yield due to their volatility, 2,3,5-trifluorotoluene as internal standard. ^d Yield of allylic alcohol owing to their instability on long silica gel column. Allylic alcohols of **3-2-1g** and **3-2-1h** contain less than 3 % of corresponding saturated alcohol. d.r. of the allylic alcohol (from **3-2-1h**) is 4:1.

Carbinol ester **3-2-3a** (entry 1) isomerizes to diene **3-2-2a** in CDCl_3 in the presence of gold catalyst **3-2-4**,⁶⁰ and diene **3-2-2a** reacts readily with Selectfluor in MeCN to yield the corresponding fluorinated ketone **3-2-1a** in excellent yield and with exclusive *E* stereoselectivity.⁶¹ It is noteworthy that dienes **3-2-2a** (*E*:*Z* = 95:5), during fluorination, yielded exclusively the *E* isomer. There are two possibilities to produce exclusively fluoroalkyl (*E*)- α,β -unsaturated ketone **3-2-1a**. First, the (*Z*)-diene **3-2-2a** may be isomerized, upon fluorination, into *E* stereomer and give *E* ketone **3-2-1a** after subsequent hydrolysis. Secondly, (*E*)-diene **3-2-2a** is probably more reactive, than (*Z*)-diene **3-2-2a**, with Selectfluor and no fluorine cation-enabled isomerization happened. The isomerization is proposed as main pathway to give exclusively *E* ketone, since unreacted (*E*)-diene **3-2-2a** and corresponding side products were failed to be isolated. Similarly, aromatic substrates **3-2-3b**, **3-2-3c** and **3-2-3d** isomerized to the corresponding dienes **3-2-2** as a mixture of *E*/*Z* isomers, but upon fluorinations, these isomers afforded **3-2-1a** in high stereoselectivity and excellent chemical yields, regardless of the type of carboxylic acid derivatives used. Aliphatic allenyl esters **3-2-5e** (entry 5) and **3-2-6f** (entry 6) were also examined. Both of them efficiently isomerized and were fluorinated to give **3-2-3e** (*E*:*Z* = 99:1) in good yield.⁶² The high molecular weight aliphatic allenyl ester **3-2-3g** (entry 7) was tested in order to isolate the desired pure product (not volatile). Because all substrates used in Gagosz's work were terminal allenyl esters, we were intrigued as to how would an internal allenyl ester such **3-2-3h** behave under our reaction conditions (entry 8). We were pleasantly surprised to discover that even though **3-2-3h** isomerized readily to give a complex mixture of stereoisomers **3-2-2h** (1:0.41:0.68:0.87), this mixture produced the desired **3-2-1h** in good yield as a single *E*-isomer. Benzoate

and acetate derivatives of 1,1-disubstituted allenyl esters **3-2-3i** and **3-2-3j** were also tested (entries 9 and 10). Their isomerizations furnished dienes **3-2-2i** and **3-2-2j**, respectively, but after fluorination, only **3-2-1i** was obtained in very good yield and with exclusive *E* selectivity.

We also investigated the possibility of preparing fluoroalkyl *E,E*- $\alpha,\beta,\gamma,\delta$ -unsaturated ketone **3-2-1k** using our protocol, but, unfortunately, this attempt failed because no isomerization of the two substrates tested, **3-2-3k** and **3-2-3l**, took place in the presence of gold catalyst **3-2-4** (Scheme 12).

Scheme 12. Limitations of Our Protocol

A plausible mechanism for the isomerization of a diene mixture is proposed in Scheme 13. The electron-rich double bond of diene **3-2-2** attacks the electrophilic fluorine in Selectfluor to form cationic intermediate **C**, which instantly isomerizes to form the more stable intermediate **D**, and after hydrolysis with trace amounts of water present in the reaction media, yields fluoroalkyl *E*- α,β -unsaturated ketone **3-2-1**.

Scheme 13. Proposed Mechanism for Fluorinated Cation Enabled Isomerization

The reported electrophilic fluorination-nucleophilic addition reactions of glycals⁶³ and other substrates⁵⁷ using Selectfluor lend support to our proposal. It is likely that in our case an electrophilic fluorinated cation-enabled isomerization occurred to form the most stable stereoisomer before nucleophilic attack of trace water from the reaction mixture happened.

3.2.3 Summary

Our protocol can efficiently fluorinate a mixture of dienes **3-2-2** under mild conditions to give fluoroalkyl *E*- α,β -unsaturated ketone **3-2-1** in good to excellent yield. Because all starting allenyl alcohols can be readily prepared using literature methodologies⁶⁴ our methodology can be used to prepare a large variety of fluoroalkyl *E*- α,β -unsaturated ketones. These ketones could be further functionalized; a case in point is the conversion to a fluorosugar precursor⁶⁵ through dihydroxylation of a double bond.^{54a}

3.2.4 Experimental Section

General

The gold complex (**3-2-4**) and Selectfluor were purchased from Aldrich. All air and/or moisture sensitive reactions were carried out under argon atmosphere. Solvents (tetrahydrofuran, ether, dichloromethane and DMF) were chemically dried using a commercial solvent purification system. All other reagents and solvents were employed without further purification. The products were purified using a Combiflash system or Chromatotron apparatus or a regular glass column. TLC was developed on Merck silica gel 60 F254 aluminum sheets. ^1H , ^{13}C and ^{19}F NMR spectra were recorded at 500 (or 400), 126 (or 100) and 470 (or 376) MHz respectively, using CDCl_3 (or CD_3CN) as a solvent. The chemical shifts are reported in δ (ppm) values relative to CHCl_3 (7.26 ppm for ^1H NMR and 77.0 ppm for ^{13}C NMR) and CFCl_3 (0 ppm for ^{19}F NMR), multiplicities are indicated by s (singlet), d (doublet), t (triplet), q (quartet), p (pentet), h (hextet), m (multiplet) and br (broad). Coupling constants, J , are reported in Hertz. RTIR spectra were recorded in ATR (attenuated total reflection) solid mode using a Perkin Elmer Spectrum 100.

Experimental procedure

1. Preparation of allenyl alcohol

Allenyl alcohol precursors of **3-2-3a**, **3-2-3e** and **3-2-3g** were prepared by Crabbe reaction.^{64a}

1-Phenylprop-2-yn-1-ol (396.0 mg, 3.0 mmol), paraformaldehyde (180.0 mg, 6.0 mmol), diisopropylamine (834.0 μ L, 6.0 mmol), CuBr (257.4 mg, 1.8 mmol) and 1,4-dioxane (20 mL) were added into 50 mL round bottom reaction flask containing stir bar, and the flask was equipped with reflux condenser. The reaction mixture was refluxed in oil bath for 2 h, cooled to room temperature and filtered through silica gel (about 15 g) and washed by mixture of hexanes and ethyl acetate (100 mL, ratio is 4:1). The solvent was removed in vacuum and the resulting residue was purified on silica gel column, which was eluted by hexanes and ethyl acetate to give 1-phenylbuta-2,3-dien-1-ol (292.0 mg, 67 %).

Allenyl alcohol precursor of **3-2-3h** was prepared by Ma's methodology.^{64b}

Hex-1-yn-3-ol (556.2 μ L, 5.0 mmol), octanal (779.5 μ L, 5.0 mmol), morpholine (691.2 μ L, 8.0 mmol), ZnI₂ (1276 mg, 4.0 mmol) and toluene (20 mL) were added into 50 mL round bottom reaction flask containing stir bar, and the flask was equipped with refluxing condenser. The reaction mixture was refluxed for 8 h in oil bath, cooled to room temperature and filtered through silica gel (about 20 g) and washed by mixture of hexanes and ethyl acetate (120 mL, ratio is 4:1). The solvent was removed in vacuum and the resulting residue was purified on silica gel column, which was eluted by hexanes and ethyl acetate to give tetradeca-5,6-dien-4-ol (262.5 mg, 25 %).

Allenyl alcohol precursor of **3-2-3i** was prepared by Harada's methodology.^{64c}

Stannous chloride (568.8 mg, 3.0 mmol), 1-bromo-2-butyne (262.55 μ L, 3.0 mmol), sodium iodide (450.0 mg, 3.0 mmol) and DMF (5 mL) were added into 10 mL reaction

flask containing stir bar, and the reaction mixture was stirred for 1.5 h at room temperature. Then the reaction mixture was cooled at 0 °C, and benzaldehyde (305.7 55 μ L, 3.0 mmol) in DMF (1 mL) was added dropwise and stirred for overnight. The reaction mixture was quenched by saturated ammonium chloride (20 mL) and followed by diethyl ether extraction (25 mL $\times$ 4). The ether layers were combined and dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure, the resulting residue was subjected to silica gel chromatography eluted by hexanes and ethyl acetate system to yield 2-methyl-1-phenylbuta-2,3-dien-1-ol (384.0 mg, 80 %).

2. Esterification of allenyl alcohol

Allenyl carbinol esters (**3-2-3**) were prepared from allenyl alcohols under standard conditions.^{59a}

1-Phenylbuta-2,3-dien-1-ol (292.1 mg, 2.0 mmol), and triethylamine (417 μ L, 3 mmol) were dissolved in dry dichloromethane (5.0 mL) at 0°C, in which the acetyl chloride (213 μ L, 3 mmol) was added slowly over a 5 min period, and stirred for 10 min at 0 °C. The resulting solution was stirred for 5 h at room temperature, and then saturated ammonium chloride solution (20 mL) was added to the reaction mixture, followed by diethyl ether extraction (25 mL $\times$ 3). The ether layers were combined and dried over anhydrous Na₂SO₄, filtered and evaporated under reduced pressure to yield **3-2-3a** (357.2 mg, 95%).

3. Isomerization of allenyl carbinol esters (3-2-3) by gold complex 3-2-4.^{59a}

Allenyl carbinol esters (3-2-3) (0.25 mmol) and gold complex (3-2-4) (1 mol %) were dissolved in CDCl_3 (1 mL), and stirred at room temperature and monitored by ^1H -NMR until the reaction was completed. All allenyl carbinol esters 3-2-3 efficiently isomerized into dienes 3-2-2, and NMR yields were at least 95 %.

4. Fluorination of dienes (3-2-2) with Selectfluor

The solvent (CDCl_3) in the above reaction was carefully removed in vacuum, and Selectfluor (106 mg, 1.2 equiv.) and acetonitrile (1.5 mL) were added and stirred at room temperature for 3 h.

For 3-2-1a, the reaction mixture was subjected to short silica gel chromatography separation, which was eluted by gradient elution of hexanes and ethyl acetate.

For 3-2-1e, α,α,α -trifluorotoluene (12.5 μL , 0.1 mmol) and CD_3CN (0.5 mL) were added into reaction mixture to measure ^{19}F NMR yield.

For 3-2-1g, 3-2-1h and 3-2-1i, the reaction mixture was filtered through approximately 10 g of silica gel, and washed by mixture of hexanes and ethyl acetate (4:1, 70 mL). Then, the solvent was evaporated, and the residue was dissolved in methanol (2-3 mL). $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (186 mg, 0.4 mmol) was added into the methanol solution and the solution was placed at 0 °C (ice bath), and NaBH_4 (15 mg, 0.4 mmol) was added over 15 min, and the reaction was stirred for another 20 min at room temperature. Then the mixture was directly subjected to silica gel chromatography, eluted by a hexane-ethyl acetate system to give the corresponding allylic alcohols.

(*E*)-1-fluoro-4-phenylbut-3-en-2-one (**3-2-1a**)

^1H NMR (500 MHz, CDCl_3): δ 7.77 (d, $J = 16.5$ Hz, 1H), 7.61 (m, 2H), 7.44 (m, 3H), 7.03 (dt, $J = 16.0, 2.0$ Hz, 1H), 5.05 (dd, $J = 47.5, 1.0$ Hz, 2H).

^{13}C NMR (125 MHz, CDCl_3): δ 194.84 (d, $J = 18.3$ Hz), 145.09 (d, $J = 3.1$ Hz), 134.06, 131.20, 129.06, 128.73, 119.77, 84.78 ($J = 184.1$ Hz)

^{19}F NMR (470 MHz): δ -228.68 (dt, $J = 47.5, 2.4$ Hz).

FTIR (ATR, Attenuated Total Reflectance)/ cm^{-1} : 3029, 2927, 1709, 1689, 1607, 1576, 1495, 1450, 1333, 1202, 1167, 1036, 998, 977, 748, 688.

(*E*)-1-fluorohept-3-en-2-one (**3-2-1e**)

NMR spectra was performed on mixture of **1e** and ethyl acetate.

^1H NMR (500 MHz, CDCl_3): (significant peaks) δ 7.05 (dt, $J = 16.0, 6.6$ Hz, 1H), 6.37 (d, $J = 16.0$ Hz, 1H), 4.97 (d, $J = 47.5$ Hz, 2H), 2.26 (m, 2H).

^{19}F NMR (470 MHz): δ -229.53 (t, $J = 47.9$ Hz).

(*E*)-1-fluoropentadec-3-en-2-ol, derived from (**3-2-1g**)

Contain less than 3 % saturated alcohol.

^1H NMR (400 MHz, CDCl_3): δ 5.83 (dt, $J = 15.2, 8.0$ Hz, 1H), 5.41 (dd, $J = 15.6, 6.4$ Hz, 1H), 4.45 (m, 0.5H), 4.34 (m, 2H), 4.20 (m, 0.5H), 2.04 (q, $J = 6.8$ Hz, 1H), 1.25-1.39 (m, 18H), 0.87 (t, $J = 7.2$ Hz, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 135.54 (d, $J = 1.5$ Hz), 125.89 (d, $J = 8.5$ Hz), 86.36 (d, $J = 171.2$ Hz), 71.48 (d, $J = 19.4$ Hz), 32.31, 31.89, 29.64, 29.61, 29.56, 29.44, 29.32, 29.12, 28.91, 22.66, 14.09.

^{19}F NMR (376 MHz): δ -225.54 (dt, $J = 48.1, 17.7$ Hz).

FTIR (ATR)/ cm^{-1} : 3420, 2924, 2854, 1466, 1012.

(*E*)-7-fluorotetradec-4-en-6-ol (dr = 4:1), derived from **(3-2-1h)**

Contain less than 3 % saturated alcohol.

Major diastereomers:

^1H NMR (400 MHz, CDCl_3): δ 5.79 (dt, $J = 15.2, 7.2$ Hz, 1H), 5.42 (dd, $J = 15.6, 7.6$ Hz, 1H), 4.37 (m, 0.5H), 4.23 (m, 0.5H), 4.05 (m, 1H), 2.04 (m, 2H), 1.26-1.62 (m, 14H), 0.87 (m, 6H).

^{13}C NMR (100 MHz, CDCl_3): δ 135.47, 127.34 (d, $J = 7.0$ Hz), 96.61 (d, $J = 171.1$ Hz), 74.72 (d, $J = 20.9$ Hz), 34.38, 31.72, 30.94, 29.34, 29.09, 24.93, 22.59, 22.08, 14.04, 13.61.

^{19}F NMR (376 MHz): δ -192.30 (m).

FTIR (ATR)/ cm^{-1} : 3427, 2926, 2857, 1464, 970.

(*E*)-1-fluoro-3-methyl-4-phenylbut-3-en-2-ol, derived from **(3-2-1i)**

^1H NMR (500 MHz, CDCl_3): δ 7.37 (m, 2H), 7.27 (m, 3H), 6.67 (s, 1H), 4.61 (dd, $J = 9.0$, 2.5 Hz, 0.5H), 4.51 (m, 2H), 4.39 (t, $J = 7.5$ Hz, 0.5H), 1.92 (d, $J = 1.5$ Hz, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 136.95, 134.67 (d, $J = 7.0$ Hz), 128.97, 128.16, 127.73 (d, $J = 1.6$ Hz), 126.79, 85.73 (d, $J = 172.2$ Hz), 75.80 (d, $J = 19.4$ Hz), 14.58.

^{19}F NMR (470 MHz): δ -225.33 (dt, $J = 47.5$, 17.4 Hz).

Stereochemistry was determined by ROSEY spectra.

FTIR (ATR)/ cm^{-1} : 3397, 2977, 1723, 1448, 1055, 903, 751, 698.

3.3. Combination of Copper and Selectfluor: a Strong yet Selective Oxidant System

3.3.1 Hydroxyl Group Directed C-H Amination

3.3.1.1 Background

Selectfluor is not only a good fluorinating agent, but also a good oxidant. Selectfluor can be an oxidant in functional group transformations,⁴⁵ halogens (iodine),⁴⁶ C-H functionalization,⁴⁷ gold-catalyzed oxidative C-C and C-heteroatom bond formation,⁴⁸ and palladium catalyzed C-C or C-heteroatom bond formation.⁴⁹ Banks and co-workers have reported a hydroxyl group directed C-H amination using Selectfluor in refluxing acetonitrile (Scheme 14).⁶⁶ (-)-Menthol **3-3-1-1a** and Selectfluor (2.2 equiv.) were refluxed in acetonitrile for 16 h, and oxazoline derivative **3-3-1-2a** was isolated in moderate yield; **3-3-1-2a** gave amination product **3-3-1-3a** after basic hydrolysis.

Scheme 14. Hydroxyl Group Directed C-H amination by Selectfluor in Acetonitrile

Although Bank's amination protocol is novel and potentially very useful, its harsh reaction conditions make it unattractive for synthetic purposes.

3.3.1.2 Copper Catalyzed C-H Amination by Selectfluor

To make Bank's amination milder, we screened various transition metal catalysts and found that copper salts could be suitable catalysts, but CuBr was the best (Table 6).

Table 6. Screening for a Suitable Transition Metal Catalyst

3-3-1-1a	3-3-1-2a	3-3-1-3a
Entry	Cat. (10 %)	Yield ^a of 3-3-1-3a
1	—	No Reaction
2	Sc(OTf) ₃	No Reaction
3	Eu(OTf) ₃	No Reaction
4	In(OTf) ₃	No Reaction
5	AgOTf	No Reaction
6	AuCl	No Reaction
7	FeCl ₂ 4H ₂ O	No Reaction
8	FeBr ₂	No Reaction
9	CuCl	43%
10	CuBr	50%
11	CuI	45%
12	CuCl ₂	43%
13	Cu(OAc) ₂	36%

^a Isolated yield, calculated on the menthol.

(-)-Menthol (1 equiv.), Selectfluor (2.2 equiv.) and CuBr (10 mol %) were mixed in acetonitrile at room temperature for 2 h, and following basic hydrolysis the reaction gave moderate yields of the amination product **3-3-1-3a** (entry 10, Table 6). We confirmed through HRMS that the reaction goes through an oxazoline intermediate. Our conditions were definitely milder than Banks' but when we examined the substrate scope using our optimized conditions, we were disappointed to find that only two alcohol substrates worked well (Table 7); most of the other alcohols tried were oxidized by the combination

of CuBr and Selectfluor. The two alcohols in Table 7 that were successfully aminated using our conditions share a similar feature, that is, they are sterically very hindered.

Table 7. Substrate Scope of Hydroxyl Group Directed C-H Amination

Entry	Alcohol (1)	Yield ^a of 3
1	 3-3-1-1a	 3-3-1-3a, 50%
2	 3-3-1-1b	 3-3-1-3b, 43%

^a Isolated yield, calculated on 3-3-1-1.

We were not the only ones that realized the importance of Bank's approach; very recently, Baran and co-workers reported a C-H amination protocol,⁶⁷ which is very similar to ours, but with wider scope. The key to their success was the use of a zinc salt in combination with copper(II) bromide. In the following sections, we explore the oxidative ability of copper and Selectfluor.

3.3.1.3 Experimental

The alcohol 3-3-1-1 (0.25 mmol, 1 equiv.), Selectfluor (0.55 mmol, 2.2 equiv.) and CuBr (0.025 mmol, 0.1 equiv.) were dissolved in acetonitrile (5 mL) and stirred at room

temperature for 2 h. NaOH (1.25 mmol, 5 equiv.) and water (5 mL) were added into the reaction mixture and stirred at room temperature for 5 h. Then, saturated ammonium chloride solution (30 mL) was added into the reaction mixture and extracted with diethyl ether (25 mL×4). The ether layers were combined and dried over Na₂SO₄, filtered, and evaporated under reduced pressure to give the crude product. Silica gel flash chromatography of the crude product [hexanes-ethyl acetate (4:1) to hexanes-ethyl acetate (2:1)] yielded the amination product **3-3-1-3**, which was **3-3-1-3** characterized by its ¹H and ¹³C NMR spectra.

N-(2-((1S,2R,4R)-2-hydroxy-4-methylcyclohexyl)propan-2-yl)acetamide (**3-3-1-3a**)

¹H NMR (500 MHz, CDCl₃): δ 0.90 (d, *J*=6.5Hz, 3H), 0.93 (m, 2H), 1.06 (m, 1H), 1.30 (s, 3H), 1.46 (s, 3H), 1.50 (m, 2H), 1.66 (m, 1H), 1.78 (m, 1H), 1.86 (s, 3H), 1.91 (m, 1H), 3.67 (dt, *J*=4.0, 10Hz, 1H), 5.52 (bs, 1H).

¹³C NMR (CDCl₃, 125 MHz): δ 21.379, 21.836, 24.965, 25.392, 25.499, 31.374, 34.387, 45.756, 51.539, 56.819, 72.186, 169.710.

N-(4-hydroxy-2,6-dimethylheptan-2-yl)acetamide (**3-3-1-3b**)

^1H NMR (500 MHz, CDCl_3): δ 0.90 (d, $J=6.5\text{Hz}$, 3H), 0.91 (d, $J=6.5\text{Hz}$, 3H), 1.20 (m, 1H), 1.38 (s, 3H), 1.42 (s, 3H), 1.44 (m, 2H), 1.70 (m, 2H), 1.87 (s, 3H), 2.96 (d, $J=4.5\text{Hz}$, 1H), 3.95 (m, 1H), 7.09 (bs, 1H).

^{13}C NMR (CDCl_3 , 125 MHz): δ 22.203, 23.072, 24.308, 24.537, 25.216, 28.398, 48.220, 48.976, 53.309, 67.096, 169.985.

3.3.2 Copper-Mediated Oxidation of Amides into Imides by Selectfluor (F-TEDA-BF₄)

3.3.2.1 Background

Imides are core structures in many therapeutic and agrochemical agents (Figure 5).⁶⁸ Imides can be prepared through condensation of carboxylic acid derivatives and ammonia or primary amine,⁶⁹ or other protocols,⁷⁰ such as cross coupling,^{70d, e} oxidation of oxazoles,^{70b} N-formylation,^{70c} and reaction of aziridines.^{70a}

Figure 5. Therapeutic and Agrochemical Agents Containing the Imide Moiety

We were interested in a direct synthetic pathway to prepare an unsymmetrical imide by direct oxidation of the α -methylene group of an amide. Several methods to accomplish this transformation have been reported.⁷¹ However, efficient methodologies for α -methylene oxidation are relatively rare;^{71e, 71i} most of the reported oxidative methods suffer from low yields and/or limited substrate scope.^{71a-d, 71f-h, 71j} For example, although RuO₄ is widely recognized as a strong oxidant for amide oxidation,^{71a, 71f, 71j} it can oxidize even unactivated tertiary C-H bonds.⁷² Of the efficient oxidative methods reported thus far, one employs a heavy metal (chromium VI) reagent,⁷¹ⁱ and although Nicolau and coworkers demonstrated that such toxic heavy metal oxidant could be supplanted by the more environmentally benign Dess–Martin periodinane (DMP), their method required heating the hypervalent iodine reagent to relatively high temperatures (80–85°C).^{71e} In this

section we report a room temperature synthesis of imides from amides using a combination of copper (I) bromide and Selectfluor (F-TEDA-BF₄).

3.3.2.2 Screening for Suitable Oxidative Conditions

Selectfluor is an easy-handling and bench-stable electrophilic fluorinating agent.³⁷ In the course of our work on Selectfluor/gold-mediated oxidative coupling³ and copper catalyzed amination,⁷³ we found that a combination of copper (I) bromide and Selectfluor is a strong oxidation system that can efficiently oxidize amides into imides. We chose amide **3-3-2-1a** as a substrate to screen for optimum oxidation conditions (Table 8). Amide **3-3-2-1a**, Selectfluor (2.2 equiv.) and copper bromide (10%) were mixed in acetonitrile at room temperature for 1 h to afford 10% yield of imide **3-3-2-2a**--the same yield as the load of the copper salt--together with unreacted starting material (entry 1). When the copper load was increased to 1 equivalent, we obtained a moderate yield of imide (entry 2); however, use of 1 equivalent of Selectfluor gave a low yield (entry 3). Entries 4 and 5 showed that both copper bromide and Selectfluor are indispensable for the oxidation to proceed. CuCl₂ alone, or together with Selectfluor (entries 6 and 7), has no effect on the transformation. Similarly, the use of different copper salts, such as CuOTf, CuBF₄, CuCl, CuI and Cu(OTf)₂ gave small amounts of imide (entries 8 to 12). Since the main difference among these copper salts is the counterion, which showed remarkably dissimilar results, and because it has been reported that bromide can be oxidized to bromine cation⁷⁴ we examined the role of bromide, and found that bromide itself (KBr) was not the active species (entry 13). Inspired by White's work,⁷⁵ CuBr was added in five portions over 32 min, improving our results (entry 14). Using a similar portion-wise addition, the highest yield of imide was attained increasing the amount of

both Selectfluor and CuBr (entry 15); hence, it was chosen as the optimized condition for this oxidation.

Table 8. Screening for Optimal Oxidation Conditions^a

3-3-2-1a **3-3-2-2a**

Entry	Selectfluor (equiv.)	Metal (or other) (equiv.)	% Yield ^b (SM) ^c
1	2.2	CuBr (0.1)	10(90)
2	2.2	CuBr (1.0)	66(33)
3	1.0	CuBr (1.0)	12(88)
4	2.2	—	0(100)
5	—	CuBr (1.0)	0(100)
6	—	CuCl ₂ (1.0)	0(100)
7	2.2	CuCl ₂ (1.0)	0(100)
8	2.2	CuOTf (1.0)	25(75)
9 ^d	2.2	CuBF ₄ (1.0)	0(100)
10	2.2	CuCl (1.0)	7(93)
11	2.2	CuI (1.0)	6(94)
12	2.2	Cu(OTf) ₂ (1.0)	19(81)
13	2.2	KBr (1.0)	0(100)
14 ^e	2.2	CuBr (1.0)	83(17)
15 ^f	2.5	CuBr (1.0)	92(8)
16 ^{f,g}	2.5	CuBr (1.0)	37(63)

^a All reactions were conducted in acetonitrile at room temperature for 1 h. ^b NMR yield. Benzyl bromide as internal standard. ^c Starting material. ^d CuBF₄ was generated in situ. ^e Copper bromide was added 5 portions over 32 min. ^f Copper bromide was added 6 portions over 40 min. ^g Acetonitrile: water=100:3 (v/v).

Although water may be a possible oxygen source in this oxidation, small amounts of water (entry 16) had a deleterious effect in this reaction.

3.3.2.3 Examination of Amide Scope

Next, we examined the substrate scope. Amide **3-3-2-1a** (Table 9, entry 1, Condition A) was readily oxidized by the combination of copper bromide and Selectfluor to give imide **3-3-2-2a** in excellent yield after 1 h, together with small amounts of unreacted starting material. Similarly, amides **3-3-2-1b**, **3-3-2-1c**, **3-3-2-1d** and **3-3-2-1e** afforded the corresponding imides in high yields (entries 2-5). Branched amide **3-3-2-1f** gave moderate yields (entry 6); this may be due to the steric hindrance, but the unreacted amide could be recycled. Amide **3-3-2-1g** also gave moderate yield of **3-3-2-2g** together with small amount of unreacted **3-3-2-1g** (entry 7), but the reason for this lower yield is unknown. The oxidation of **3-3-2-1h** gave **3-3-2-2h** in high yield (entry 8); and the reaction with amidoesters **3-3-2-1i**, **3-3-2-1j** was chemoselective (entries 9 and 10). Unfortunately, the presence of hydroxyl groups, double or triple bonds is not tolerated in this reaction due to a possible over oxidation.

3.3.2.4 Investigation of the Oxygen Source

There are two possible sources of oxygen for the newly incorporated oxygen in imide **3-3-2-2**, namely dioxygen in air and trace amounts of water in the reaction media. We conducted the reaction under nitrogen, and found that 50% of the imide was formed.⁷⁶ Since commercial Selectfluor always contains trace amounts of water,⁷⁷ we could not conduct the reaction under strictly anhydrous conditions. On the other hand, when substrate **3-3-2-1b** was oxidized in acetonitrile (dried shortly before use) containing 0.1 % H₂¹⁸O, we observed ¹⁸O (45%) incorporation in imide **3-3-2-2b** (confirmed by HRMS and ¹³C NMR of **3-3-2-2b**) (Figure 6 and 7 in experimental section), which

clearly demonstrated that trace water in the reaction acts as the oxygen source (Scheme 15).

Since all amides in Table 9 are secondary amides, a tertiary amide **3-3-2-1k** was employed to examine whether the N-H moiety is important (Scheme 16). Most of the unreacted amide **3-3-2-1k** was recycled and only small amounts of oxidation product **3-3-2-2b** (20%) were isolated, together with an unknown by-product, the yield being lower than that obtained with a similar substrate in entry 2 of Table 9, hinting to the possibility that N-H played a role in the oxidation.

Scheme 15. Evidence of Trace H₂O as Ultimate Oxygen Source

Scheme 16. Tertiary Amide not Suitable for This Oxidation

3.3.3 Copper Catalyzed Oxidation of Amide into Imide by F-TEDA-PF₆

3.3.3.1 Examination of Amide Substrate Scope

We reported above that a combination of stoichiometric amounts of CuBr and excess Selectfluor (F-TEDA-BF₄) can oxidize amides into imides at room temperature.⁷⁸ Although this method is efficient, the use of a stoichiometric amount of copper makes the methodology unattractive for process chemists, so we investigated alternative conditions that would permit use of a catalytic amount of a cuprous salt. Here we are pleased to disclose an exciting finding: a simple counterion exchange on Selectfluor (F-TEDA-PF₆ instead of F-TEDA-BF₄) permits the amount of copper catalyst to be reduced significantly (to 10 mol %). The oxidizing power of F-TEDA-PF₆, which can be prepared easily from Selectfluor, is at least as potent as Selectfluor itself; amides are oxidized to imides as rapidly and efficiently as we reported previously despite the significant reduction in CuBr loading.

The results of amide oxidation experiments are summarized in Table 9, Condition **B**. First, amide **3-3-2-1a**, CuBr (10 mol %) and F-TEDA-PF₆ (2.2 equiv.) reacted in acetonitrile for 5 h, giving a similar yield of imide (**3-3-2-2a**) (entry 1) (condition **B**) to our previous result using condition **A**, albeit utilizing longer reaction times. Similarly, amides **3-3-2-1b** (entry 2) and **3-3-2-1c** (entry 3) were also efficiently oxidized to the corresponding imides in good yield under the new condition **B**. Entry 4 (amide **3-3-2-1d**) demonstrated that condition **B** was superior to condition **A**. In entries 5, 6 and 7, amides worked well with the new protocol, without exceptions. The highest yield was observed with fluorinated amide **3-3-2-1h** (entry 8), and esters groups tolerated both, the new and the former oxidative systems (entries 9 and 10). Unfortunately, both oxidative systems,

the combination of CuBr (1.2 equiv.) and F-TEDA-BF₄ or the combination of CuBr (0.1 equiv.) and F-TEDA-PF₆, failed to oxidize lactam **3-3-2-11** (Scheme 17), indicating that only acyclic amides are suitable substrate for these two oxidative systems.

Scheme 17. Limitation of The Copper Catalyzed/Mediated oxidations

3.3.3.2 Mechanistic Investigations

Nicolaou has proposed a mechanism^{71e} for amide oxidation by DMP, in which the amide was dehydrogenated to form imine intermediate **3-3-2-3**. Compound **3-3-2-3** reacts with water to generate hemiaminal **3-3-2-4**, which can be further oxidized by DMP to give the final product **3-3-2-2** (Scheme 18). Initially, we thought that our oxidation probably proceeded through a similar reaction mechanism since the ultimate source of the newly formed oxygen comes from trace water in the reaction mixture.

Scheme 18. Nicolaou's Proposal of Amide Oxidation by DMP

In order to confirm the above assumption, we added an aromatic compound with moderate electron density, N-arylacetamide, to the reaction mixture to capture intermediate **3-3-2-3**; two attempts are outlined in Scheme 19.

Scheme 19. Attempt to Capture Possible Imine Intermediate

Amide **3-3-2-1h** and N-arylacetamide were mixed using condition **B** in Table 9 (eq. 1, Scheme 17). Surprisingly, amide **3-3-2-1h** was not oxidized into **3-3-2-2h**; instead N-arylacetamide was fluorinated and no imine-captured product was observed. Then we added N-arylacetamide to the reaction mixture 30 min later (eq. 2). Imide (25%) and fluorinated N-arylacetamide were observed, but no imine-captured product was detected. The above two observations indicate that the oxidation pathway probably does not involve any imine intermediate. Lactam **3-3-2-1l** can not be oxidized in both systems (Scheme 17), which may be owed to that imine intermediate of lactam **3-3-2-1l** is much harder to be formed compared to acyclic amide. Therefore, possible formation of imine intermediate **3-3-2-3** can not be excluded.

It is not clear to us yet what is the role of copper salt in this oxidation. We observed that ^1H NMR of the reaction was complicated by line broadening effects when Selectfluor (or F-TEDA- PF_6) was added; however, we failed to detect the presence of paramagnetic Cu(II) by EPR at room temperature. Therefore, high valent Cu(III), possibly formed from oxidation of Cu(I) by Selectfluor, is proposed to be a possible oxidant for this reaction. Considering that a Cu(III) species may not be very stable (isolable only using specific ligands), a multi-portion addition of CuBr (in the case of Selectfluor) could reduce the decomposition of Cu(III) complex and thus improve the yield of **3-3-2-2** (see entries 2 and 14 in Table 8). A premixed solution of CuBr and Selectfluor is still active, although it gave lower yields.⁷⁹

While Selectfluor (F-TEDA- BF_4) uses stoichiometric amount of CuBr to oxidize amide, F-TEDA- PF_6 needs only catalytic amount of CuBr (10%). Browsing through literature reports of fluorination reactions, the difference between PF_6^- and BF_4^- is not significant.⁸⁰ Our rationale to explain the above difference in reactivity is the following: we assume that the different copper loading is due to the stability of the active Cu(III) species (**A**) and (**B**) (Schemes 20). Although both complexes **A** and **B** may have oxidative activity, high valence copper in **A** easily removes a fluoride ion from BF_4^- to form complex **C**, which is probably less reactive than **A**. Therefore, large amounts of CuBr are needed for oxidation in the case of F-TEDA- BF_4 . Another possibility is that the smaller water content in F-TEDA- PF_6 compared with commercial Selectfluor could account for the improved imide yields.

Scheme 20. Proposed Rationale for Different Effect of BF_4^- and PF_6^-

However, for the high valence copper in **B** it is difficult to take up the fluoride ion from PF_6^- to form complex **C** due to the stronger P-F bond in the PF_6^- , therefore, **B** remains active in the reaction system. Computational studies reported by Christe and co-workers,⁸¹ support our rationale: PF_5 has a much stronger affinity (94.9 kcal/mol) towards fluoride ion than BF_3 (83.1 kcal/mol).

3.3.3.3 Summary

We have developed a mild and efficient methodology of copper-mediated oxidation of amide into imide by Selectfluor. Simply changing the counterion on Selectfluor (F-TEDA- PF_6 instead of F-TEDA- BF_4) can significantly reduce the amount of copper (CuBr, 10 mol %) needed. The oxidative ability of copper bromide (10 mol %) and F-TEDA- PF_6 is as efficient as that of stoichiometric amount of copper bromide and Selectfluor. Both methodologies could be useful to synthesize the imide group, but more detailed mechanistic studies are needed to clarify the effects of the anion BF_4^- viz-a-viz PF_6^- .

3.3.3.4 Experimental Section

General

^1H NMR (400 MHz), ^{13}C NMR (100 MHz) and ^{19}F NMR (376 MHz) spectra were recorded on a Varian MR400 NMR spectrometer. Chemical shifts (δ) were reported as part per million (ppm). δ 7.26, δ 77.00 of CHCl_3 , 0.00 of CFCl_3 were used as internal standards for ^1H NMR, ^{13}C -NMR and ^{19}F -NMR spectra, respectively. High-resolution mass spectra (HRMS) were performed at mass spectrometry facility of Center for Regulatory and Environmental Analytical Metabolomics, University of Louisville. Melting points were measured using a DigiMelt MPA160 melting point apparatus. RTIR spectra were recorded in ATR (attenuated total reflection) solid mode using a Perkin Elmer Spectrum 100.

General procedure for the preparation of amides 3-3-2-1

The corresponding amine (3 mmol, 1.0 equiv.) and triethylamine (3 mmol, 1.0 equiv.) were dissolved in dry dichloromethane (6 mL) at 0°C , in which the corresponding acid chloride (3 mmol, 1 equiv.) was added slowly over a 10 min period, and stirred for 10 min at 0°C . The resulting solution was stirred for 30 min at room temperature, and then saturated ammonium chloride solution (15 mL) was added to the reaction mixture, followed by diethyl ether extraction (25 mL $\times$ 4). The ether layers were combined and dried over anhydrous Na_2SO_4 , filtered and evaporated under reduced pressure to yield amide **3-3-2-1**.

General procedure for copper-mediated oxidation of amide 3-3-2-1 to imide 3-3-2-2 by Selectfluor (F-TEDA- BF_4) (Condition A)

Amide **3-3-2-1** (0.25 mmol, 1 equiv.) and Selectfluor (0.625 mmol, 2.5 equiv.) were dissolved in acetonitrile (5 mL) at room temperature, and CuBr (0.3 mmol, 1.2 equiv.)

was added over a 40 min period in 6 portions. After all the CuBr was added, the resulting mixture was stirred for extra 20 min, and then acetonitrile was evaporated under reduced pressure. Saturated ammonium chloride solution (20 mL) was added into reaction mixture and extracted with diethyl ether (25 mL×4); the ether layers were combined and dried over Na₂SO₄, filtered, evaporated under reduced pressure to give the crude product. Silica gel flash chromatography of the crude product [hexanes-ethyl acetate (10:1) to hexanes-ethyl acetate (4:1)] yielded pure imide **3-3-2-2**.

Preparation of F-TEDA-PF₆ from Selectfluor (F-TEDA-BF₄)

The procedure followed a literature report.⁸⁰ Since a large amount of BF₄⁻ has a deleterious effect on CuBr loading, it is strongly recommended that F-TEDA-PF₆ contain small amounts of BF₄, and that the ratio of PF₆ and BF₄ is at least 5.5:1.

Selectfluor (1062 mg, 3.00 mmol) and ammonium hexafluorophosphate (2934 mg, 18.0 mmol) were dissolved in H₂O (10 mL) at room temperature and stirred for 1.5 h. The mixture was filtered and washed with H₂O (10 × 5 mL) and Et₂O (30 mL) to afford F-TEDA-PF₆ (1400 mg, 97 %).

General procedure for copper-catalyzed oxidation of amide **3-3-2-1 to imide **3-3-2-2** by F-TEDA-PF₆ (Condition B)**

Amide **3-3-2-1** (0.25 mmol, 1 equiv.), F-TEDA-PF₆ (0.625 mmol, 2.5 equiv.) and CuBr (0.025 mmol, 0.1 equiv.) were dissolved in acetonitrile (5 mL) and stirred at room temperature for 3-6 h, monitored by TLC until the reaction showed no further progress. The work-up was the same as in Condition A.

N-(3-Methyl-butyl)-benzamide, (**3-3-2-2a**)^{70b}

¹H NMR (400 MHz, CDCl₃): δ 8.92 (bs, 1H), 7.82 (d, *J* = 7.6 Hz, 2H), 7.51 (t, *J* = 6.4 Hz, 1H), 7.42 (t, *J* = 7.6 Hz, 2H), 2.82 (d, *J* = 6.8 Hz, 2H), 2.18 (m, 1H), 0.96 (d, *J* = 6.4 Hz, 6H).

¹³C NMR (100 MHz, CDCl₃): δ 175.9, 165.6, 133.1, 132.9, 128.9, 127.7, 46.2, 24.8, 22.52.

N-Acetyl-benzamide, (**3-3-2-2b**)^{70b}

¹H NMR (400 MHz, CDCl₃): δ 9.20 (bs, 1H), 7.88 (d, *J* = 7.6 Hz, 2H), 7.58 (t, *J* = 7.2 Hz, 1H), 7.47 (t, *J* = 7.2 Hz, 2H), 2.58 (s, 3H).

¹³C NMR (100 MHz, CDCl₃): δ 173.9, 165.9, 133.2, 132.6, 128.9, 127.8, 25.6.

HRMS *m/z* (ES⁺) calcd for C₉H₉NO₂Na ([M+Na]⁺) 186.0525, found 186.0525.

3-Methyl-*N*-octanoyl-butamide (**3-3-2-2c**)

White crystal, mp: 52-54°C.

^1H NMR (400 MHz, CDCl_3): δ 8.70 (bs, 1H), 2.57 (t, $J = 6.0$ Hz, 2H), 2.43 (d, $J = 5.6$ Hz, 2H), 2.12 (m, 1H), 1.60 (m, 2H), 1.25-1.29 (m, 8H), 0.95-0.97 (m, 6H), 0.85 (m, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 174.7, 173.6, 46.3, 37.4, 31.6, 29.0, 28.9, 25.2, 24.3, 22.6, 22.4, 14.0.

FTIR (neat)/ cm^{-1} : 3269, 3175, 2957, 2927, 1732, 1505, 1161.

HRMS m/z (ES^+) calcd for $\text{C}_{13}\text{H}_{25}\text{NO}_2\text{Na}$ ($[\text{M}+\text{Na}]^+$) 250.1777, found 250.1777.

N-Propionyl-benzamide (**3-3-2-2d**)^{70b}

^1H NMR (400 MHz, CDCl_3): δ 9.32 (bs, 1H), 7.90 (d, $J=8.4$ Hz, 2H), 7.56 (t, $J = 7.6$ Hz, 1H), 7.46 (t, $J = 7.6$ Hz, 2H), 3.00 (q, $J = 7.2$ Hz, 2H), 1.18 (t, $J = 7.2$ Hz, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 177.7, 165.8, 133.1, 132.8, 128.8, 127.8, 31.2, 8.2.

N-Benzoyl-benzamide (**3-3-2-2e**)^{70b}

^1H NMR (400 MHz, CDCl_3): δ 9.20 (bs, 1H), 7.81 (d, $J = 8.0$ Hz, 4H), 7.53 (t, $J = 7.2$ Hz, 2H), 7.42 (t, $J = 7.6$ Hz, 4H).

^{13}C NMR (100 MHz, CDCl_3): δ 166.6, 133.3, 132.9, 128.8, 127.9.

Heptanoic acid (2-ethyl-hexanoyl)-amide (**3-3-2-2f**)

White crystal, mp: 45-47°C.

^1H NMR (400 MHz, CDCl_3): δ 8.75 (bs, 1H), 2.74 (t, $J = 7.6$ Hz, 2H), 2.42 (m, 1H), 1.57-1.66 (m, 4H), 1.36-1.47 (m, 2H), 1.22-1.26 (m, 10H), 0.84-0.90 (m, 9H).

^{13}C NMR (100 MHz, CDCl_3): δ 176.3, 176.0, 49.4, 37.5, 31.7, 31.5, 29.5, 28.8, 25.4, 24.2, 22.7, 22.5, 13.9, 13.8, 11.7.

FTIR (neat)/ cm^{-1} : 2959, 2931, 2860, 1738, 1686, 1510, 1166.

HRMS m/z (ES^+) calcd for $\text{C}_{15}\text{H}_{29}\text{NO}_2\text{H}$ ($[\text{M}+\text{H}]^+$) 256.2271, found 256.2271. Calcd for $\text{C}_{15}\text{H}_{29}\text{NO}_2\text{Na}$ ($[\text{M}+\text{Na}]^+$) 278.2090, found 278.2090.

Cyclohexanecarboxylic acid 4-fluoro-benzoylamide (**3-3-2-2g**)

White crystal, mp: 136-138°C.

^1H NMR (400 MHz, CDCl_3): δ 8.92 (bs, 1H), 7.89-7.93 (m, 2H), 7.12-7.17 (m, 2H), 3.32-3.38 (m, 1H), 1.22-1.98 (m, 11H).

^{13}C NMR (100 MHz, CDCl_3): δ 179.4, 165.6 ($J = 253.8$ Hz), 164.4, 130.4 ($J = 8.9$ Hz), 129.2 ($J = 3.7$ Hz), 116.0 ($J = 22.3$ Hz), 44.5, 28.9, 25.8, 25.5.

^{19}F NMR (376 MHz): δ -105.1 (m).

FTIR (neat)/ cm^{-1} : 3426, 2932, 2855, 1680, 1602, 1492, 1237, 1157.

HRMS m/z (ES^+) calcd for $\text{C}_{14}\text{H}_{16}\text{FNO}_2\text{H}$ ($[\text{M}+\text{H}]^+$) 250.1237, found 250.1237. Calcd for $\text{C}_{14}\text{H}_{16}\text{FNO}_2\text{Na}$ ($[\text{M}+\text{Na}]^+$) 272.1057, found 272.1056.

4-Fluoro-*N*-propionyl-benzamide (**3-3-2-2h**)

White crystal, mp: 118-120°C.

^1H NMR (400 MHz, CDCl_3): δ 9.48 (bs, 1H), 7.95-7.98 (m, 2H), 7.12-7.16 (m, 2H), 3.00 (q, $J = 7.2$ Hz, 2H), 1.18 (t, $J = 7.2$ Hz, 3H).

^{13}C NMR (100 MHz, CDCl_3): δ 178.1, 165.6 ($J = 253.8$ Hz), 164.8, 130.6 ($J = 9.7$ Hz), 128.9 ($J = 3.8$ Hz), 115.9 ($J = 22.3$ Hz), 31.2, 8.1.

^{19}F NMR (376 MHz): δ -105.1 (m).

FTIR (neat)/ cm^{-1} : 1690, 1604, 1469, 1370, 1239.

HRMS m/z (ES^+) calcd for $\text{C}_{10}\text{H}_{10}\text{FNO}_2\text{H}$ ($[\text{M}+\text{H}]^+$) 196.0768, found 196.0768. Calcd for $\text{C}_{10}\text{H}_{10}\text{FNO}_2\text{Na}$ ($[\text{M}+\text{Na}]^+$) 218.0587, found 218.0587.

Benzoic acid 6-benzoylamino-6-oxo-hexyl ester (**3-3-2-2i**)

White crystal, mp: 86-88°C.

^1H NMR (400 MHz, CDCl_3): δ 9.33 (bs, 1H), 8.00 (d, $J = 8.0$ Hz, 2H), 7.89 (d, $J = 8.0$ Hz, 2H), 7.37-7.57 (m, 6H), 4.30 (t, $J = 6.8$ Hz, 2H), 3.02 (t, $J = 7.2$ Hz, 2H), 1.73-1.83 (m, 4H), 1.52-1.57 (m, 2H).

^{13}C NMR (100 MHz, CDCl_3): δ 176.7, 166.6, 165.9, 133.1, 132.8, 132.7, 130.4, 129.5, 128.8, 128.3, 127.9, 64.8, 37.5, 28.6, 25.6, 23.7.

FTIR (neat)/ cm^{-1} : 3426, 1716, 1683, 1457, 1275, 1244.

HRMS m/z (ES^+) calcd for $\text{C}_{20}\text{H}_{21}\text{NO}_4\text{H}$ ($[\text{M}+\text{H}]^+$) 340.1543, found 340.1544. Calcd for $\text{C}_{20}\text{H}_{21}\text{NO}_4\text{Na}$ ($[\text{M}+\text{Na}]^+$) 362.1363, found 362.1363.

4-Benzoylamino-4-oxo-butyric acid methyl ester (**3-3-2-j**)⁸²

^1H NMR (400 MHz, CDCl_3): δ 9.49 (bs, 1H), 7.90 (d, $J = 6.8$ Hz, 2H), 7.57 (t, $J = 7.2$ Hz, 1H), 7.47 (t, $J = 7.6$ Hz, 1H), 3.63 (s, 3H), 3.31 (t, $J = 6.0$ Hz, 1H), 2.67 (t, $J = 6.4$ Hz, 1H).

^{13}C NMR (100 MHz, CDCl_3): δ 175.5, 172.9, 165.9, 133.1, 132.5, 128.8, 127.9, 51.8, 32.9, 28.1.

Figure 6. HRMS of Imide **3-3-2-2b** and O^{18} labeled **3-3-2-2b**

Figure 7. ^{13}C NMR of Carbonyl Groups in Imide **3-3-2-2b** and O^{18} labeled **3-3-2-2b**

4. SEARCH FOR CYTOTOXIC AGENTS FROM THE AMAZONIAN RAINFOREST

4.1. Background

Nature has been a major source of inspiration for the development of front-line-drugs for cancer, Alzheimer's and other diseases. Some well-known drugs or drug candidates isolated from natural source are shown in Figure 8.

Figure 8. Drugs and Drug Candidates Isolated from Natural Source

All of them have interesting biological activities and complex architecturally skeletons, but they pose a synthetic challenge.⁸³ A multidisciplinary and international program that included the Hammond group--formed in the early 1990's--isolated many bioactive compounds from different plant sources in the Amazonian rainforest. In some cases bioactive analogs were synthesized in the Hammond group and our collaborators tested their biological activities. This chapter reports our results with 13 targeted plants from the Amazonian rainforest (Table 10).

Table 10. List of Targeted Plants

Plant	Extract No.	Voucher No.	3T3	H460	ME180	DU145	MCF-7	M-14	HT-29	PC3	K562
<i>Crematosperma microcarpum</i> (Annonaceae)	2967	19759	5	7	5	6	7	5	7	5	7
<i>Physalis angulata</i> (Solanaceae)	1159	18597	9	7	9	7	7	7	9	7	7
<i>Humiria balsamifera</i> (Humiriaceae)	2278	19478	5		1		1	5		3	5
<i>Rollinia andicola</i> (Annonaceae)	2900	19877	7	9	5	3	7	5	5	5	5
<i>Bocconia intengrifolia</i> (Papaveraceae)	2367	19644	9	9	9	9	9	9	9	9	9
<i>Hyptis Lantanaefolia</i> (Lamiaceae)	2352	19547	9	9	9	9	9	9	9	9	9
<i>Hippeastrum puniceum</i> (Amaryllidaceae)	435	17506	9	9	9	9	7	9	9	9	9
<i>Aniba riparia</i> (Lauraceae)	2239	19366	7	7	7	7	7	7	7	7	1
<i>Terminalia Amazonia</i> (Combretaceae)	3531	20515	5	5	5	3	5	5	5	5	3
<i>Piper concinnatoris</i> (missing)	1962	19065	9	9	9	7	9	7	9	9	9
<i>Paullinia</i> (missing) (Sapindaceae)	2940	19705	9	7	7	7	7	7	6	7	7
<i>Plukenetia volubillis</i> (Euphorbiaceae)	3498	20465	7	7	7	7	9	7	7	9	7
<i>Scutia</i> (missing) (Phammaceae)	3579	19883	9	9		9	9		9	9	9

GI50	
1	> 0.5 mg/mL
2	= 0.5 mg/mL
3	between 0.125 and 0.5 mg/mL
4	= 0.125 mg/mL
5	between 0.031 and 0.125 mg/mL
6	= 0.031 mg/mL
7	between 0.008 and 0.031 mg/mL
8	= 0.008 mg/mL
9	< 0.008 mg/mL

3T3, BALB/3T3 clone A31 embryonic mouse fibroblast cells;
H460, human large cell lung cancer;
ME180, human cervical carcinoma;
DU145, human prostate carcinoma;
MCF-7, human breast adenocarcinoma;
M-14, human melanoma;
HT-29, human colon adenocarcinoma;
PC3, human prostate adenocarcinoma;
K562, human chronic myelogenous leukemia cells.

4.2. *Physalis angulata* L. (Solanaceae)

The genus *Physalis* (Solanaceae) is represented by almost 90 species distributed throughout the tropical and subtropical regions of the world where it has been widely used in folk medicine by developing countries.⁸⁴ As a result of its medicinal value, there has been significant interest to evaluate the phytochemical and pharmaceutical properties of *Physalis angulata* L. Previously, physanolide,⁸⁵ and other withanolides,⁸⁶ including withangulatin,^{86b, c, 86f} physangulin,^{86a, 86d, 86k, l} and physalin,^{86c, 86g-i, 87} in addition to other constituents^{86e} isolated from *P. angulata*, were found to show significant biological activity.⁸⁸

Here we disclose the discovery of three new anti-proliferative withanolides with an unusual carbon framework, namely, physangulidines A (**1**), B (**2**) and C (**3**), isolated from *P. angulata* L. using a bioassay-directed isolation technique. The ethanol extract of the dried plant was partitioned between water and dichloromethane, the active organic layer subjected to silica gel column chromatography and the most active fractions further purified via HPLC to afford the three active components.

Physangulidine A (**1**), mp is 213.0-215.0°C (MeOH), was isolated as white crystalline needles, whose molecular formula was determined to be C₂₈H₃₆O₈ by HRMS (m/z 501.2493 [M+H]⁺, 523.2311 [M+Na]⁺). Simple analysis of ¹H, ¹³C and HSQC spectra revealed four methyls, seven methylenes, eight methines and nine quaternary carbons, and suggested 2 two hydroxyl groups, consistent with this formula. Subsequent 2D analysis (gCOSY, ROESY and HMBC) lead to the structure **1** shown in Figure 9 (For numbering see Figure 10) and the complete NMR assignment is summarized in Table 11.

Figure 9. Structures of Physangulidines A (1), B (2) and C (3)

Important structural keys included two olefinic protons H-2 (δ 6.08, dd, $J=9.8$, 2.8 Hz, 1H) and H-3 (δ 6.86, ddd, $J=9.8$, 6.3, 2.8 Hz, 1H) conjugated to carbonyl carbon C-1 (δ 201.825), consistent with an α,β -unsaturated ketone.^{86b, c, 86f} This moiety is also supported by IR (1714 cm^{-1}) and UV (220 nm).^{86b} In addition, a C-5,6 epoxide, and ketal carbon C-17 (δ 109.709) were surmised from the NMR spectra. Another critical structural feature centered around carbonyl carbon C-26 (δ 176.961) and was indicative of an *isolated* bridged δ -lactone moiety, containing two methyls and two hydroxyl groups. The structure and stereochemistry of **1** was determined by X-ray crystallography (Figure 11) and ROESY 2D NMR analysis.

Compound **1** contains 28 carbons in which C-22 and C-26⁸⁹ have been oxygenated to form a δ -lactone putting it into the class of steroids known as withanolides. Withanolides belong to a group of naturally occurring steroids having 28 carbons and derived from an

Figure 10. Carbon Numbering and Crucial ROESY Interactions of Physangulidines A (1), B (2) and C (3)

Figure 11. An *ORTEP*-3 Diagram of 1 Showing 40% Ellipsoids

Note: H atoms are shown as small spheres of arbitrary radii. Selected bond lengths (Å) and angles (deg): O1-C1, 1.178(5); O5-C26, 1.204(4); O7-C20, 1.418(3); O8-C24, 1.437(4); C13-C18, 1.535(4); C12-C13-C18, 112.5(3). The absolute structure configuration for physangulidine A (1) has been reliably determined using Cu radiation including 13 stereocenters which are as follow: (chirality at C5) S, (C6) R, (C7) S, (C8) S, (C9) S, (C10) R, (C13) R, (C14) S, (C17) R, (C20) S, (C22) R, (C24) R, (C25) R.

ergostane skeleton, in which carbons 22 and 26 (Type A) or 23 and 26 (Type B) were oxidized to form lactones (Figure 12).

Figure 12. Main structures of Withanolides Have Been Isolated

Although, to date, about 650 withanolides have been isolated from different plant sources,⁹⁰ **1** is the first withanolide having a disconnection between C-13 and C-17, which typically forms ring D of the ergostane skeleton.

Physangulidine B (**2**) (mp above 260°C) and physangulidine C (**3**) were also isolated as white needles, and both were found to have the molecular formula of $C_{28}H_{36}O_8$ by HRMS (m/z 523.2311 $[M+Na]^+$ for **2**; m/z 523.2313 $[M+Na]^+$ for **3**), indicating both are isomers of physangulidine A (**1**). After careful analysis of the NMR data (Table 12 and 13), we found that in the structures of **2** and **3**, C-13 is hydroxylated instead of at C-20 in physangulidine A (**1**), and that **2** and **3** differed from each other only at the stereochemistry at C-13 (Figure 9).

As with **1**, compounds **2** and **3** also contained the bridged δ lactone moiety and the disconnection between C-13 and C-17. Physangulidine B (**2**) was subjected to X-ray

crystallography, thus confirming the aforementioned C-20 vs C-13 hydroxyl “migration” (Figure 13) and further demonstrated that with the exception of the orientation of the C-13 methyl group, the overall conformation of **1** and **2** were very similar.

Figure 13. An *ORTEP*-3 Diagram of **2** Showing 40% Ellipsoids

Note: H atoms are shown as small spheres of arbitrary radii. Selected bond lengths (Å) and angles (deg): O1-C1, 1.188(5); O5-C26, 1.218(7); O7-C13, 1.433 (5); C13-C18, 1.527(6); O7-C13-C18, 110.4(4); C12-C13-C18, 109.4(4). The absolute structure configuration for physangulidine B (**2**) has been reliably determined using Cu radiation including 13 stereocenters which are as follow: (chirality at C5) S, (C6) R, (C7) S, (C8) S, (C9) S, (C10) R, (C13) R, (C14) R, (C17) R, (C20) R, (C22) R, (C24) R, (C25) R.

Crystals of physangulidine B (**2**) adequate for X-ray diffraction studies were grown from methylene chloride/hexane. $C_{28}H_{36}O_8$: colorless needle, 0.42 x 0.02 x 0.02 mm³, orthorhombic, space group $P2_12_12_1$, $a = 7.6354(9)$ Å, $b = 12.043(2)$ Å, $c = 27.411(5)$ Å, $\alpha = \beta = \gamma = 90^\circ$, $V = 3068.53(18)$ Å³, $D_{\text{calc}} = 1.319$ Mg/m³, $Z = 4$. For 4132 reflections $I > 2\sigma(I)$ [R(int) 0.049] the final anisotropic full matrix least-squares refinement on F^2 for 334 variables converged at $R1 = 0.055$ and $wR2 = 0.08$ with a GOF of 1.03. The absolute structure was determined by refinement of the Hooft parameter 0.0(4).

Crystals of physangulidine A (**1**), $C_{28}H_{36}O_8$: colorless prism, $0.42 \times 0.22 \times 0.09 \text{ mm}^3$, orthorhombic, space group $P2_12_12_1$, $a = 10.0880(4) \text{ \AA}$, $b = 10.5957(3) \text{ \AA}$, $c = 24.5357(11) \text{ \AA}$, $\alpha = \beta = \gamma = 90^\circ$, $V = 2622.60(18) \text{ \AA}^3$, $D_{\text{calc}} = 1.270 \text{ Mg/m}^3$, $Z = 4$. For 5195 reflections $I > 2\sigma(I)$ [$R(\text{int})$ 0.038] the final anisotropic full matrix least-squares refinement on F^2 for 425 variables converged at $R1 = 0.052$ and $wR2 = 0.118$ with a GOF of 1.03 and Hooft parameter of 0.11(6).

Unfortunately, a single crystal of **3** suitable for X-ray crystallography was failed to be isolated, owing to its unstability at room temperature for long time (three weeks). However, the ROESY 2D NMR comparison between **2** and **3** did confirm that the C-18 methyl of physangulidine C is located axially and the hydroxyl group is equatorially positioned on C-13 of **3**, while in physangulidine B (**2**) the C-18 methyl is equatorial on C-13 and the hydroxyl is axial (Figure 9). Accordingly, CH_3 -18 shows a strong correlation with H-8 and H-15 (both α and β) in compound **3**, however, CH_3 -18 of physangulidine B (**2**) has only single interaction with H-15 β observed in the ROESY spectra. A similar interaction was observed in physangulidine A: CH_3 -18 has interaction with H-8 and H-15 (α and β) on ROESY, indicating the preferred axial orientation of the methyl group in both **1** and **3**.

Table 11. All NMR Data of Physangulidine A (**1**)

Position	¹³ C NMR	HSQC	COSY	ROESY	HMBC
1	201.825				3, 4eq, 9, 19
2	129.261	6.08 (dd, <i>J</i> =9.8, 2.8 Hz, 1H)	3, 4ax	3	4
3	143.748	6.86(ddd, <i>J</i> =9.8, 6.3, 2.8 Hz, 1H)	2, 4eq	2	4
4	32.064	2.95(ax, dt, <i>J</i> =18.2, 2.8 Hz, 1H) 1.93(eq, dd, <i>J</i> =18.2, 6.3 Hz, 1H)	Hax: 2 Heq: 3	Hax: 19 Heq: 6	2, 3, 6, 7
5	62.596				3, 4, 19
6	63.008	3.14(d, <i>J</i> =2.1 Hz, 1H)	7	4eq, 7	4, 7
7	67.070	4.32(t, <i>J</i> =2.8 Hz, 1H)	6, 8	6, 8, 15α, 16	6, 8
8	33.974	1.45(m, 1H)	7, 9	7, 15α, 18, 19	
9	34.112	1.82(m, 1H)	8	12ax	6, 7, 19
10	47.459				4eq, 19
11	20.488	1.35(ax, dq, <i>J</i> =14.0, 2.8 Hz, 1H) 2.10(eq, m, 1H)		Hax: 18, 19 Heq: 12ax, 19	9, 19
12	28.535	1.48(ax, m, 1H), 1.73(eq, m, 1H)		Hax: 9, 11eq	11, 18
13	37.345	1.76(m, 1H)	18	18	18
14	90.027				8, 15, 16, 18
15	32.957	1.71(α, m, 1H) 1.85(β, m, 1H)	16	Hα: 7, 8, 16, 18 Hβ: 16, 18, 21ax	16
16	31.462	2.17(m, 2H)	15	7, 15, 21ax	
17	109.709				8, 15, 16, 21
18	14.470	0.95(d, <i>J</i> =7.0 Hz, 3H)	13	8, 11ax, 13, 15	
19	14.228	1.21(s, 3H)		4ax, 8, 11	4ax
20	75.065				21, 22, 23
21	36.926	2.40(ax, d, <i>J</i> =15.4 Hz, 1H) 1.45(eq, m, 1H)		Hax: 15β, 16, 27 Heq: 27	22, 27
22	78.743	4.62(dd, <i>J</i> =3.5, 1.4 Hz, 1H)	23	23	23
23	40.499	2.41(ax, dd, <i>J</i> =16.1, 1.4 Hz, 1H) 2.09(eq, m, 1H)	22	Hax: 22 Heq: 22, 28	28
24	69.496				21eq, 22, 23, 27, 28
25	48.461				21, 27, 28
26	176.961				21, 22, 27
27	14.041	1.14(s, 3H)		21	21
28	27.044	1.17(s, 3H)		23eq	23

Table 12. All NMR Data of Physangulidine B (2)

Position	¹³ C NMR	HSQC	COSY	ROESY	HMBC
1	201.973				3, 19
2	129.287	6.10(dd, <i>J</i> =9.8, 2.8 Hz, 1H)	3, 4ax	3	4
3	143.856	6.88(ddd, <i>J</i> =9.8, 6.3, 2.1 Hz, 1H)	2, 4eq	2, 4ax	4
4	31.978	2.96 (ax,dt, <i>J</i> =18.2, 2.8 Hz, 1H) 1.95 (eq, m, 1H)	Hax: 2 Heq: 3	Hax: 19 Heq: 3, 6	2, 3
5	62.709				3, 4ax, 19
6	62.937	3.11 (d, <i>J</i> =2.8 Hz, 1H)	7	4eq, 7	4eq, 7
7	67.063	4.39 (dd, <i>J</i> =3.5, 2.8 Hz, 1H)	6, 8	6, 8, 15α, 16α	6
8	35.278	1.68 (m, 1H)	7, 9	7, 15α	15
9	33.338	1.83 (dt, <i>J</i> =11.9, 3.8 Hz, 1H)	8, 11	12ax	19
10	47.601				4eq, 19
11	21.632	1.53(ax, m, 1H), 2.17 (eq, m, 1H)	9, 12	Hax: 19	19
12	36.766	1.65 (ax, m, 1H), 1.70 (eq, m, 1H)	11	Hax: 9	18
13	71.219				15β, 18
14	88.323				15β, 18
15	27.993	1.79 (α, m, 1H) 2.17 (β, m, 1H)	16	Hα: 7, 8, 16α Hβ: 16β, 18	16
16	33.686	2.07 (α, m, 1H) 1.94 (β, m, 1H)	15	Hα: 7, 15α Hβ: 15β	15
17	108.110				15β, 16, 20, 21
18	23.916	1.21 (s, 3H)		15β	
19	14.291	1.24 (s, 3H)		4ax, 11ax	
20	40.585	2.61 (m, 1H)	21, 22	21eq, 22	15β, 16β, 21, 23eq
21	26.453	2.00(ax, dd, <i>J</i> =14.0, 5.6 Hz, 1H) 1.59 (eq, m, 1H)	20	Hax: 27 Heq: 20, 27	20, 27
22	74.149	4.76 (dt, <i>J</i> =3.5, 2.1 Hz, 1H)	20, 23	20, 23	20, 21
23	40.933	2.03(eq, ddd, <i>J</i> =15.4, 3.5, 2.1 Hz, 1H) 2.23 (ax, dd, <i>J</i> =15.4, 2.1 Hz, 1H)	22	Hax: 22 Heq: 22, 28	20, 28
24	70.651				23, 27, 28
25	47.351				21, 27, 28
26	177.626				21, 22, 23, 27
27	14.284	1.11 (s, 3H)		21	21eq
28	27.399	1.14 (s, 3H)		23eq	

Table 13. All NMR Data of Physangulidine C (3)

Position	¹³ C NMR	HSQC	COSY	ROESY	HMBC
1	201.805				3, 9, 19
2	129.310	6.09(dd, <i>J</i> =9.8, 3.5 Hz, 1H)	3, 4ax	3	4
3	143.860	6.89(ddd, <i>J</i> =9.8, 6.3, 2.1 Hz, 1H)	2, 4eq	2	4
4	31.899	2.95 (ax, dt, <i>J</i> =18.2, 2.8 Hz, 1H) 1.95 (eq, m, 1H)	Hax: 2 Heq: 3	Hax: 19 Heq: 6	2, 3, 6
5	62.910				3, 4, 19
6	62.963	3.12(d, <i>J</i> =2.1 Hz, 1H)	7	4eq, 7	7
7	67.973	4.34(t, <i>J</i> =2.8 Hz, 1H)	6, 8	6, 8, 15α, 16α	6
8	36.496	1.40(dd, <i>J</i> =11.9, 3.5 Hz, 1H)	7, 9	7, 15α, 18, 19	6, 11ax, 15, 18
9	33.394	1.88(dt, <i>J</i> =12.6, 2.8 Hz, 1H)	8	11eq	7, 8, 11, 19
10	47.358				2, 4eq, 19
11	23.897	1.20 (ax, m, 1H) 2.27(eq, m, 1H)	12	Hax: 12 Heq: 9, 12	12
12	37.614	1.69(m, 2H)	11	11, 18	18
13	71.066				12, 15β, 18
14	89.003				12, 15β, 16β, 18
15	27.197	1.63(α, m, 1H) 2.31(β, m, 1H)	16	Hα: 7, 8, 18 Hβ: 16β, 18	16, 18
16	33.383	2.03(α, m, 1H) 1.97(β, m, 1H)	15	Hα: 7 Hβ: 15β	15
17	107.837				15β, 16, 20, 21
18	21.871	1.53 (s, 3H)		8, 12, 15	12
19	14.205	1.19(s, 3H)		4ax, 8	9
20	41.311	2.54(m, 1H)	21, 22	21eq, 22	21, 23eq
21	26.648	2.16(ax, dd, <i>J</i> =14.0, 6.3 Hz, 1H) 1.55(eq, dd, <i>J</i> =14.0, 11.9 Hz, 1H)	20	Hax: 27 Heq: 20, 27	27
22	74.415	4.73(dt, <i>J</i> =3.5, 1.4 Hz, 1H)	20, 23	20, 23	20, 21eq, 23eq
23	40.578	2.40(ax, dd, <i>J</i> =15.4, 1.4 Hz, 1H) 1.99(eq, m, 1H)	22	Hax: 22 Heq: 22, 28	20, 28
24	70.838				21, 22, 23eq, 28
25	47.605				21, 23eq, 27
26	177.574				21, 22, 27
27	14.265	1.14(s, 3H)		21	21
28	28.042	1.19(s, 3H)		23eq	23eq

Physangulidine A (**1**) was found to have significant *in vitro* antiproliferative activity against DU145 cancer cell line in bioassay (GI_{50} estimated to be 3.0 μ M) and also RWPE-1 prostate epithelial cell (GI_{50} =2.4 μ M). GI_{50} values of physangulidine B (**2**) and physangulidine C (**3**) were found to be very similar and comparatively lower than physangulidine A (**1**).

GI_{50} values of physangulidine B were determined to be 6.0 and 6.8 μ M against RWPE-1 and DU145, while in physangulidine C they were 6.6 and 6.0 μ M against RWPE-1 and DU145, respectively. Their antiproliferative activities are comparable to, or higher than, related withanolides isolated from *P. angulata*.^{86c} Because physangulidine A was the most active and abundant of the three bioactive compounds isolated from *P. angulata*, it was tested on additional cell lines, using 5-fluorouracil as positive control. The results are summarized in Table 14. Compared to 5-fluorouracil, physangulidine A has less antiproliferative activity against nonmalignant 3T3 cell and more antiproliferative activity against HT-29 and K562 cells (Table 14).

Table 14. GI_{50} Values (μ M) of Physangulidine A^a and 5-Fluorouracil on Different Cells

	3T3	H460	HuTu 80	DU145	MCF-7	M-14	HT-29	K562
Physangulidine A	4.12	3.59	4.18	3.56	5.26	4.66	2.73	2.73
5-Fluorouracil	0.28	1.67	4.92	3.11	2.18	5.03	5.92	30.32

^a GI_{50} values of physangulidine A in Table 14 are average values of three tests.

3T3: Nontumorigenic, BALB/c mouse embryo cells. H460: human lung large cell carcinoma.

HuTu 80: human duodenum Adenocarcinoma. DU145: human prostate carcinoma.

MCF-7: human breast adenocarcinoma. M-14: human amelanotic melanoma.

HT-29: human colon adenocarcinoma. K562: human chronic myelogenous leukemia.

In conclusion, we have isolated three new bioactive withanolides, physangulidines A, B and C, from *P. angulata* L. using a bioassay-guided isolation technique. Their structures,

determined by NMR and X-ray crystallography, have an unusual disconnection between C-13 and C-17, a structural feature observed for the first time in the withanolides family.

Figure 14. Selected HMBC Interactions in Physangulidines A, B and C

Figure 15. Selected ROESY Interactions in Physangulidines A, B and C

4.3. *Crematosperma microcarpum* (Annonaceae)

The genus *Crematosperma* (Annonaceae) is represented by 31 species⁹¹ but has been very poorly investigated phytochemically. There is only one study on *C. polyphlebum*,⁹² and no investigation has been conducted on *C. microcarpum*. We investigated *C. microcarpum* and isolated (±)-*trans*-dehydrodiisoeugenol (Figure 16) as the main cytotoxic agent via bioassay-directed fractionation techniques.

Figure 16. Structure of Trans-dehydrodiisoeugenol

(±)-*Trans*-dehydrodiisoeugenol has been studied previously for its antioxidant and anticancer properties on salivary gland tumor,⁹³ and it has been recently applied on a skin formulation patent.⁹⁴ We found that (±)-*trans*-dehydrodiisoeugenol has moderate cytotoxic activity on different cell lines, compared to a positive control (Doxorubicin, DOXO) (Table 15). (±)-*trans*-Dehydrodiisoeugenol was identified by comparison of its NMR data with literature reports (Table 16).⁹⁵

Table 15. GI₅₀ Values (μM) of (±)-*Trans*-dehydrodiisoeugenol^a and Doxorubicin on Different Cells

Compound	3T3	H460	DU145	MCF-7	M-14	HT-29	K562	Vero
Trans-dehydrodiisoeugenol	63.4	60.5	85.3	38.7	68.8	93.6	48.8	67.8
DOXO	0.029	0.044	0.055	0.074	0.092	0.13	0.044	0.020

^a GI₅₀ values of (±)-*Trans*-dehydrodiisoeugenol are average values of two tests.

Table 16. NMR data for (±)-*Trans*-dehydrodiisoeugenol and Literature Data

position	¹ H NMR (400 MHz)		¹³ C NMR (100 MHz)	
	ref ^{95a}	(±)- <i>Trans</i> - dehydrodiisoeugenol	ref ^{95b}	(±)- <i>Trans</i> - dehydrodiisoeugenol
2	5.14 (d, <i>J</i> =9.2 Hz, 1H)	5.08 (d, <i>J</i> =9.2 Hz, 1H)	93.79	93.77
3	3.48 (dq, <i>J</i> =9.2, 6.7 Hz, 1H)	3.43 (dq, <i>J</i> =8.8, 6.6 Hz, 1H)	45.62	45.59
4	6.75-7.00 (m, 1H)	6.75 (s, 1H)	113.31	113.28
5	-	-	132.19	132.18
6	6.75-7.00 (m, 1H)	6.77 (s, 1H)	109.23	109.20
7	-	-	144.15	144.13
1'	-	-	132.08	132.07
2'	6.75-7.00 (m, 1H)	6.96 (s, 1H)	108.92	108.89
3'	-	-	146.67	146.64
4'	-	-	145.77	145.75
5'	6.75-7.00 (m, 1H)	6.88 (m, 1H)	114.08	114.03
6'	6.75-7.00 (m, 1H)	6.88 (m, 1H)	119.96	119.96
OCH ₃ (on C3')	3.93 (s, 3H)	3.88 (s, 3H)	55.98	55.96
OCH ₃ (7)	3.90 (s, 3H)	3.87 (s, 3H)	55.94	55.91
3a	-	-	133.27	133.25
7a	-	-	146.56	146.55
α	6.40 (d, <i>J</i> =15.5 Hz, 1H)	6.34 (d, <i>J</i> =15.6 Hz, 1H)	130.92	130.89
β	6.11 (dq, <i>J</i> =15.5, 5.2 Hz, 1H)	6.09 (dq, <i>J</i> =15.6, 6.4 Hz, 1H)	123.48	123.48
γ	1.92 (d, <i>J</i> =5.2 Hz, 3H)	1.86 (dd, <i>J</i> =6.8, 1.6 Hz, 3H)	18.38	18.35
CH ₃ (on C3)	1.43 (d, <i>J</i> =6.7 Hz, 3H)	1.36 (d, <i>J</i> =6.8 Hz, 3H)	17.57	17.54
OH	5.69 (s, 1H)	5.61 (s, 1H)	-	-

4.4. *Hyptis lantanaefolia* (Lamiaceae)

Hyptis is a genus of the *Lamiaceae* family, widely spread in the tropical regions of the Americas. There are 300-400 species which may be annual or perennial, appearing as

small herbs or large shrubs.⁹⁶ *H. lantanaefolia* is a very poorly investigated species; there is only one report on the xanthine oxidase inhibitory activity of the ethanol extract of *H. lantanaefolia*.⁹⁷ We found that the ethanol extract of *H. lantanaefolia* has high cytotoxic activity and the bioassay-directed fractionation afforded semi-purified chromatographic fractions LHL-14 and LHL-15 both of which had extremely high activities on different cell lines (IC_{50} is lower than 50 ng/mL) (Figure 17). The fractions LHL-14 and LHL-15 are currently under investigation.

Figure 17. Cytotoxic Activity of Fractions LHL-14 and LHL-15

4.5 *Bocconia intengrifolia* (Papaveraceae)

Bocconia is a small genus in the *Papaveraceae* family, representing 10 species,⁹⁸ but up to now there has been no investigation conducted on *B. intengrifolia*. Our collaborators in Peru (Professor Vaisberg's group at UPCH) collected this plant species and found that the ethanol extract had very good cytotoxic activity. Bioassay directed fractionation allowed us to find the most active chromatographic fraction PBI-13, which is currently under investigation.

4.6 Other Plant Species Studied

We have active fractions of the following plant extracts: *Humiria balsamifera* (Humiriaceae), *Rollinia andicola* (Annonaceae), *Hippeastrum puniceum* (Amaryllidaceae), *Aniba riparia* (Lauraceae), and I am instructing other members in Dr. Hammond's group on how to purify them further. Other plant extracts lost their bioactivity during separation, and we failed to isolate any bioactive compounds. These plants are: *Terminalia Amazonia* (Combretaceae), *Piper concinnatoris* (missing), *Paullinia* (missing) (Sapindaceae), *Plukenetia volubilis* (Euphorbiaceae) and *Scutia sp.* (Phammaceae). Their bioactivity loss may be due to two factors: first, the plants were collected from the late 1990's to early 2000's; the active components could have decomposed during the long storage period. Second, the cytotoxic activity we determined could have come from a synergistic effect of different components, and therefore the activity was lost during purification.

4.7. Summary

We have isolated three novel antiproliferative withanolides with an unusual carbon framework, namely, physangulidines A, B and C, isolated from *Physalis angulata* L. using a bioassay-directed isolation technique, and their structures were confirmed by NMR spectroscopy and X-ray crystallography techniques. All three withanolides showed significant antiproliferative activity ($GI_{50} < 4.0 \mu\text{g/mL}$) on DU145 and RWPE-1 cells. Compared to 5-fluorouracil, physangulidine A has less antiproliferative activity against nonmalignant 3T3 cell and more antiproliferative activity against HT-29 and K562 cells. The antiproliferative mechanism of physangulidine A is currently under investigation.

Similarly, ($\pm$)-*trans*-dehydrodiisoeugenol was isolated from *Crematosperma microcarpum*, as the main cytotoxic and identified by comparison of its NMR data with a literature report. *Hyptis lantanaefolia* and *Bocconia intengrifolia* have been found to contain cytotoxic agents; LHL-14, LHL-15 and PBI-13 are the most cytotoxic fractions. We are working on LHL-14, LHL-15 and PBI-13 to find pure cytotoxic compound(s).

4.8. Experimental Section

General

Plant material was collected in the Aguaruna region of northeastern Perú, from the late 1990's to early 2000's. Methanol for HPLC separation was HPLC grade, and the rest of the solvents were ACS-grade. Column chromatography was carried out using silica gel (230-400 mesh, Silicycle; Quebec, Canada) and octadecyl-functionalized silica gel (C18, 200-400 mesh, Aldrich; St. Louis, MO, USA). HPLC was performed with a Waters 600E equipped with rheodyne injector and a PDA detector, using a Nova-Pak column (Waters, 6 μ m, 7.8x300 mm, flow rate 1.0 mL/min) or a SymmetryPrep C₁₈ column (Waters, 7 μ m, 300x19 mm, flow rate 4.0 mL/min) and water and methanol as solvent system. NMR spectra of physangulidines A, B and C were collected at 25.0 °C in CDCl₃ at 699.81 MHz in a 5 mm ¹H(¹³C/¹⁵N) (¹³C enhanced) Cold Probe on a VNMRs700 Varian (Agilent) Spectrometer. ¹H NMR spectra of physangulidines A, B and C were recorded at 700 MHz, while their ¹³C NMR spectra were recorded at 175 MHz. UV spectra of physangulidines A and B were recorded in methanol by a Varian Cary 50 Bio UV-Visible Spectrophotometer. FTIR spectra of physangulidines A and B were recorded in ATR (attenuated total reflection) solid mode using a Perkin Elmer Spectrum 100. Melting

points of physangulidines A and B were measured in a DigiMelt PMA 160 melting point apparatus. Crystallographic data of physangulidines A and B were collected by an Agilent Technologies/Oxford Diffraction Gemini CCD diffractometer at 293 K using CuK α radiation. Crystallographic data of physangulidines A (CCDC-858987) and B (CCDC-858988) can be obtained free of charge from the Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

NMR spectra of ($\pm$)-*trans*-dehydrodiisoeugenol were collected at 25.0 °C in CDCl₃ in a Varian Inova 400 MHz Spectrometer. ¹H NMR spectrum of ($\pm$)-*trans*-dehydrodiisoeugenol was recorded at 400 MHz, while its ¹³C NMR spectrum was recorded at 100 MHz. Cytotoxic activity of *Physalis angulata*, fractions and physangulidine A, B and C were evaluated by MTT assay. Cytotoxic activity of other plant extracts was evaluated using the sulforhodamine B (SRB) assay method.

The MTT assay

The DU145 and RWPE-1 cell lines were purchased from American Type Tissue Collection (ATCC, Manassas, VA). The DU145 cell line represents human hormone-insensitive metastatic prostate cancer cells and RWPE-1 is derived from histologically normal human prostate epithelial cells that were immortalized by transfection with HPV-18 DNA. Cells were maintained in standard culture in a humidified incubator at 37°C with 5% CO₂. DU145 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (GIBCO, Grand Island, NY) supplemented with 10% heat-inactivated fetal bovine serum (FBS, 15 min at 65°C, GIBCO) and 1% penicillin/streptomycin (10,000 units each per 1 mL; GIBCO). RWPE-1 cells were cultured in Keratinocyte SFM medium (GIBCO) containing Human Keratinocyte Growth Supplement (GIBCO). For

antiproliferative activity assays, cells were plated in 96 well plates (1,000 cells per well) and were treated the following day by addition of the experimental fraction or vehicle as control. Plant extracts or fractions were resuspended in dimethylsulfoxide (DMSO) to give a concentration of 2 mg/mL and then diluted in cell culture medium then added to triplicate wells to give final concentrations of 4, 2, 1, 0.5, 0.25, or 0 μ g/mL. Cells were cultured for 5 days and then relative cell number was determined using 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, as described previously.⁹⁹ The GI₅₀ value (the concentration required for 50% inhibition of cell proliferation) was estimated from the resulting dose response graphs.

The Sulforhodamine B (SRB) assay method¹⁰⁰

To determine the cytotoxicity of the compounds, cells were plated into 96-well tissue culture plates and in their corresponding growth medium at approximately 10% confluency (BALB/3T3 at 3,500 cells/well, Vero at 2,500 cells/well, H460 at 1,500 cells/well, M-14 at 5,000 cells/well, DU145 at 3,500 cells/well, MCF-7 at 5,000 cells/well, HT-29 at 3,000 cells/well, and K562 at 4,000 cells/well), and incubated at 37°C in a 5% CO₂ and 95 % air humidified atmosphere for 24 h to allow cells to attach. A plate containing each of these cells was fixed in situ with trichloroacetic acid (TCA) in order to obtain the cell values at zero time before adding the compounds. The rest of the plates containing the different cell lines received serial dilutions of the compound to be tested at the following final concentrations: 62.5, 15.625, 3.90625, and 0.97656 μ g/mL. The plates were then incubated at 37 °C in a 5% CO₂ and 95 % air humidified atmosphere for 48 h. The assay was terminated by the addition of cold TCA. TCA treated plates were incubated at 4 °C for 1 hour and then washed five times with tap water to

remove TCA and air dried. Background optical densities were measured in wells incubated with growth medium without cells. TCA-fixed cells were stained for 20 minutes with 0.4% (w/v) SRB dissolved in 1% acetic acid. At the end of the staining period unbound dye was removed by washing four times with 1% acetic acid. After air drying the plates, bound dye was solubilized with 10 mM Tris base (pH 10.5) and the absorbance read on an automated plate reader at a wavelength of 550 nm. The GI₅₀ value was defined as the concentration of test sample resulting in a 50% reduction of absorbance as compared with untreated controls that received a serial dilution of the solvent in which the test samples were dissolved, and was determined by linear regression analysis.

For K562 cells, which grow in suspension, instead of fixing and staining with SRB, cells were counted using a Coulter counter.

Physalis angulata

Leaves, stem and fruits of *Physalis angulata* L were collected in Rio Domingosa, Kuith/Barranquita, Amazonas Department, Northeastern Perú, and identified by the team led by Walter Lewis (voucher number: 18597, extract number: 1159) in April 3, 2002. 46 g of the dried and grounded material was extracted by ethanol (1 L, 95%) at room temperature for 7 days to give 6.26 g extract, which was denoted as SPA. An aliquot (1.5 g) of SPA was partitioned between dichloromethane (40 mL) and water (40 mL), which afforded two fractions, namely organic (SPA1, 500 mg) and aqueous (SPA2, 600 mg) fraction (Bioassay directed fractionation, Scheme 21). In vitro cytotoxic study showed that organic fraction (SPA1) is active (Figure 18), and which was separated on silica gel chromatography and eluted by hexane, ethyl acetate and methanol in gradient mode to

give further four more fractions, SPA11 (15 mg), SPA12 (117 mg), SPA13 (75 mg) and SPA14 (275 mg). Then the most active SPA14 (Figure 19) was subjected to the same chromatography separation as above to further give four fractions: SPA14-1 (15 mg), SPA14-2 (27 mg), SPA14-3 (97 mg) and SPA14-4 (130 mg). Unfortunately, we failed to isolate any pure compound from SPA14-1, SPA14-2 and SPA14-4. SPA14-3 (30 mg) was separated on HPLC reverse phase C18 column eluted in gradient mode of methanol and water (from 80% methanol to 100% methanol) and detected at 254 nm, yielding SPA14-3-1 (0 mg), SPA14-3-2 (1.2 mg) (physangulidine B), SPA14-3-3 (4.1 mg) (physangulidine C) and SPA14-3-4 (18 mg) (physangulidine A). All three of them showed cytotoxic activity against DU145 and RWPE-1 (Figure 20). Their structures were determined by NMR spectra analysis and X-ray crystallography.

Scheme 21. Bioassay Directed Fractionation of *P. angulata* L

Figure 18. Cytotoxic Activity of SPA, SPA1 and SPA2

Figure 19. Cytotoxic Activity of SPA11, SPA12, SPA13 and SPA14

Figure 20. Cytotoxic Activity of SPA14-3-2, SPA14-3-3 and SPA14-3-4

Crematosperma microcarpum

Leaflets and stems of *C. microcarpum* were collected at Copallin, province of Bagua, Amazonas Department, Perú, in May 1999 (voucher number: 19759, extract number: 2967). Air-dried and ground leaflets and stems of *C. microcarpum* (50 g) were extracted with ethanol (1 L, 95%) at room temperature for 7 days. The ethanolic solution was concentrated to dryness under reduced pressure to yield 3.66 g of extract, denoted as ACM. An aliquot (2.0 g) was partitioned between dichloromethane (45 mL) and water (45 mL), which afforded an organic fraction (ACM1, 1240 mg), insoluble residue

(ACM2, 100 mg) and an aqueous fraction (ACM3, 108 mg). *In vitro* cytotoxic studies indicated that ACM1 was the only active fraction. Hence, it was purified on silica gel chromatography and eluted with hexanes, ethyl acetate and methanol in gradient mode to yield five sub-fractions: ACM11 (163 mg), ACM12 (165 mg), ACM13 (650 mg), ACM14 (123 mg) and ACM15 (110 mg). An aliquot of ACM13 (10 mg) was the most cytotoxic fraction, and An aliquot (10 mg) was further chromatographed using a reverse phase C18 column in a solvent gradient of methanol-water (80%, V/V) to pure methanol, yielding ($\pm$)-*trans*-dehydrodiisoeugenol (7.0 mg) (Scheme 22).

Scheme 22. Bioassay Directed Fractionation of *C. microcarpum*

Hyptis lantanaefolia

Leaves, stem and flower of *H. lantanaefolia* were collected at road from Chiple to Cutervo, province of Jaén, Cajamarca Department, Perú in Oct 6, 2005. (Voucher number: 19547, extract number: 2352). Dried leaflets, stems and flower of *H. lantanaefolia* (70 g) were extracted with ethanol (1L, 95%) at room temperature for 7 days. The ethanolic solution was concentrated to dryness under reduced pressure to yield the extract (5.12 g), denoted as LHL. An aliquot of LHL (4.0 g) was partitioned between dichloromethane (60mL) and water (65 mL), affording an organic fraction (LHL1, 840 mg) and an aqueous fraction (LHL2, 2.0 g) (Scheme 23). LHL1 was more active on different cell

Scheme 23. Bioassay Directed Fractionation of *H. lantanaefolia*

lines than LHL2, and LHL1 was further subjected to silica gel chromatographic separation to give six fractions: LHL11 (41 mg), LHL12 (150 mg), LHL13 (75 mg), LHL14 (105 mg), LHL15 (145 mg) and LHL16 (66 mg). Of all fractions LHL14 and LHL15 were extremely high in vitro cytotoxic activities. We are currently working on both LHL14 and LHL15 to isolate active agents.

Bocconia intengrifolia

Bark of *B. intengrifolia* was collected on the road from Santo Domingo de la Capilla to San Pedro, province of Cutervo, Cajamarca Department, Perú (Voucher number: 19644, extract number: 2367). Dried and grounded bark of *B. intengrifolia* (50) was extracted by ethanol (1L, 95%) at room temperature for 7 days to give crude extract (4.47 g) after dryness of ethanol. An aliquot of PBI (2.3 g) was partitioned between dichloromethane (50 mL) and water (50 mL), affording an organic fraction (PBI1, 181 mg) and an aqueous fraction (PBI2, 1.5 g) (Scheme 24). *In vitro* cytotoxicity study showed that the organic fraction PBI1 is active; following fractionation of PBI1 on silica gel furnished four fractions: PBI11 (13 mg), PBI12 (43 mg), PBI13 (119 mg) and PBI14 (7 mg), of which PBI13 is most active fraction. PBI13 is currently under investigation.

Scheme 24. Bioassay Directed Fractionation of *B. intengrifolia*

Ongoing plant studies

Bark and leaves of *Humiria balsamifera* (Humiriaceae) were collected in the trail to Quebrada Sunsunza, El Porvenir, Aramango, province of Bagua, Amazonas department, Perú (Voucher number: 19478, extract number: 2278). Dried and grounded bark and leaves of *H. balsamifera* (50 g) were extracted with ethanol (1L, 95%) to give crude extract (6.69 g) after filtration and dryness of ethanol solution. Then, the extract was subjected to simple partition between water and dichloromethane; the bioactive organic layer is currently under study.

Fruits of *Rollinia andicola* (Annonaceae) were collected at El Tigre, Cambio-Pitec, Copallín, province of Bagua, Amazonas Department, Perú (Voucher number: 19877, extract number: 2900). Dried and grounded flowers of *R. andicola* (61 g) were extracted with ethanol (1L, 95%) to give crude extract (7.71 g). The bioactive organic layer is currently under study.

Bulbs of *Hippeastrum puniceum* (Amaryllidaceae) were collected at Kunchin, Imaza, province of Bagua, Amazonas Department, Perú in Aug 9, 2001 (Voucher number: 17506, extract number: 435). Dried and grounded *H. puniceum* (43 g) was extracted with ethanol (1L, 95%) to give crude extract (3.36 g). Bioactive organic layer is currently under investigation.

The bark of *Aniba riparia* (Lauraceae) was collected at Wichim, Imaza, province of Bagua, Amazonas Department, Perú in Nov 26, 2002 (Voucher number: 19366, extract number: 2239). Dried and grounded bark of *A. riparia* (50 g) was extracted with ethanol (1L, 95%) to yield crude extract (4.13 g) after dryness. The bioactive organic layer is currently under investigation.

REFERENCES

1. (a) Amblard, F.; Cho, J. H.; Schinazi, R. F., Cu(I)-Catalyzed Huisgen Azide-Alkyne 1,3-Dipolar Cycloaddition Reaction in Nucleoside, Nucleotide, and Oligonucleotide Chemistry. *Chem. Rev.* **2009**, *109*, 4207-4220; (b) Harvey, D. F.; Sigano, D. M., Carbene-Alkyne-Alkene Cyclization Reactions. *Chem. Rev.* **1996**, *96*, 271-288; (c) Meldal, M.; Tornøe, C. W., Cu-Catalyzed Azide-Alkyne Cycloaddition. *Chem. Rev.* **2008**, *108*, 2952-3015; (d) Willis, M. C., Transition Metal Catalyzed Alkene and Alkyne Hydroacylation. *Chem. Rev.* **2009**, *110*, 725-748.
2. (a) Han, J.; Xu, B.; Hammond, G. B., Synthesis of α -CN and α -CF₃ N-Heterocycles through Tandem Nucleophilic Additions. *Org. Lett.* **2011**, *13*, 3450-3453; (b) Liu, L.-P.; Hammond, G. B., Highly Efficient and Tunable Synthesis of Dioxabicyclo[4.2.1] Ketals and Tetrahydropyrans via Gold-Catalyzed Cycloisomerization of 2-Alkynyl-1,5-diols. *Org. Lett.* **2009**, *11*, 5090-5092; (c) Liu, L.-P.; Hammond, G. B., Gold-Catalyzed Cascade Annulations of 2-(Ynol)aryl Aldehydes: Facile Synthesis of Benzochromanes and Benzobicyclo[n.3.1]acetals. *Org. Lett.* **2010**, *12*, 4640-4643; (d) Liu, L.-P.; Xu, B.; Hammond, G. B., Michael Addition of Allenates to Electron-Deficient Olefins: Facile Synthesis of 2-Alkynyl-Substituted Glutaric Acid Derivatives. *Org. Lett.* **2008**, *10*, 3887-3890; (e) Wang, W.; Xu, B.; Hammond, G. B., Synthesis of Functionalized α,α -Disubstituted β -Alkynyl Esters from Allenates through an Alkynyleneolate Intermediate. *Org. Lett.* **2008**, *10*, 3713-3716; (f) Yang, H.; Xu, B.; Hammond, G. B., Highly Regioselective Fluorination and Iodination of Alkynyl Enolates. *Org. Lett.* **2008**, *10*, 5589-5591.
3. Wang, W.; Jasinski, J.; Hammond, G. B.; Xu, B., Fluorine-Enabled Cationic Gold Catalysis: Functionalized Hydration of Alkynes. *Angew. Chem. Int. Ed.* **2010**, *49*, 7247-7252.
4. Jin, Z.; Aponte, J. C.; Vaisberg, A.; Sauvain, M.; Gilman, R. H.; Hammond, G. B. In *Antileishmanial, trypanocidal, antituberculosis and cytotoxic activities of targeted plant species from the Peruvian rain forest*, American Chemical Society: 2010; pp AGFD-202.
5. Newman, D. J.; Cragg, G. M., Natural Products as Sources of New Drugs over the Last 25 Years. *J. Nat. Prod.* **2007**, *70*, 461-477.
6. Aponte, J. C.; Vaisberg, A. J.; Rojas, R.; Sauvain, M.; Lewis, W. H.; Lamas, G.; Sarasara, C. s.; Gilman, R. H.; Hammond, G. B., A Multipronged Approach to the Study of Peruvian Ethnomedicinal Plants: A Legacy of the ICBG-Peru Project Δ . *J. Nat. Prod.* **2009**, *72*, 524-526.

7. (a) Aponte, J. C.; Estevez, Y.; Gilman, R. H.; Lewis, W. H.; Rojas, R.; Sauvain, M.; Vaisberg, A. J.; Hammond, G. B., Anti-infective and cytotoxic compounds present in *Blepharodon nitidum*. *Planta Med.* **2008**, *74*, 407-410; (b) Aponte, J. C.; Jin, Z.; Vaisberg, A. J.; Castillo, D.; Malaga, E.; Lewis, W. H.; Sauvain, M.; Gilman, R. H.; Hammond, G. B., Cytotoxic and anti-infective phenolic compounds isolated from *Mikania decora* and *Crematosperma microcarpum*. *Planta Med.* **2011**, *77*, 1597-1599; (c) Aponte, J. C.; Vaisberg, A. J.; Rojas, R.; Caviedes, L.; Lewis, W. H.; Lamas, G.; Sarasara, C.; Gilman, R. H.; Hammond, G. B., Isolation of Cytotoxic Metabolites from Targeted Peruvian Amazonian Medicinal Plants. *J. Nat. Prod.* **2007**, *71*, 102-105; (d) Aponte, J. C.; Yang, H.; Vaisberg, A. J.; Castillo, D.; Malaga, E.; Verastegui, M.; Casson, L. K.; Stivers, N.; Bates, P. J.; Rojas, R.; Fernandez, I.; Lewis, W. H.; Sarasara, C.; Sauvain, M.; Gilman, R. H.; Hammond, G. B., Cytotoxic and anti-infective sesquiterpenes present in *Plagiochila disticha* (plagiochilaceae) and *Ambrosia peruviana* (asteraceae). *Planta Med.* **2010**, *76*, 705-707.
8. (a) Aponte, J. C.; Castillo, D.; Estevez, Y.; Gonzalez, G.; Arevalo, J.; Hammond, G. B.; Sauvain, M., In vitro and in vivo anti-Leishmania activity of polysubstituted synthetic chalcones. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 100-103; (b) Aponte, J. C.; Vaisberg, A. J.; Castillo, D.; Gonzalez, G.; Estevez, Y.; Arevalo, J.; Quiliano, M.; Zimic, M.; Verastegui, M.; Malaga, E.; Gilman, R. H.; Bustamante, J. M.; Tarleton, R. L.; Wang, Y.; Franzblau, S. G.; Pauli, G. F.; Sauvain, M.; Hammond, G. B., Trypanoside, anti-tuberculosis, leishmanicidal, and cytotoxic activities of tetrahydrobenzothienopyrimidines. *Bioorg. Med. Chem.* **2010**, *18*, 2880-2886; (c) Aponte, J. C.; Verastegui, M.; Malaga, E.; Zimic, M.; Quiliano, M.; Vaisberg, A. J.; Gilman, R. H.; Hammond, G. B., Synthesis, Cytotoxicity, and Anti-Trypanosoma cruzi Activity of New Chalcones. *J. Med. Chem.* **2008**, *51*, 6230-6234.
9. Cremlyn, R. J. W., *An introduction to organosulfur chemistry*. Wiley: Chichester ; New York, 1996; p xii, 250 p.
10. (a) Cao, C.; Fraser, L. R.; Love, J. A., Rhodium-Catalyzed Alkyne Hydrothiolation with Aromatic and Aliphatic Thiols. *J. Am. Chem. Soc.* **2005**, *127*, 17614-17615; (b) Kondo, T.; Mitsudo, T.-a., Metal-Catalyzed Carbon-Sulfur Bond Formation. *Chem. Rev.* **2000**, *100*, 3205-3220; (c) Ogawa, A.; Ikeda, T.; Kimura, K.; Hirao, T., Highly Regio- and Stereocontrolled Synthesis of Vinyl Sulfides via Transition-Metal-Catalyzed Hydrothiolation of Alkynes with Thiols. *J. Am. Chem. Soc.* **1999**, *121*, 5108-5114; (d) Weiss, C. J.; Wobser, S. D.; Marks, T. J., Organoactinide-Mediated Hydrothiolation of Terminal Alkynes with Aliphatic, Aromatic, and Benzylic Thiols. *J. Am. Chem. Soc.* **2009**, *131*, 2062-2063.
11. (a) Beckerle, K.; Manivannan, R.; Lian, B.; Meppelder, G.-J. M.; Raabe, G.; Spaniol, T. P.; Ebeling, H.; Pelascini, F.; Muelhaupt, R.; Okuda, J., Stereospecific styrene enchainment at a titanium site within a helical ligand framework: evidence for the formation of homochiral polystyrene. *Angew. Chem., Int. Ed.* **2007**, *46*, 4790-4793; (b) Cohen, A.; Yeori, A.; Goldberg, I.; Kol, M., Group 4 Complexes of a New [OSSO]-Type Dianionic Ligand. Coordination Chemistry and Preliminary Polymerization Catalysis Studies. *Inorg. Chem.* **2007**, *46*, 8114-

8116; (c) Gall, B. T.; Pelascini, F.; Ebeling, H.; Beckerle, K.; Okuda, J.; Muelhaupt, R., Molecular Weight and End Group Control of Isotactic Polystyrene Using Olefins and Nonconjugated Diolefins as Chain Transfer Agents. *Macromolecules (Washington, DC, U. S.)* **2008**, *41*, 1627-1633; (d) Ishii, A.; Toda, T.; Nakata, N.; Matsuo, T., Zirconium Complex of an [OSSO]-Type Diphenolate Ligand Bearing trans-1,2-Cyclooctanediylbis(thio) Core: Synthesis, Structure, and Isospecific 1-Hexene Polymerization. *J. Am. Chem. Soc.* **2009**, *131*, 13566-13567; (e) Lian, B.; Beckerle, K.; Spaniol, T. P.; Okuda, J., Regioselective 1-hexene oligomerization using cationic bis(phenolato) group 4 metal catalysts: switch from 1,2- to 2,1-insertion. *Angew. Chem., Int. Ed.* **2007**, *46*, 8507-8510.

12. (a) Caserio, M. C.; Fisher, C. L.; Kim, J. K., Boron trifluoride catalyzed addition of disulfides to alkenes. *J. Org. Chem.* **1985**, *50*, 4390-4393; (b) Kitamura, T.; Matsuyuki, J.; Taniguchi, H., 1,2-Disulfenylation of alkenes induced by a hypervalent iodine(III) reagent [PhIO-TfOH]. *J. Chem. Soc., Perkin Trans. 1* **1991**, 1607-1608; (c) Kondo, T.; Uenoyama, S.-y.; Fujita, K.-i.; Mitsudo, T.-a., First Transition-Metal Complex Catalyzed Addition of Organic Disulfides to Alkenes Enables the Rapid Synthesis of vicinal-Dithioethers. *J. Am. Chem. Soc.* **1999**, *121*, 482-483; (d) Nishimura, T.; Yoshinaka, T.; Uemura, S., Metal cation-exchanged montmorillonite-catalyzed addition of organic disulfides to alkenes. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 1138-1141; (e) Usugi, S.; Yorimitsu, H.; Shinokubo, H.; Oshima, K., Disulfidation of alkynes and alkenes with gallium trichloride. *Org. Lett.* **2004**, *6*, 601-603; (f) Yamagiwa, N.; Suto, Y.; Torisawa, Y., Convenient method for the addition of disulfides to alkenes. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6197-6201.

13. (a) Alonso, F.; Beletskaya, I. P.; Yus, M., Transition-metal-catalyzed addition of heteroatom-hydrogen bonds to alkynes. *Chem. Rev.* **2004**, *104*, 3079-3159; (b) Amin, S. B.; Marks, T. J., Versatile pathways for in situ polyolefin functionalization with heteroatoms: catalytic chain transfer. *Angew. Chem., Int. Ed.* **2008**, *47*, 2006-2025; (c) Hartwig, J. F., Carbon-heteroatom bond formation catalyzed by organometallic complexes. *Nature (London, U. K.)* **2008**, *455*, 314-322.

14. (a) Blomquist, A. T.; Wolinsky, J., Addition of ethyl mercaptan to acetylenic compounds. *J. Org. Chem.* **1958**, *23*, 551-554; (b) Demchuk, D. V.; Lazareva, M. I.; Lindeman, S. V.; Khrustalyov, V. N.; Struckhov, Y. T.; Ismagilov, R. F.; Troyansky, E. I.; Nikishin, G. I., Synthesis of 12- and 13-membered sulfur-containing lactones by homolytic macrocyclization of mercaptoacetic esters with alkynes. *Synthesis* **1995**, 307-311; (c) Il'yasov, E. A.; Galust'yan, G. G., Homolytic addition of 1-alkanethiols to 5-ethynyl-2-methylpyridine. *Chem. Heterocycl. Compd.* **2000**, *35*, 1187-1189; (d) Stacey, F. W.; Harris, J. F., Jr., Radiation-induced addition of hydrogen sulfide to substituted acetylenes. Synthesis of vinylthiols. *J. Am. Chem. Soc.* **1963**, *85*, 963-965; (e) Troyanskii, E. I.; Lazareva, M. I.; Demchuk, D. V.; Samoshin, V. V.; Strelenko, Y. A.; Nikishin, G. I., Free radical intermolecular macrocyclization: an extremely facile route to 12-membered thiacyclic ethers. *Synlett* **1992**, 233-234; (f) Troyansky, E. I.; Ismagilov, R. F.; Strelenko, Y. A.; Samoshin, V. V.; Demchuk, D. V.; Nikishin, G.

- I.; Lindeman, S. V.; Khrustalyov, V. V.; Struchkov, Y. T., Remote asymmetric induction in free radical cycloaddition leading to trans-cyclohexano-fused 12-membered crown thioethers. *Tetrahedron Lett.* **1995**, *36*, 2293-2294; (g) Yamagishi, K.; Tanaka, T.; Hoshino, T., Photochemical addition reactions. IV. Addition of thioacetic acid to propargyl alcohol and some propargyl esters. *Bull. Chem. Soc. Jpn.* **1957**, *30*, 455-458.
15. (a) Yorimitsu, H.; Wakabayashi, K.; Shinokubo, H.; Oshima, K., Radical addition of 2-iodoalkanamide or 2-iodoalkanoic acid to alkenols using a water-soluble radical initiator in water. A facile synthesis of γ -lactones. *Tetrahedron Lett.* **1999**, *40*, 519-522; (b) Yorimitsu, H.; Wakabayashi, K.; Shinokubo, H.; Oshima, K., Radical addition of 2-iodoalkanamide or 2-iodoalkanoic acid to alkenes with a water-soluble radical initiator in aqueous media: facile synthesis of γ -lactones. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 1963-1970.
16. (a) Chan, J. W.; Yu, B.; Hoyle, C. E.; Lowe, A. B., Convergent synthesis of 3-arm star polymers from RAFT-prepared poly(N,N-diethylacrylamide) via a thiol-ene click reaction. *Chem. Commun. (Cambridge, U. K.)* **2008**, 4959-4961; (b) Killops, K. L.; Campos, L. M.; Hawker, C. J., Robust, Efficient, and Orthogonal Synthesis of Dendrimers via Thiol-ene "Click" Chemistry. *J. Am. Chem. Soc.* **2008**, *130*, 5062-5064.
17. (a) Almutairi, A.; Guillaudeu, S. J.; Berezin, M. Y.; Achilefu, S.; Frechet, J. M. J., Biodegradable pH-sensing dendritic nanoprobe for near-infrared fluorescence lifetime and intensity imaging. *J. Am. Chem. Soc.* **2008**, *130*, 444-445; (b) Grayson, S. M.; Frechet, J. M., Convergent dendrons and dendrimers: from synthesis to applications. *Chem. Rev.* **2001**, *101*, 3819-3868; (c) Helms, B.; Meijer, E. W., Chemistry. Dendrimers at work. *Science* **2006**, *313*, 929-930.
18. (a) Fokin, V. V.; Sharpless, K. B., A practical and highly efficient aminohydroxylation of unsaturated carboxylic acids. *Angew. Chem., Int. Ed.* **2001**, *40*, 3455-3457; (b) Narayan, S.; Muldoon, J.; Finn, M. G.; Fokin, V. V.; Kolb, H. C.; Sharpless, K. B., "On water": Unique reactivity of organic compounds in aqueous suspension. *Angew. Chem., Int. Ed.* **2005**, *44*, 3275-3279.
19. Yadav, J. S.; Reddy, B. V. S.; Raju, A.; Ravindar, K.; Baishya, G., Hydrothiolation of unactivated alkynes catalyzed by indium(III) bromide. *Chem. Lett.* **2007**, *36*, 1474-1475.
20. Ionita, P., Is DPPH stable free radical a good scavenger for oxygen active species? *Chem. Pap.* **2005**, *59*, 11-16.
21. (a) Ranu, B. C.; Banerjee, S.; Jana, R., Ionic liquid as catalyst and solvent: the remarkable effect of a basic ionic liquid, [bmim]OH on Michael addition and alkylation of active methylene compounds. *Tetrahedron* **2006**, *63*, 776-782; (b) Santos, L. L.; Ruiz, V. R.; Sabater, M. J.; Corma, A., Regioselective transformation of alkynes into cyclic acetals and thioacetals with a gold(I) catalyst: comparison with Bronsted acid catalysts. *Tetrahedron* **2008**, *64*, 7902-7909; (c) Xu, C.; Bartley, J. K.; Enache, D. I.; Knight, D. W.; Lunn, M.; Lok, M.; Hutchings,

G. J., On the synthesis of β -keto-1,3-dithianes from conjugated ynones catalyzed by magnesium oxide. *Tetrahedron Lett.* **2008**, 49, 2454-2456.

22. Ranu, B. C.; Mandal, T., Water-promoted highly selective anti-Markovnikov addition of thiols to unactivated alkenes. *Synlett* **2007**, 925-928.

23. (a) Andriyankova, L. V.; Zhivet'ev, S. A.; Mal'kina, A. G.; Kudryakova, P. N.; Kositsyna, E. I.; Il'icheva, L. N.; Ushakov, I. A.; Afonin, A. V.; Trofimov, B. A., Cyanoacetylene and its derivatives: XXXI. Nucleophilic addition of 2- and 4-mercaptopyridines to cyanacetylenes. *Russ. J. Org. Chem.* **2002**, 38, 1681-1685; (b) Bierer, D. E.; Dener, J. M.; Dubenko, L. G.; Gerber, R. E.; Litvak, J.; Peterli, S.; Peterli-Roth, P.; Truong, T. V.; Mao, G.; Bauer, B. E., Novel 1,2-Dithiins: Synthesis, Molecular Modeling Studies, and Antifungal Activity. *J. Med. Chem.* **1995**, 38, 2628-48; (c) Kondoh, A.; Takami, K.; Yorimitsu, H.; Oshima, K., Stereoselective Hydrothiolation of Alkynes Catalyzed by Cesium Base: Facile Access to (Z)-1-Alkenyl Sulfides. *J. Org. Chem.* **2005**, 70, 6468-6473; (d) Koreeda, M.; Wang, Y., Novel Synthesis of 3,6-Disubstituted 1,2-Dithiin Molecules Involving a Direct Oxidative Deprotection-Cyclization Sequence from 1,4-Bis(tert-butylthio)-1,3-butadiene Precursors. *J. Org. Chem.* **1997**, 62, 446-447; (e) Koreeda, M.; Yang, W., Chemistry of 1,2-Dithiins. Synthesis of the Potent Antibiotic Thiarubrine A. *J. Am. Chem. Soc.* **1994**, 116, 10793-10794; (f) Koreeda, M.; Yang, W., The chemistry of 1,2-dithiins: synthesis of 1,2-dithiin and 3,6-disubstituted 1,2-dithiins. *Synlett* **1994**, 201-203; (g) Waters, M. S.; Snelgrove, K.; Maligres, P., 1-Hydroxy-3-phenyl-2-propanone. *Org. Synth.* **2003**, 80, 190-194; (h) Skvortsov, Y. M.; Mal'kina, A. G.; Moshchevitina, E. I.; Modonov, V. B.; Trofimov, B. A., Nucleophilic addition of thiols to 4-hydroxy-2-alkynenitriles and furanization of the adducts. *Sulfur Lett.* **1989**, 9, 141-148; (i) Trzhtsinskaya, B. V.; Abramova, N. D.; Teterina, L. F.; Andriyankova, L. V.; Afonin, A. V., Reaction of 3-indolethiols with acetylenes. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1989**, 416-419; (j) Galambos, G.; Csokasi, P.; Szantay, C., Jr.; Czira, G.; Szantay, C., Addition of thiols to acetylene-substituted indoles. *Heterocycles* **1994**, 38, 1459-1464; (k) Galambos, G.; Csokasi, P.; Szantay, C., Jr.; Szantay, C., Synthesis of 4,5-disubstituted 1H-1,3,4,5-tetrahydrobenz[cd]indole derivatives. *Liebigs Ann./Recl.* **1997**, 1969-1978.

24. Miura, K.; Oshima, K.; Utimoto, K., Triethylborane induced stereoselective radical addition of R_3SiH to acetylenes and stereoselective reduction of alkenyl iodides with tris(trimethylsilyl)silane. *Bull. Chem. Soc. Jpn.* **1993**, 66, 2356-64.

25. http://en.wikipedia.org/wiki/Carbon-sulfur_bond.

26. http://en.wikipedia.org/wiki/Bond_length.

27. Dubbaka, S. R.; Vogel, P., Organosulfur compounds: Electrophilic reagents in transition-metal-catalyzed carbon-carbon bond-forming reactions. *Angew. Chem., Int. Ed.* **2005**, 44, 7674-7684.

28. (a) Sabarre, A.; Love, J., Synthesis of 1,1-Disubstituted Olefins via Catalytic Alkyne Hydrothiolation/Kumada Cross-Coupling. *Org. Lett.* **2008**, 10, 3941-3944; (b) Wenkert, E.; Chianelli, D., Nickel-catalyzed decarbonylation of

thioesters. *J. Chem. Soc., Chem. Commun.* **1991**, 627-628; (c) Wenkert, E.; Michelotti, E. L.; Swindell, C. S., Nickel-induced conversion of carbon-oxygen into carbon-carbon bonds. One-step transformations of enol ethers into olefins and aryl ethers into biaryls. *J. Am. Chem. Soc.* **1979**, *101*, 2246-2247.

29. Ma, S.; Ren, H.; Wei, Q., Highly Regio- and Stereoselective Halohydroxylation Reaction of 1,2-Allenyl Phenyl Sulfoxides. Reaction Scope, Mechanism, and the Corresponding Pd- or Ni-Catalyzed Selective Coupling Reactions. *J. Am. Chem. Soc.* **2003**, *125*, 4817-4830.

30. (a) Hadfield, J. A.; Gaukroger, K.; Hirst, N.; Weston, A. P.; Lawrence, N. J.; McGown, A. T., Synthesis and evaluation of double bond substituted combretastatins. *Eur. J. Med. Chem.* **2005**, *40*, 529-541; (b) Haidoune, M. B.; Raynaud, I.; O'Connor, N.; Richomme, P.; Mornet, R.; Laloue, M., Synthesis and cytokinin activity of new zeatin derivatives. *J. Agric. Food Chem.* **1998**, *46*, 1577-1582; (c) Miyauchi, H.; Nakamura, T.; Ohashi, N., Asymmetric synthesis of SM-9164, a biologically active enantiomer of antifungal agent SM-8668. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 2625-2632; (d) Pinard, E.; Gaudry, M.; Henot, F.; Thellend, A., Asymmetric total synthesis (+)-pisatin, a phytoalexin from garden peas (*Pisum sativum* L.). *Tetrahedron Lett.* **1998**, *39*, 2739-2742; (e) Spino, C.; Gobdout, C., A Stereodivergent Approach to Chiral Nonracemic N-Protected α,α -Dialkylated Amino Acids. *J. Am. Chem. Soc.* **2003**, *125*, 12106-12107.

31. (a) Adam, W.; Alsters, P. L.; Neumann, R.; Saha-Moeller, C. R.; Seebach, D.; Beck, A. K.; Zhang, R., Chiral Hydroperoxides as Oxygen Source in the Catalytic Stereoselective Epoxidation of Allylic Alcohols by Sandwich-Type Polyoxometalates: Control of Enantioselectivity through a Metal-Coordinated Template. *J. Org. Chem.* **2003**, *68*, 8222-8231; (b) Adam, W.; Beck, A. K.; Pichota, A.; Saha-Moller, C. R.; Seebach, D.; Vogl, N.; Zhang, R., Control of enantioselectivity through a hydrogen-bonded template in the vanadium(V)-catalyzed epoxidation of allylic alcohols by optically active hydroperoxides. *Tetrahedron: Asymmetry* **2003**, *14*, 1355-1361; (c) Lattanzi, A.; Piccirillo, S.; Scettri, A., Vanadium-catalyzed asymmetric epoxidation of allylic alcohols mediated by (+)-norcamphor-derived hydroperoxide. *Eur. J. Org. Chem.* **2005**, 1669-1674; (d) Malkov, A. V.; Bourhani, Z.; Kocovsky, P., Amino acid-derived hydroxamic acids as chiral ligands in the vanadium catalyzed epoxidation. *Org. Biomol. Chem.* **2005**, *3*, 3194-3200; (e) Mantilli, L.; Gerard, D.; Torche, S.; Besnard, C.; Mazet, C., Iridium-Catalyzed Asymmetric Isomerization of Primary Allylic Alcohols. *Angew. Chem., Int. Ed.* **2009**, *48*, 5143-5147; (f) Zhang, W.; Basak, A.; Kosugi, Y.; Hoshino, Y.; Yamamoto, H., Enantioselective epoxidation of allylic alcohols by a chiral complex of vanadium: An effective controller system and a rational mechanistic model. *Angew. Chem., Int. Ed.* **2005**, *44*, 4389-4391.

32. Brandsma, L.; Heus-Kloos, Y. A.; Heiden, R. v. d.; Verkruijsse, H. D., *Preparative acetylenic chemistry*. 2nd ed.; Elsevier: Amsterdam ; New York, 1988; p xii, 321 p.

33. (a) Liu, L.-P.; Malhotra, D.; Paton, R. S.; Houk, K. N.; Hammond, G. B., The [4+2], not [2+2], Mechanism Occurs in the Gold-Catalyzed Intramolecular

Oxygen Transfer Reaction of 2-Alkynyl-1,5-Diketones. *Angew. Chem., Int. Ed.* **2010**, *49*, 9132-9135; (b) Xu, B.; Hammond, G. B., Thermodynamically favored aldol reaction of propargyl or allenyl esters: regioselective synthesis of carbinol allenoates. *Angew. Chem., Int. Ed.* **2008**, *47*, 689-692.

34. (a) Di, Y.-T.; He, H.-P.; Lu, Y.; Yi, P.; Li, L.; Wu, L.; Hao, X.-J., Alkaloids from the Leaves of *Daphniphyllum longeracemosum*. *J. Nat. Prod.* **2006**, *69*, 1074-1076; (b) Ferreira, M. A.; Alves, A. C.; Costa, M. A. C.; Paul, M. I., Naphthoquinone dimers and trimers from *Euclea natalensis*. *Phytochemistry* **1977**, *16*, 117-120; (c) Gu, J.-Q.; Graf, T. N.; Lee, D.; Chai, H.-B.; Mi, Q.; Kardono, L. B. S.; Setyowati, F. M.; Ismail, R.; Riswan, S.; Farnsworth, N. R.; Cordell, G. A.; Pezzuto, J. M.; Swanson, S. M.; Kroll, D. J.; Falkinham, J. O., III; Wall, M. E.; Wani, M. C.; Kinghorn, A. D.; Oberlies, N. H., Cytotoxic and Antimicrobial Constituents of the Bark of *Diospyros maritima* Collected in Two Geographical Locations in Indonesia. *J. Nat. Prod.* **2004**, *67*, 1156-1161; (d) Solomon, P. H.; Nakanishi, K.; Fallon, W. E.; Shimizu, Y., Confirmation of 18-norsteroidal structure of fukujusonorones. Sensitivity of CMR[carbon magnetic resonance] to configurational changes. *Chem. Pharm. Bull.* **1974**, *22*, 1671-1673.

35. (a) Jin, T.; Yamamoto, Y., Gold-Catalyzed Intramolecular Carbocyclization of Alkynyl Ketones Leading to Highly Substituted Cyclic Enones. *Org. Lett.* **2007**, *9*, 5259-5262; (b) Jin, T.; Yamamoto, Y., Gold-Catalyzed Synthesis of Polycyclic Enones from Carbon Tethered 1,3-Enynyl Carbonyls via Tandem Heteroenyne Metathesis and Nazarov Reaction. *Org. Lett.* **2008**, *10*, 3137-3139; (c) Jin, T.; Yang, F.; Liu, C.; Yamamoto, Y., TfOH-catalyzed intramolecular alkyne-ketone metathesis leading to highly substituted five-membered cyclic enones. *Chem. Commun. (Cambridge, U. K.)* **2009**, 3533-3535.

36. Lal, G. S.; Pez, G. P.; Syvret, R. G., Electrophilic NF Fluorinating Agents. *Chem. Rev.* **1996**, *96*, 1737-1755.

37. (a) Banks, R. E.; Mohialdin-Khaffaf, S. N.; Lal, G. S.; Sharif, I.; Syvret, R. G., 1-Alkyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane salts: a novel family of electrophilic fluorinating agents. *J. Chem. Soc., Chem. Commun.* **1992**, 595-596; (b) Nyffeler, P. T.; Duron, S. G.; Burkart, M. D.; Vincent, S. P.; Wong, C.-H., Selectfluor: Mechanistic insight and applications. *Angew. Chem., Int. Ed.* **2005**, *44*, 192-212.

38. Hart, J. J.; Syvret, R. G., Industrial scale production of Selectfluor fluorination agent: from initial concept to full scale commercial production in a 5 year period. *J. Fluorine Chem.* **1999**, *100*, 157-161.

39. (a) Singh, R. P.; Shreeve, J. n. M., Recent Highlights in Electrophilic Fluorination with 1-Chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane Bis(tetrafluoroborate). *Acc. Chem. Res.* **2004**, *37*, 31-44; (b) Stavber, S.; Zupan, M., Selectfluor F-TEDA-BF₄ as a versatile mediator or catalyst in organic chemistry. *Acta Chim. Slov.* **2005**, *52*, 13-26.

40. (a) Laali, K. K.; Borodkin, G. I., First application of ionic liquids in electrophilic fluorination of arenes; Selectfluor (F-TEDA-BF₄) for "green"

fluorination. *J. Chem. Soc., Perkin Trans. 2* **2002**, 953-957; (b) Takeuchi, Y.; Tarui, T.; Shibata, N., A Novel and Efficient Synthesis of 3-Fluorooxindoles from Indoles Mediated by Selectfluor. *Org. Lett.* **2000**, 2, 639-642.

41. (a) Chambers, R. D.; Kenwright, A. M.; Parsons, M.; Sandford, G.; Moilliet, J. S., Elemental fluorine. Part 14.1 Electrophilic fluorination and nitrogen functionalization of hydrocarbons. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2190-2197; (b) Chambers, R. D.; Parsons, M.; Sandford, G.; Bowden, R., Electrophilic fluorination at saturated sites. *Chem. Commun.* **2000**, 959-960; (c) Ladame, S.; Willson, M.; Perie, J., A convenient synthesis of dibenzyl α,α -difluoromethyl- β -ketophosphonates. *Eur. J. Org. Chem.* **2002**, 2640-2648.

42. (a) Majumder, U.; Armantrout, J. R.; Williams, R. V.; Shreeve, J. n. M., N,N-Difluorotris(tert-butyl)silylamine - The First Organosilyl Difluoroamine. Synthesis and Computational Studies. *J. Org. Chem.* **2002**, 67, 8435-8439; (b) Singh, R. P.; Shreeve, J. n. M., The first application of Selectfluor in electrophilic fluorination of amines: a new route to -NF₂, -NHF, and >NF compounds. *Chem. Commun.* **2001**, 1196-1197.

43. (a) Greedy, B.; Gouverneur, V., Fluorodesilylation of alkenyltrimethylsilanes: a new route to fluoroalkenes and difluoromethyl-substituted amides, alcohols or ethers. *Chem. Commun.* **2001**, 233-234; (b) Lal, G. S., Site-selective fluorination of organic compounds using 1-alkyl-4-fluoro-1,4-diazabicyclo[2.2.2]octane salts (selectfluor reagents). *J. Org. Chem.* **1993**, 58, 2791-2796; (c) Stavber, S.; Sotler, T.; Zupan, M., A mild, selective method for preparation of vicinal fluoro ethers using "F-Teda-BF₄". *Tetrahedron Lett.* **1994**, 35, 1105-1108; (d) Stavber, S.; Sotler-Pecan, T.; Zupan, M., Stereochemistry and some kinetic aspects of fluorination of phenyl-substituted alkenes with Selectfluor reagent F-TEDA-BF₄. *Bull. Chem. Soc. Jpn.* **1996**, 69, 169-175.

44. (a) Cahard, D.; Audouard, C.; Plaquevent, J. C.; Toupet, L.; Roques, N., N-Fluorocinchonidinium tetrafluoroborate F-CD-BF₄: purification and structure elucidation of this novel enantioselective electrophilic fluorinating agent. *Tetrahedron Lett.* **2001**, 42, 1867-1869; (b) Cahard, D.; Audouard, C.; Plaquevent, J.-C.; Roques, N., Design, Synthesis, and Evaluation of a Novel Class of Enantioselective Electrophilic Fluorinating Agents: N-Fluoro Ammonium Salts of cinchona Alkaloids (F-CA-BF₄). *Org. Lett.* **2000**, 2, 3699-3701; (c) Mohar, B.; Baudoux, J.; Plaquevent, J.-C.; Cahard, D., Electrophilic fluorination mediated by cinchona alkaloids: highly enantioselective synthesis of α -fluoro- α -phenylglycine derivatives. *Angew. Chem., Int. Ed.* **2001**, 40, 4214-4216; (d) Shibata, N.; Suzuki, E.; Asahi, T.; Shiro, M., Enantioselective Fluorination Mediated by Cinchona Alkaloid Derivatives/Selectfluor Combinations: Reaction Scope and Structural Information for N-Fluoro cinchona Alkaloids. *J. Am. Chem. Soc.* **2001**, 123, 7001-7009.

45. (a) Banks, R. E.; Lawrence, N. J.; Popplewell, A. L., Oxidation of benzylic alcohols and benzaldehydes with the Selectfluor reagent F-TEDA-BF₄ {1-chloromethyl-4-fluoro-1,4-diazoniabicyclo[2.2.2]octane bis(tetrafluoroborate)}. *Synlett* **1994**, 831-832; (b) Jing, Y.; Huang, X., Fluorous thiols in oligosaccharide

- synthesis. *Tetrahedron Lett.* **2004**, *45*, 4615-4618; (c) Kirihaara, M.; Naito, S.; Ishizuka, Y.; Hanai, H.; Noguchi, T., Oxidation of disulfides with Selectfluor: concise syntheses of thiosulfonates and sulfonyl fluorides. *Tetrahedron Lett.* **2011**, *52*, 3086-3089; (d) Pavlinac, J.; Zupan, M.; Stavber, S., Effect of Water on the Functionalization of Substituted Anisoles with Iodine in the Presence of F-TEDA-BF₄ or Hydrogen Peroxide. *J. Org. Chem.* **2006**, *71*, 1027-1032.
46. (a) Stavber, S.; Jereb, M.; Zupan, M., Selectfluor F-TEDA-BF₄ mediated and solvent directed iodination of aryl alkyl ketones using elemental iodine. *Chem. Commun.* **2002**, 488-489; (b) Stavber, S.; Kralj, P.; Zupan, M., Selective and effective iodination of alkyl-substituted benzenes with elemental iodine activated by selectfluor F-TEDA-BF₄. *Synlett* **2002**, 598-600; (c) Stavber, S.; Kralj, P.; Zupan, M., Progressive direct iodination of sterically hindered alkyl substituted benzenes. *Synthesis* **2002**, 1513-1518.
47. (a) Kralj, P.; Zupan, M.; Stavber, S., Remarkable Effect of Water on Functionalization of the Phenyl Ring in Methyl-Substituted Benzene Derivatives with F-TEDA-BF₄. *J. Org. Chem.* **2006**, *71*, 3880-3888; (b) Stavber, S.; Kralj, P.; Zupan, M., Effective and versatile functionalisation of hexamethylbenzene using N-F reagents. *Synlett* **2001**, 1152-1154; (c) Stavber, S.; Kralj, P.; Zupan, M., Selectfluor F-TEDA-BF₄ mediated introduction of perfluoroalkyl-containing groups in the benzylic position of hexamethylbenzene. *Acta Chim. Slov.* **2002**, *49*, 553-560.
48. (a) Brenzovich, J. W. E.; Benitez, D.; Lackner, A. D.; Shunatona, H. P.; Tkatchouk, E.; Goddard, I. I. I. W. A.; Toste, F. D., Gold-Catalyzed Intramolecular Aminoarylation of Alkenes: C-C Bond Formation through Bimolecular Reductive Elimination. *Angew. Chem., Int. Ed.* **2010**, *49*, 5519-5522; (b) Cui, L.; Zhang, G.; Zhang, L., Homogeneous gold-catalyzed efficient oxidative dimerization of propargylic acetates. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 3884-3887; (c) Melhado, A. D.; Brenzovich, W. E., Jr.; Lackner, A. D.; Toste, F. D., Gold-Catalyzed Three-Component Coupling: Oxidative Oxyarylation of Alkenes. *J. Am. Chem. Soc.* **2010**, *132*, 8885-8887; (d) Peng, Y.; Cui, L.; Zhang, G.; Zhang, L., Gold-Catalyzed Homogeneous Oxidative C-O Bond Formation: Efficient Synthesis of 1-Benzoxovinyl Ketones. *J. Am. Chem. Soc.* **2009**, *131*, 5062-5063; (e) Zhang, G.; Cui, L.; Wang, Y.; Zhang, L., Homogeneous Gold-Catalyzed Oxidative Carboheterofunctionalization of Alkenes. *J. Am. Chem. Soc.* **2010**, *132*, 1474-1475; (f) Zhang, G.; Peng, Y.; Cui, L.; Zhang, L., Gold-Catalyzed Homogeneous Oxidative Cross-Coupling Reactions. *Angew. Chem., Int. Ed.* **2009**, *48*, 3112-3115.
49. Qian, D.; Zhang, J., Au(I)/Au(III)-catalyzed Sonogashira-type reactions of functionalized terminal alkynes with arylboronic acids under mild conditions. *Beilstein J. Org. Chem.* **2011**, *7*, 808-812, No. 92.
50. Coleman, P. J.; Cox, C. D. Preparation of diarylpyrazolyfluoroalkylamines as mitotic kinesin inhibitors. WO2006110390A1, 2006.
51. Wu, J.; Liu, Y.; Deng, Y.; Chen, W.; Zhu, Y. Process for preparation of prostaglandin derivative and its intermediate. CN102101835A, 2011.

52. Ueno, R.; Ueno, R.; Kato, I.; Oda, T. Prostaglandins E and antiulcer containing compositions containing same. EP284180A1, 1988.
53. Seth, P. P.; Swayze, E. E. Preparation and RNA hybridization studies of cyclohexenyl nucleic acid analogs. WO2010036696A1, 2010.
54. (a) Bouvet, D.; O'Hagan, D., The synthesis of 1-fluoro- and 1,1-difluoro-analogues of 1-deoxy-d-xylulose. *Tetrahedron* **1999**, *55*, 10481-10486; (b) Bravo, P.; Piovosi, E.; Resnati, G., Homochiral fluoro-organic compounds. Part 10. A new protocol for the synthesis of variously functionalized α -fluoro-ketones and α -fluoro-alcohols. *J. Chem. Res., Synop.* **1989**, 134-135; (c) Chung, W. J.; Ngo, S. C.; Higashiya, S.; Welch, J. T., Lewis acid promoted aldol reaction of fluorinated silyl enol ethers from new fluoroacetylsilane derivatives. *Tetrahedron Lett.* **2004**, *45*, 5403-5406; (d) Leroy, J.; Wakselman, C., Synthesis of alkenyl fluoromethyl ketones via a phosphorus ylid. *Synthesis* **1982**, 496-497; (e) Linderman, R. J.; Graves, D. M., Oxidation of fluoroalkyl-substituted carbinols by the Dess-Martin reagent. *J. Org. Chem.* **1989**, *54*, 661-668; (f) Palacios, F.; Ochoa de Retana, A. M.; Pascual, S.; Oyarzabal, J., Preparation of Fluoroalkyl Imines, Amines, Enamines, Ketones, α -Amino Carbonyls, and α -Amino Acids from Primary Enamine Phosphonates†. *J. Org. Chem.* **2004**, *69*, 8767-8774; (g) Palacios, F.; Pascual, S.; Oyarzabal, J.; Ochoa de Retana, A. M., Fluoroalkyl α,β -Unsaturated Imines. Valuable Synthetic Intermediates from Primary Fluorinated Enamine Phosphonates. *Org. Lett.* **2002**, *4*, 769-772; (h) Yang, Y.; Zheng, F.; Qing, F.-L., Synthesis of 2',3'-dideoxy-6'-fluorocarbocyclic nucleosides via Reformatskii - Claisen rearrangement. *Tetrahedron* **2011**, *67*, 3388-3394.
55. (a) Bergmann, E. D.; Cohen, S., Organic fluorine compounds. IX. Preparation of fluoroacetone and 1,3-difluoroacetone. *J. Chem. Soc.* **1958**, 2259-2262; (b) Chaabouni, M. M.; Baklouti, A., Ring cleavage of (fluoroalkyl)- and (chloroalkyl)oxiranes by amine hydrofluorides. *Bull. Soc. Chim. Fr.* **1989**, 549-553; (c) Cherbuliez, E.; Yazgi, A.; Rabinowitz, J., Formation and transformation of organic derivatives of fluorine. II. Synthesis and esterification of fluorinated amino alcohols. *Helv. Chim. Acta* **1960**, *43*, 1135-1142; (d) Ichihara, J.; Hanafusa, T., The combination of hydrogen fluoride salt and aluminum fluoride: an efficient solid reagent for epoxide opening to give fluorohydrins. *J. Chem. Soc., Chem. Commun.* **1989**, 1848-1850.
56. (a) Arthurs, C. L.; Wind, N. S.; Whitehead, R. C.; Stratford, I. J., Analogues of 2-crotonyloxymethyl-(4R,5R,6R)-4,5,6-trihydroxycyclohex-2-enone (COTC) with anti-tumor properties. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 553-557; (b) Cui, H.-F.; Yang, Y.-Q.; Chai, Z.; Li, P.; Zheng, C.-W.; Zhu, S.-Z.; Zhao, G., Enantioselective Synthesis of Functionalized Fluorinated Cyclohexenones via Robinson Annulation Catalyzed by Primary-Secondary Diamines. *J. Org. Chem.* **2009**, *75*, 117-122; (c) Franck-Neumann, M.; Geoffroy, P.; Gumery, F., Aldol condensation reactions of tricarbonyliron complexes. Towards building blocks for the synthesis of carbomycin B/tylosin macrolide antibiotics and fluorinated analogs. *Tetrahedron Lett.* **2000**, *41*, 4219-4222; (d) Hollowood, C. J.; Ley, S. V.; Wright, E. A., 1,5-Asymmetric induction of chirality using [small pi]-

allyltricarbonyliron lactone complexes: highly diastereoselective synthesis of [small alpha]-functionalised carbonyl compounds. *Org. Biomol. Chem.* **2003**, *1*, 3208-3216; (e) Stavber, G.; Zupan, M.; Stavber, S., Micellar-system-mediated direct fluorination of ketones in water. *Synlett* **2009**, 589-594; (f) Stavber, S.; Jereb, M.; Zupan, M., Direct α -fluorination of ketones using N-F reagents. *Synthesis* **2002**, 2609-2615; (g) Stavber, S.; Zupan, M., High yield direct fluorofunctionalisation of ketones using AccufluorTM-NFTh fluorinating reagent. *Tetrahedron Lett.* **1996**, *37*, 3591-3594.

57. Hopkinson, M. N.; Giuffredi, G. T.; Gee, A. D.; Gouverneur, V., Gold-catalyzed diastereoselective synthesis of α -fluoroenones from propargyl acetates. *Synlett* **2010**, 2737-2742.

58. de, H. T.; Nevado, C., Domino gold-catalyzed rearrangement and fluorination of propargyl acetates. *Chem. Commun.* **2011**, *47*, 248-249.

59. (a) Buzas, A. K.; Istrate, F. M.; Gagosz, F., Gold(I)-Catalyzed Isomerization of Allenyl Carbinol Esters: An Efficient Access to Functionalized 1,3-Butadien-2-ol Esters. *Org. Lett.* **2007**, *9*, 985-988; (b) Zhang, X.; Fu, C.; Ma, S., Highly Stereoselective Facile Synthesis of 2-Acetoxy-1,3(E)-alkadienes via a Rh(I)-Catalyzed Isomerization of 2,3-Allenyl Carboxylates. *Org. Lett.* **2011**, *13*, 1920-1923.

60. We modified Gagosz's reaction condition by replacing CH_2Cl_2 with CDCl_3 considering two reasons: 1) Gold catalyst **4**, purchased from Aldrich, is more efficient in CDCl_3 than in CH_2Cl_2 . 2) Reaction can be easily monitored by ^1H -NMR.

61. We chose two steps in one pot reaction, because the gold catalyzed isomerization is not effective in CD_3CN and Selectfluor has low solubility in CDCl_3 .

62. ^1H and ^{19}F -NMR spectra of mixture of **3e** and ethyl acetate are available in supporting information.

63. Vincent, S. P.; Burkart, M. D.; Tsai, C.-Y.; Zhang, Z.; Wong, C.-H., Electrophilic Fluorination-Nucleophilic Addition Reaction Mediated by Selectfluor: Mechanistic Studies and New Applications. *J. Org. Chem.* **1999**, *64*, 5264-5279.

64. (a) Crabbe, P.; Fillion, H.; Andre, D.; Luche, J. L., Efficient homologation of acetylenes to allenes. *J. Chem. Soc., Chem. Commun.* **1979**, 859-862; (b) Kuang, J.; Ma, S., One-Pot Synthesis of 1,3-Disubstituted Allenes from 1-Alkynes, Aldehydes, and Morpholine. *J. Am. Chem. Soc.* **2010**, *132*, 1786-1787; (c) Mukaiyama, T.; Harada, T., A convenient synthesis of α -hydroxyallenes by the reaction of propargyl iodides with aldehydes in the presence of stannous halide. *Chem. Lett.* **1981**, 621-624.

65. (a) Burkart, M. D.; Vincent, S. P.; Düffels, A.; Murray, B. W.; Ley, S. V.; Wong, C.-H., Chemo-enzymatic synthesis of fluorinated sugar nucleotide: useful mechanistic Probes for glycosyltransferases. *Bioorg. Med. Chem.* **2000**, *8*, 1937-1946; (b) Dax, K.; Albert, M.; Ortner, J.; Paul, B. J., Synthesis of deoxyfluoro sugars from carbohydrate precursors. *Carbohydr. Res.* **2000**, *327*, 47-86.

66. Banks, R. E.; Lawrence, N. J.; Besheesh, M. K.; Popplewell, A. L.; Pritchard, R. G., Remote functionalization of (-)-menthol - synthesis of 4a,5,6,7,8,8a-hexahydro-4H-benzo[1,3]oxazine derivatives with the Selectfluor reagent F-TEDA-BF₄. *Chem. Commun.* **1996**, 1629-1630.
67. Michaudel, Q.; Thevenet, D.; Baran, P. S., Intermolecular Ritter-Type C-H Amination of Unactivated sp³ Carbons. *J. Am. Chem. Soc.* **2012**, *134*, 2547-2550.
68. (a) Krohn, K.; Franke, C.; Jones, P. G.; Aust, H. J.; Draeger, S.; Schulz, B., Biologically active metabolites of fungi. I. Isolation, synthesis, and biological activity of coniothyriomycin as well as bioassay of analogous open-chain imides. *Liebigs Ann. Chem.* **1992**, 789-798; (b) Maruyama, H. B.; Suhara, Y.; Suzuki-Watanabe, J.; Maeshima, Y.; Shimizu, N.; Ogura-Hamada, M.; Fujimoto, H.; Takano, K., New antibiotic, fumaramidmycin. I. Production, biological properties, and characterization of producer strain. *J. Antibiot.* **1975**, *28*, 636-647; (c) Ross, L.; Guarino, L. A., Cycloheximide inhibition of delayed early gene expression in baculovirus-infected cells. *Virology* **1997**, *232*, 105-113; (d) Suhara, Y.; Maruyama, H. B.; Kotoh, Y.; Miyasaka, Y.; Yokose, K.; Shirai, H.; Takano, K.; Quitt, P.; Lanz, P., New antibiotic, fumaramidmycin. II. Isolation, structure, and syntheses. *J. Antibiot.* **1975**, *28*, 648-655; (e) Thirkettle, J.; Alvarez, E.; Boyd, H.; Brown, M.; Diez, E.; Hueso, J.; Elson, S.; Fulston, M.; Gershater, C.; Morata, M. L.; Perez, P.; Ready, S.; Sanchez-Puelles, J. M.; Sheridan, R.; Stefanska, A.; Warr, S., SB-253514 and analogues; novel inhibitors of lipoprotein-associated phospholipase A₂. Produced by *Pseudomonas fluorescens* DSM 11579. I. Fermentation of producing strain, isolation and biological activity. *J. Antibiot.* **2000**, *53*, 664-669.
69. (a) Eliel, E. L.; McBride, R. T.; Kaufmann, S., Abnormal Chichibabin Reactions. The Condensation of Phenylacetaldehyde and Homoveratric Aldehyde with Ammonia¹. *J. Am. Chem. Soc.* **1953**, *75*, 4291-4296; (b) Ficken, G. E.; Linstead, R. P., 945. Conjugated macrocycles. Part XXIII. Tetracyclohexeno-tetrazaporphin. *J. Chem. Soc.* **1952**, 4846-4854.
70. (a) Alper, H.; Mahatantila, C.; Einstein, F.; Willis, A., A novel and stereospecific imide synthesis. *J. Am. Chem. Soc.* **1984**, *106*, 2708-2710; (b) Evans, D. A.; Nagorny, P.; Xu, R., Ceric Ammonium Nitrate Promoted Oxidation of Oxazoles. *Org. Lett.* **2006**, *8*, 5669-5671; (c) Gledhill, A. P.; McCall, C. J.; Threadgill, M. D., The formation and metabolism of N-hydroxymethyl compounds. Part 8. The oxidative decarboxylation of N-aroyleglycines to N-(acetoxymethyl)benzamides and N-formylbenzamides with lead(IV) acetate. *J. Org. Chem.* **1986**, *51*, 3196-3201; (d) Schnyder, A.; Indolese, A. F., Synthesis of Unsymmetrical Aroyl Acyl Imides by Aminocarbonylation of Aryl Bromides. *J. Org. Chem.* **2002**, *67*, 594-597; (e) Wang, F.; Liu, H.; Fu, H.; Jiang, Y.; Zhao, Y., Highly Efficient Iron(II) Chloride/N-Bromosuccinimide-Mediated Synthesis of Imides and Acylsulfonamides. *Adv. Synth. Catal.* **2009**, *351*, 246-252.
71. (a) Altomare, C.; Carotti, A.; Casini, G.; Cellamare, S.; Ferappi, M.; Gavuzzo, E.; Mazza, F.; Pantaleoni, G.; Giorgi, R., Synthesis and cognition-

activating properties of some mono- and bicyclic lactam derivatives. *J. Med. Chem.* **1988**, *31*, 2153-2158; (b) Gorka, A.; Hazai, L.; Szantay, C., Jr.; Hada, V.; Szabo, L.; Szantay, C., Synthesis of spiro-substituted benzo[c]azepinones. *Heterocycles* **2005**, *65*, 1359-1371; (c) Minisci, F.; Punta, C.; Recupero, F.; Fontana, F.; Pedulli, G. F., Aerobic Oxidation of N-Alkylamides Catalyzed by N-Hydroxyphthalimide under Mild Conditions. Polar and Enthalpic Effects. *J. Org. Chem.* **2002**, *67*, 2671-2676; (d) Nakayama, H.; Itoh, A., Aerobic photooxidation of benzylamide in the presence of catalytic iodine. *Synlett* **2008**, 675-678; (e) Nicolaou, K. C.; Mathison, C. J. N., Synthesis of Imides, N-Acyl Vinylogous Carbamates and Ureas, and Nitriles by Oxidation of Amides and Amines with Dess–Martin Periodinane. *Angew. Chem. Int. Ed.* **2005**, *44*, 5992-5997; (f) Papillon, J. P. N.; Taylor, R. J. K., The First Syntheses of the 1-Oxo-2-oxa-5-azaspiro[3.4]octane Ring System Found in Oxazolomycin. *Org. Lett.* **2000**, *2*, 1987-1990; (g) Sueda, T.; Kajishima, D.; Goto, S., Mechanistic Investigations on the Reaction between Amines or Amides and an Alkylperoxy- λ 3-iodane. *J. Org. Chem.* **2003**, *68*, 3307-3310; (h) Wang, J.-R.; Liu, L.; Wang, Y.-F.; Zhang, Y.; Deng, W.; Guo, Q.-X., Aerobic oxidation with N-hydroxyphthalimide catalysts in ionic liquid. *Tetrahedron Lett.* **2005**, *46*, 4647-4651; (i) Xu, L.; Zhang, S.; Trudell, M. L., Novel chromium(vi) catalyzed oxidation of N-alkylamides to imides with periodic acid. *Chem. Commun.* **2004**, 1668-1669; (j) Yoshifuji, S.; Arakawa, Y., Ruthenium tetroxide oxidation of 3,4-dihydroisoquinolin-1(2H)-ones: an efficient synthesis of isoquinoline-1,3,4(2H)-triones. *Chem. Pharm. Bull.* **1989**, *37*, 3380-3381.

72. McNeill, E.; Bois, J. D., Ruthenium-Catalyzed Hydroxylation of Unactivated Tertiary C–H Bonds. *J. Am. Chem. Soc.* **2010**, *132*, 10202-10204.

73. Han, J.-B.; Xu, B.; Hammond, G. B., Highly efficient Cu(I)-catalyzed synthesis of N-heterocycles through a cyclization-triggered addition of alkynes. *J. Am. Chem. Soc.* **2010**, *132*, 916-917.

74. Ye, C.; Shreeve, J. n. M., Structure-Dependent Oxidative Bromination of Unsaturated C–C Bonds Mediated by Selectfluor. *J. Org. Chem.* **2004**, *69*, 8561-8563.

75. (a) Chen, M. S.; White, M. C., A Predictably Selective Aliphatic C–H Oxidation Reaction for Complex Molecule Synthesis. *Science* **2007**, *318*, 783-787; (b) Chen, M. S.; White, M. C., Combined Effects on Selectivity in Fe-Catalyzed Methylene Oxidation. *Science* **2010**, *327*, 566-571.

76. Amide **1a** (1 equiv), copper (I) bromide (1.2 equiv, one portion) and Selectfluor (2.5 equiv) were stirred in acetonitrile under nitrogen atmosphere for 1 h; a 50% (NMR) yield of imide **2a** was recorded, which is comparable to entry 2 (Condition A) in Table 2.

77. The trace amount of water in commercial Selectfluor can not be removed by 50°C/1 mmHg vacuum overnight.

78. Jin, Z.; Xu, B.; Hammond, G. B., Copper mediated oxidation of amides to imides by Selectfluor. *Tetrahedron Lett.* **2011**, *52*, 1956-1959.

79. CuBr and Selectfluor were pre-mixed in acetonitrile for 1h, the resulting solution still can oxidize amide 1a into imide 2a (50% yield by ¹H-NMR).
80. Furuya, T.; Strom, A. E.; Ritter, T., Silver-Mediated Fluorination of Functionalized Aryl Stannanes. *J. Am. Chem. Soc.* **2009**, *131*, 1662-1663.
81. Christe, K. O.; Dixon, D. A.; McLemore, D.; Wilson, W. W.; Sheehy, J. A.; Boatz, J. A., On a quantitative scale for Lewis acidity and recent progress in polynitrogen chemistry. *J. Fluorine Chem.* **2000**, *101* (2), 151-153.
82. Krohn, K.; Elsasser, B.; Antus, S.; Konya, K.; Ammermann, E., Synthesis and structure-activity relationship of antifungal coniothyriomycin analogues. *J. Antibiot.* **2003**, *56*, 296-305.
83. (a) Cheng, B.; Wu, F.; Yang, X.; Zhou, Y.; Wan, X.; Zhai, H., Efficient Total Synthesis of Marine Alkaloid (-)-Nakadomarin A. *Chem.--Eur. J.* **2011**, *17*, 12569-12572; (b) Fuerstner, A.; Mathes, C.; Lehmann, C. W., Alkyne metathesis: development of a novel molybdenum-based catalyst system and its application to the total synthesis of epothilone A and C. *Chem.--Eur. J.* **2001**, *7*, 5299-5317; (c) Jakubec, P.; Kyle, A. F.; Calleja, J.; Dixon, D. J., Total synthesis of (-)-nakadomarin A: alkyne ring-closing metathesis. *Tetrahedron Lett.* **2011**, *52*, 6094-6097; (d) Kyle, A. F.; Jakubec, P.; Cockfield, D. M.; Cleator, E.; Skidmore, J.; Dixon, D. J., Total synthesis of (-)-nakadomarin A. *Chem. Commun.* **2011**, *47*, 10037-10039; (e) Morihira, K.; Hara, R.; Kawahara, S.; Nishimori, T.; Nakamura, N.; Kusama, H.; Kuwajima, I., Enantioselective Total Synthesis of Taxol. *J. Am. Chem. Soc.* **1998**, *120*, 12980-12981; (f) Mukaiyama, T.; Shiina, I.; Iwadare, H.; Saitoh, M.; Nishimura, T.; Ohkawa, N.; Sakoh, H.; Nishimura, K.; Tani, Y.-I.; Hasegawa, M.; Yamada, K.; Saitoh, K., Asymmetric total synthesis of taxol. *Chem.--Eur. J.* **1999**, *5*, 121-161; (g) Storer, R. I.; Takemoto, T.; Jackson, P. S.; Brown, D. S.; Baxendale, I. R.; Ley, S. V., Multi-step application of immobilized reagents and scavengers: A total synthesis of epothilone C. *Chem.--Eur. J.* **2004**, *10*, 2529-2547; (h) Trost, B. M.; Dong, G., Total synthesis of bryostatin 16 using atom-economical and chemoselective approaches. *Nature (London, U. K.)* **2008**, *456*, 485-488.
84. Martinez, M., Revision of Physalis Section Epeteiorhiza (Solanaceae). *Anales del Instituto de Biologia, Universidad Nacional Autonoma de Mexico: Serie Botanica*, 1998; p. 71.
85. Kuo, P.-C.; Kuo, T.-H.; Damu, A. G.; Su, C.-R.; Lee, E. J.; Wu, T.-S.; Shu, R.; Chen, C.-M.; Bastow, K. F.; Chen, T.-H.; Lee, K.-H., Physanolide A, a Novel Skeleton Steroid, and Other Cytotoxic Principles from Physalis angulata. *Org. Lett.* **2006**, *8*, 2953-2956.
86. (a) Abe, F.; Nagafuji, S.; Okawa, M.; Kinjo, J., Trypanocidal constituents in plants 6. Minor withanolides from the aerial parts of Physalis angulata. *Chem. Pharm. Bull.* **2006**, *54*, 1226-1228; (b) Chen, C. M.; Chen, Z. T.; Hsieh, C. H.; Li, W. S.; Wen, S. Y., Constituents of Formosan antitumor folk medicine. III. Withangulatin A, a new withanolide from Physalis angulata. *Heterocycles* **1990**, *31*, 1371-1375; (c) Damu, A. G.; Kuo, P.-C.; Su, C.-R.; Kuo, T.-H.; Chen, T.-H.;

Bastow, K. F.; Lee, K.-H.; Wu, T.-S., Isolation, Structures, and Structure-Cytotoxic Activity Relationships of Withanolides and Physalins from *Physalis angulata*. *J. Nat. Prod.* **2007**, *70*, 1146-1152; (d) He, Q.-P.; Ma, L.; Luo, J.-Y.; He, F.-Y.; Lou, L.-G.; Hu, L.-H., Cytotoxic withanolides from *Physalis angulata* L. *Chem. Biodiversity* **2007**, *4*, 443-449; (e) Ismail, N.; Alam, M., A novel cytotoxic flavonoid glycoside from *Physalis angulata*. *Fitoterapia* **2001**, *72*, 676-679; (f) Lee, S.-W.; Pan, M.-H.; Chen, C.-M.; Chen, Z.-T., Withangulatin I, a new cytotoxic withanolide from *Physalis angulata*. *Chem. Pharm. Bull.* **2008**, *56*, 234-236; (g) Makino, B.; Kawai, M.; Ogura, T.; Nakanishi, M.; Yamamura, H.; Butsugan, Y., Structural revision of physalin H isolated from *Physalis angulata*. *J. Nat. Prod.* **1995**, *58*, 1668-1674; (h) Row, L. R.; Reddy, K. S.; Sarma, N. S.; Matsuura, T.; Nakashima, R., New physalins from *Physalis angulata* and *Physalis lancifolia*. Structure and reactions of physalins D, I, G, and K. *Phytochemistry* **1980**, *19*, 1175-81; (i) Row, L. R.; Sarma, N. S.; Matsuura, T.; Nakashima, R., New physalins from *Physalis angulata* and *P. lancifolia*. Part 1. Physalins E and H, new physalins from *Physalis angulata* and *P. lancifolia*. *Phytochemistry* **1978**, *17*, 1641-1645; (j) Row, L. R.; Sarma, N. S.; Reddy, K. S.; Matsuura, T.; Nakashima, R., New physalins from *Physalis angulata* and *P. lancifolia*. Part 2. The structure of physalins F and J from *Physalis angulata* and *P. lancifolia*. *Phytochemistry* **1978**, *17*, 1647-1650; (k) Shingu, K.; Marubayashi, N.; Ueda, I.; Yahara, S.; Nohara, T., Constituents of solanaceous plants. XXI. Physagulin C, a new withanolide from *Physalis angulata* L. *Chem. Pharm. Bull.* **1991**, *39*, 1591-1593; (l) Shingu, K.; Yahara, S.; Okabe, H.; Nohara, T., Constituents of solanaceous plants. XXIV. Three new withanolides, physagulins E, F and G from *Physalis angulata* L. *Chem. Pharm. Bull.* **1992**, *40*, 2448-2451.

87. Row, L. R.; Sarma, N. S.; Reddy, K. S.; Matsuura, T.; Nakashima, R., The structure of Physalins F and J from *Physalis angulata* and *P. Lancifolia*. *Phytochemistry* **1978**, *17*, 1647-1650.

88. (a) Januario, A. H.; Rodrigues, F. E.; Pietro, R. C. L. R.; Kashima, S.; Sato, D. N.; Franca, S. C., Antimycobacterial physalins from *Physalis angulata* L. (Solanaceae). *Phytother. Res.* **2002**, *16*, 445-448; (b) Magalhaes, H. I. F.; Veras, M. L.; Torres, M. R.; Alves, A. P. N. N.; Pessoa, O. D. L.; Silveira, E. R.; Costa-Lotufo, L. V.; Odorico, d. M. M.; Pessoa, C., In-vitro and in-vivo antitumour activity of physalins B and D from *Physalis angulata*. *J. Pharm. Pharmacol.* **2006**, *58*, 235-241; (c) Pinto, N. B.; Morais, T. C.; Carvalho, K. M. B.; Silva, C. R.; Andrade, G. M.; Brito, G. A. C.; Veras, M. L.; Pessoa, O. D. L.; Rao, V. S.; Santos, F. A., Topical anti-inflammatory potential of Physalin E from *Physalis angulata* on experimental dermatitis in mice. *Phytomedicine* **2010**, *17*, 740-743; (d) Sa, M. S.; de, M. M. N.; Krettli, A. U.; Ribeiro, I. M.; Tomassini, T. C. B.; Ribeiro, d. S. R.; de, A. W. F.; Soares, M. B. P., Antimalarial Activity of Physalins B, D, F, and G. *J. Nat. Prod.* **2011**, *74*, 2269-2272.

89. Carbon numbering followed steroid numbering system since we assumed that left part of Physangulidine A was derived from steroid, and D ring of the steroid is disconnected by C-13 and C-17. Similar numbering was applied to Physalin, in which steroid is disconnected by C-13 and C-14.

90. (a) Chen, L.-X.; He, H.; Qiu, F., Natural withanolides: an overview. *Nat. Prod. Rep.* **2011**, *28*, 705-740; (b) Glotter, E., Withanolides and related ergostane-type steroids. *Nat. Prod. Rep.* **1991**, *8*, 415-440; (c) Misico, R. I.; Nicotra, V. E.; Oberti, J. C.; Barboza, G.; Gil, R. R.; Burton, G., Withanolides and related steroids. *Fortschr Chem Org Naturst* **2011**, *94*, 127-229.
91. <http://www.theplantlist.org/browse/A/Annonaceae/Crematosperma/>.
92. Cava, M. P.; Wakisaka, K.; Noguchi, I.; Edie, D. L.; DaRocha, A. I., Phlebicine, a new biphenylbisbenzylisoquinoline alkaloid from *Crematosperma polyphlebum*. *J. Org. Chem.* **1974**, *39*, 3588-3590.
93. (a) Atsumi, T.; Fujisawa, S.; Satoh, K.; Sakagami, H.; Iwakura, I.; Uehai, T.; Sugita, Y.; Yokoe, I., Cytotoxicity and radical intensity of eugenol, isoeugenol or related dimers. *Anticancer Res.* **2000**, *20*, 2519-2524; (b) Bortolomeazzi, R.; Verardo, G.; Liessi, A.; Callea, A., Formation of dehydrodiisoeugenol and dehydrodieugenol from the reaction of isoeugenol and eugenol with DPPH radical and their role in the radical scavenging activity. *Food Chem.* **2009**, *118*, 256-265; (c) Le, Q. P. W.; Larrahondo, J. E.; Raffauf, R. F., Antitumor plants. X. Constituents of *Nectandra rigida*. *J. Nat. Prod.* **1980**, *43*, 353-359.
94. Meyer, I.; Koch, O.; Krutmann, J. Use of dihydrodehydrodiisoeugenol as skin care preparations. 2010.
95. (a) Aiba, C. J.; Correa, R. G. C.; Gottlieb, O. R., Chemistry of Brazilian Lauraceae. XXII. Natural occurrence of Erdtman's dehydrodiisoeugenol. *Phytochemistry* **1973**, *12*, 1163-1164; (b) Enriquez, R. G.; Chavez, M. A.; Reynolds, W. F., Phytochemical investigations of plants of the genus *Aristolochia*, 1. Isolation and NMR spectral characterization of eupomatenoid derivatives. *J. Nat. Prod.* **1984**, *47*, 896-899.
96. <http://en.wikipedia.org/wiki/Hyptis>.
97. Gonzalez, A. G.; Bazzocchi, I. L.; Moujir, L.; Ravelo, A. G.; Correa, M. D.; Gupta, M. P., Xanthine oxidase inhibitory activity of some Panamanian plants from Celastraceae and Lamiaceae. *J Ethnopharmacol* **1995**, *46*, 25-29.
98. <http://en.wikipedia.org/wiki/Bocconia>.
99. Choi, E. W.; Nayak, L. V.; Bates, P. J., Cancer-selective antiproliferative activity is a general property of some G-rich oligodeoxynucleotides. *Nucleic Acids Res.* **2010**, *38*, 1623-1635.
100. (a) Boyd, M. R.; Paull, K. D., Some practical considerations and applications of the National Cancer Institute in vitro anticancer drug discovery screen. *Drug Dev. Res.* **1995**, *34*, 91-109; (b) Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bokesch, H.; Kenney, S.; Boyd, M. R., New colorimetric cytotoxicity assay for anticancer-drug screening. *J. Natl. Cancer Inst.* **1990**, *82*, 1107-1112.

APPENDIX A: LIST OF ABBREVIATIONS

DMP: Dess-Martin Periodinane
DMSO: Dimethylsulfonyl Oxide
dr: diastereomeric ratio
FTIR: Fourier transform infrared spectroscopy
ee: Enantiomeric excess
EI: Electrospray Ionization
EPR: Electron paramagnetic resonance
Eq: Equation
Equiv: Equivalence
h: Hour
HPLC: High performance liquid chromatography
HRMS: High resolution mass spectroscopy
Hz: Hertz
IR: infrared
LDA: Lithium diisopropylamide
M: Molar
***m*:** meta
mg: milligram
min: minute
mL: milliliter
mmol: millimole
NMR: Nuclear magnetic resonance spectroscopy
***o*:** ortho
***p*:** para

ppm: Parts per million

r.t.: room temperature

TBS: *tert*-Butyldimethylsilyl

***tert*:** tertiary

TABF: Tetrabutylammonium fluoride

TIPS: Triisopropylsilyl

TMS: Trimethylsilyl

THF: Tetrahydrofuran

TLC: Thin layer chromatography

APPENDIX B: COPYRIGHT PERMISSION

JOHN WILEY AND SONS LICENSE TERMS AND CONDITIONS

Mar 06, 2012

This is a License Agreement between Zhuang jin ("You") and John Wiley and Sons ("John Wiley and Sons") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by John Wiley and Sons, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

License Number	2851400919014
License date	Feb 17, 2012
Licensed content publisher	John Wiley and Sons
Licensed content publication	European Journal of Organic Chemistry
Licensed content title	Green Synthesis of Vicinal Dithioethers and Alkenyl Thioethers from the Reaction of Alkynes and Thiols in Water
Licensed content author	Zhuang Jin, Bo Xu, Gerald B. Hammond
Licensed content date	Jan 1, 2010
Start page	168
End page	173
Type of use	Dissertation/Thesis
Requestor type	Author of this Wiley article
Format	Print
Portion	Full article
Will you be translating?	No
Order reference number	
Total	0.00 USD

Terms and Conditions

TERMS AND CONDITIONS

This copyrighted material is owned by or exclusively licensed to John Wiley & Sons, Inc. or one of its group companies (each a "Wiley Company") or a society for whom a Wiley Company has exclusive publishing rights in relation to a particular journal (collectively WILEY"). By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the billing and payment terms and conditions established by the Copyright Clearance Center Inc., ("CCC's Billing and Payment terms and conditions"), at the time that you opened your Rightslink account (these are available at any time at <http://myaccount.copyright.com>)

**JOHN WILEY AND SONS LICENSE
TERMS AND CONDITIONS**

Mar 06, 2012

This is a License Agreement between Zhuang jin ("You") and John Wiley and Sons ("John Wiley and Sons") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by John Wiley and Sons, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

License Number	2851410409242
License date	Feb 17, 2012
Licensed content publisher	John Wiley and Sons
Licensed content publication	Chemistry - A European Journal
Licensed content title	Gold-Catalyzed, Intramolecular, Oxygen-Transfer Reactions of 2-Alkynyl-1,5-diketones or 2-Alkynyl-5-ketoesters: Scope, Expansion, and Mechanistic Investigations on a New [4 2] Cycloaddition
Licensed content author	Le-Ping Liu, Deepika Malhotra, Zhuang Jin, Robert S. Paton, K. N. Houk, Gerald B. Hammond
Licensed content date	Sep 12, 2011
Start page	10690
End page	10699
Type of use	Dissertation/Thesis
Requestor type	Author of this Wiley article
Format	Print and electronic
Portion	Full article
Will you be translating?	No
Order reference number	
Total	0.00 USD

Terms and Conditions

TERMS AND CONDITIONS

This copyrighted material is owned by or exclusively licensed to John Wiley & Sons, Inc. or one of its group companies (each a "Wiley Company") or a society for whom a Wiley Company has exclusive publishing rights in relation to a particular journal (collectively WILEY). By clicking "accept" in connection with completing this licensing transaction, you agree that the following terms and conditions apply to this transaction (along with the billing and payment terms and conditions established by the Copyright Clearance Center Inc., ("CCC's Billing and Payment terms and conditions"), at the time that you opened your Rightslink account (these are available at any time at <http://mvaccount.copyright.com>)

**ELSEVIER LICENSE
TERMS AND CONDITIONS**

Mar 06, 2012

This is a License Agreement between Zhuang jin ("You") and Elsevier ("Elsevier") provided by Copyright Clearance Center ("CCC"). The license consists of your order details, the terms and conditions provided by Elsevier, and the payment terms and conditions.

All payments must be made in full to CCC. For payment instructions, please see information listed at the bottom of this form.

Supplier	Elsevier Limited The Boulevard, Langford Lane Kidlington, Oxford, OX5 1GB, UK
Registered Company Number	1982084
Customer name	Zhuang jin
Customer address	788 raymond kent ct apt3 louisville, Ky 40217
License number	2851410063072
License date	Feb 17, 2012
Licensed content publisher	Elsevier
Licensed content publication	Tetrahedron Letters
Licensed content title	Copper mediated oxidation of amides to imides by Selectfluor
Licensed content author	Zhuang Jin, Bo Xu, Gerald B. Hammond
Licensed content date	20 April 2011
Licensed content volume number	52
Licensed content issue number	16
Number of pages	4
Start Page	1956
End Page	1959
Type of Use	reuse in a thesis/dissertation
Portion	full article
Format	both print and electronic
Are you the author of this Elsevier article?	Yes
Will you be translating?	No
Order reference number	
Title of your thesis/dissertation	Copper mediated oxidation of amides to imides by Selectfluor

RightsLink®

Home

Account
Info

Help

ACS Publications Title:
High quality. High impact.

Physangulidines A, B, and C:
Three New Antiproliferative
Withanolides from *Physalis*
angulata L.

Logged in as:
Zhuang jin
Account #:
3000500119

Author: Zhuang Jin et al.

Publication: Organic Letters

Publisher: American Chemical Society

Date: Mar 1, 2012

Copyright © 2012, American Chemical Society

LOGOUT

PERMISSION/LICENSE IS GRANTED FOR YOUR ORDER AT NO CHARGE

This type of permission/license, instead of the standard Terms & Conditions, is sent to you because no fee is being charged for your order. Please note the following:

- Permission is granted for your request in both print and electronic formats.
- If figures and/or tables were requested, they may be adapted or used in part.
- Please print this page for your records and send a copy of it to your publisher/graduate school.
- Appropriate credit for the requested material should be given as follows: "Reprinted (adapted) with permission from (COMPLETE REFERENCE CITATION). Copyright (YEAR) American Chemical Society." Insert appropriate information in place of the capitalized words.
- One-time permission is granted only for the use specified in your request. No additional uses are granted (such as derivative works or other editions). For any other uses, please submit a new request.

BACK

CLOSE WINDOW

Copyright © 2012 Copyright Clearance Center, Inc. All Rights Reserved. [Privacy statement](#).
Comments? We would like to hear from you. E-mail us at customercare@copyright.com

Dear Jin,

We herewith grant you permission to use the following article as part of your cumulative dissertation:

Aponte, José C.; Jin, Zhuang; Vaisberg, Abraham J.; Castillo, Denis; Málaga, Edith; Lewis, Walter H.; Sauvain, Michel; Gilman, Robert H.; Hammond, Gerald B.: Cytotoxic and Anti-infective Phenolic Compounds Isolated from *Mikania decora* and *Crematosperma microcarpum*" Planta Med 2011; 77: 1597-1599.

Permission is granted provided that:

- the dissertation is not published, neither in print nor online
- Thieme will be informed should publication occur

Please note that this is a permission to reproduce the article; copyright remains with the publisher.

With my very best regards and best wishes for your dissertation,
Regina Wenzel

Dr.Regina Wenzel
Programmplannung / Publishing Editor

Georg Thieme Verlag KG
Rüdigerstraße 14 | 70469 Stuttgart

Fon +49[0]711/8931-102
Fax +49[0]711/8931-392
Regina.Wenzel@thieme.de
www.thieme.de

Georg Thieme Verlag KG | Rüdigerstr. 14 | 70469 Stuttgart Rechtsform: KG | Sitz und Handelsregister: Stuttgart, HRA 3499

APPENDIX C: NMR SPECTRA FOR NEW COMPOUNDS

zj-2-53-1

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

zj-2-53-1

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

4J-4-33 4
Pulse Sequence: s2pu1

d

zj-2-57-2

Pulse Sequence: s2pu1

^1H NMR Spectrum of

2-1-3d

zj-c-s/-c
Pulse Sequence: s2pu1

2-1-3d

^{13}C NMR Spectrum of

zj-2-88-2
Pulse Sequence: s2pu1

zj-2-88-2

Pulse Sequence: s2pul

¹H NMR Spectrum of
2-1-3g, Diastereomers 1

Diastereomer1
Archive directory: /export/home/hammond/vmrays/data
Sample directory:
File: PROTON
Pulse Sequence: s2pul

9

zj-2-60-2-1
Pulse Sequence: szpul

Diastereomer2

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: p820N

Pulse Sequence: s2pul

2-1-3g, Diastereomers 2

¹H NMR Spectrum of

diastereomer 2

9

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

^{13}C NMR Spectrum of

2-1-3g, Diastereomers 2

sj-2-74-1-1

h

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: PROTON

Pulse Sequence: s2pul

^1H NMR Spectrum of

2-1-3h

xj-2-74-1-1

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

^{13}C NMR Spectrum of

2-1-3h

zj-2-77-1-2

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: PROTON

Pulse Sequence: s2pul

¹H NMR Spectrum of

sj-2-77-1-2

Archive directory: /export/home/hammond/vnmrsys/data

Sample directory:

File: CARBON

Pulse Sequence: s2pul

^{13}C NMR Spectrum of

[illegible]

zj-2-89-1-1
Pulse Sequence: s2pu1

zj-2-93-2

Pulse Sequence: s2pu1

u

^1H NMR Spectrum of

2-1-4a (E/Z=9:1)

E=Z=9:1

154

zj-2-93-2
Pulse Sequence: szpul

129.112
129.165
130.035
130.973
132.055
132.957
133.217

134.018
133.476
133.346

77.252
77.000
76.746

65.990

135.910

59.192

15.313
14.632

130 120 110 100 90 80 70 60 50 40 30 20 ppm

zj-2-94-1

Pulse Sequence: s2pul

0

E:Z = 9:1

zj-2-94-1
Pulse Sequence: s2pul

Ab

zj-2-94-2

Pulse Sequence: s2pu1

6c

E:Z = 12.5:1

¹H NMR Spectrum of

2-1-4c (E/Z=12.5:1)

158

zj-2-94-2

Pulse Sequence: s2pu

¹³C NMR Spectrum of

2-1-4c (E/Z=12.5:1)

159

zj-2-95-1

Pulse Sequence: s2pu1

d

zj-2-95-1

Pulse Sequence: s2pu1

4d

7e

zj-2-96-2
 exp1 s2pu1
 SAMPLE
 date Jun 19 2009 temp 25.0
 solvent CDC13 gain not used
 file exp spin 0
 ACQUISITION
 sw 5995.7 hst 0.008
 at 2.733 pw90 7.900
 np 32768 alfa 0
 fb 3000 il
 bs 32 in
 dl 1.000 dp
 nt 8 hs
 ct 8
 TRANSMITTER fn not used
 tn H1
 sfrq 499.643 sp
 tof 0.0 wp 5096.3
 tpwr 59 rfl 499.6
 pw 3.950 rfp 0
 DECOUPLER C13 rp -6.6
 dn 0 lp -21.4
 dof 0
 dm nnn wc 250
 dmm c sc 0
 dpwr 45 vs 134
 dmf 17094 th 12
 a1 cdc ph

¹H NMR Spectrum of

2-1-4e (E/Z=12.5:1)

E:Z = 12.5:1

4e
zj-2-96-2

Pulse Sequence: s2pu1

^{13}C NMR Spectrum of CCCC=C(Cc1ccccc1S)O
2-1-4e (E/Z=12.5:1)

2J-2-101-1
Pulse Sequence: szput

164

zj-2-101-1

Pulse Sequence: s2pu1

¹³C NMR Spectrum of

2-1-4f (E/Z=17:1)

zj-2-101-2

Pulse Sequence: s2pul

4-9

¹H NMR Spectrum of

2-1-4g (E/Z=17:1)

zj-2-101-2
Pulse Sequence: s2pul

2-1-4g (E/Z=17:1)

zj-1-102-2

expl s2pu1

4h

SAMPLE		SPECIAL	
date	Jul 7 2009	temp	25.0
solvent	CDC13	gain	not used
file	exp	spin	0
ACQUISITION		hst	0.008
sw	5995.7	pw90	7.900
at	2.733	alfa	0
np	32768	FLAGS	
fb	3000	il	nn
bs	32	in	nn
dl	1.000	dp	nn
nt	8	hs	nn
ct	8	PROCESSING	
TRANSMITTER		fn	not used
tn	H1	DISPLAY	
strq	499.643	sp	-99.3
tof	0.0	wp	5096.3
tpwr	59	rfl	499.6
pw	3.950	rpf	-10.7
DECOUPLER		rp	-12.1
dn	C13	lp	
dof	0	PLOT	
dm	nnn	wc	250
dmm	c	sc	0
dpwr	45	vs	149
dmf	17094	th	9
	af	cdc	ph

¹H NMR Spectrum of

2-1-4h (E/Z=17:1)

zj-1-102-2
Pulse Sequence: s2pu1

2-1-4h (E/Z=17:1)

^{13}C NMR Spectrum of

fi

```

zj-2-103-2
exp1 s2pul

SAMPLE
date Jul 9 2009 temp 25.0
solvent CDCl3 gain not used
file exp spin 0
ACQUISITION hst 0.008
sw 5995.7 pw90 7.900
at 2.733 alfa 0
np 32768
fb 3000 il n
bs 32 in n
dl 1.000 dp y
nt 8 hs nn
ct 8

TRANSMITTER fn not used
tn H1
sfrq 499.643 sp
tof 0.0 wp 3442.7
tpwr 59 rfl 499.6
pw 3.950 rfp 0
DECOUPLER C13 lp -5.8
dn 0
dof 0 PLOT -22.5
dm nnn wc 250
dmm c sc 0
dpwr 45 vs 116
dref 17094 th 6
af cdc ph
  
```

E:Z = 20:1

zj-2-103-2

Pulse Sequence: zgpg30

AV

4)

zj-2-102-1-1

Pulse Sequence: s2pu1

A)

¹³C NMR Spectrum of

2-1-4j (E/Z=100:0)

zj-2-104-1

Pulse Sequence: s2pu1

4k

^1H NMR Spectrum of

2-1-4k (*E/Z*=100:0)

zj-2-104-1
Pulse Sequence: s2pul

AK

¹³C NMR Spectrum of

2-1-4k (E/Z=100:0)

zj-2-105-2
Pulse Sequence: s2pu1

2-1-4I (E/Z=100:0)

¹H NMR Spectrum of

zj-2-105-2
Pulse Sequence: s2pul1

AV

2J-2-185-1

Pulse Sequence: s2pu1

507

¹³C NMR Spectrum of

2-1-4m (E/Z=50:1)

zj-2-107-1

Pulse Sequence: s2pul

5

¹H NMR Spectrum of

2-1-5 (E/Z=10:1)

zj-2-107-1
Pulse Sequence: s2pu1

2-1-5 (E/Z=10:1)

¹³C NMR Spectrum of

zj-2-109-1

Pulse Sequence: s2pu1

Solvent: CDCl₃
Temp. 25.0 C / 298.1 K
INOVA-500 "ulnar3"

PULSE SEQUENCE

Relax. delay 1.000 sec

Pulse 45.0 degrees

Acq. time 2.045 sec

Width 3997.2 Hz

8 repetitions

OBSERVE H1, 499.6405729 MHz

DATA PROCESSING

FT size 16384

Total time 0 min, 24 sec

¹H NMR Spectrum of

2-1-6 (E/Z=12.5:1)

zj-2-109-1
Pulse Sequence: s2pu1

¹³C NMR Spectrum of

2-1-6 (E/Z=12.5:1)

Sample: ZJ-3-61-1
Sample ID: ZhuangJ_20100917_11
File: ZhuangJ_20100917_11/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603207 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: ZJ-3-61-1
Sample ID: ZhuangJ_20100917_12
File: ZhuangJ_20100917_12/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
256 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 19 min, 10 sec

¹³C NMR Spectrum of

Sample: Zj-3-63-1
Sample ID: ZhuangJ_20100922_08
File: ZhuangJ_20100922_08/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603522 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: zj-3-63-1
Sample ID: ZhuangJ_20100922_10
File: ZhuangJ_20100922_10/data/Carbon_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
1000 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 38 min, 21 sec

¹³C NMR Spectrum of

Sample: zj-3-64-1
Sample ID: ZhuangJ_20100924_01
File: ZhuangJ_20100924_01/data/Proton_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603503 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: zj-3-64-1
Sample ID: ZhuangJ_20100924_02
File: ZhuangJ_20100924_02/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRB-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
64 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 38 min, 21 sec

¹³C NMR Spectrum of

Sample: ZJ-3-65-2
Sample ID: ZhuangJ_20101001_06
File: ZhuangJ_20101001_06/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603207 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

^1H NMR Spectrum of

2-2-2d

Sample: ZJ-3-65-2
Sample ID: ZhuangJ_20101001_09
File: /home/walkup/vnmrsys/data/auto_2010.09.27/ZhuangJ_20101001_09/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Sample #37, Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
8000 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 5 hr, 6 min, 45 sec

¹³C NMR Spectrum of

Sample: ZJ-3-68-2
Sample ID: ZhuangJ_20101006_07
File: ZhuangJ_20101006_07/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdc13
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603207 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: ZJ-3-68-2
Sample ID: ZhuangJ_20101006_08
File: ZhuangJ_20101006_08/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
64 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 5 hr, 6 min, 45 sec

¹³C NMR Spectrum of

2-2-2e

Sample: ZJ-3-70-1
Sample ID: ZhuangJ_20101018_04
File: ZhuangJ_20101018_04/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
INOVA-500 "ulnmr500"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 7994.4 Hz
8 repetitions
OBSERVE H1, 499.6404224 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 31 sec

^1H NMR Spectrum of

194

Sample: ZJ-3-70-2
Sample ID: ZhuangJ_20101018_05
File: ZhuangJ_20101018_05/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
INOVA-500 "ulnmr500"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 30154.5 Hz
64 repetitions
OBSERVE C13, 125.6346841 MHz
DECOUPLE H1, 499.6429206 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 2.0 Hz
FT size 131072
Total time 8 hr, 20 min, 22 sec

561

Sample: ZJ-3-74-1
Sample ID: ZhuangJ_20101023_06
File: ZhuangJ_20101023_06/data/Proton_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7535764 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: zj-3-74-1
Sample ID: ZhuangJ_20101026_03
File: ZhuangJ_20101026_03/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Sample #25, Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
12000 repetitions
OBSERVE C13, 100.5181167 MHz
DECOUPLE H1, 399.7555752 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 7 hr, 40 min, 8 sec

¹³C NMR Spectrum of

Sample: ZJ-3-73-1-2
Sample ID: ZhuangJ_20101022_03
File: ZhuangJ_20101022_03/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
INOVA-500 "ulnmr500"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 7994.4 Hz
8 repetitions
OBSERVE H1, 499.6404224 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
PT size 65536
Total time 0 min, 31 sec

¹H NMR Spectrum of

2-2-2h

Sample: ZJ-3-73-1-2
Sample ID: ZhuangJ_20101022_04
File: ZhuangJ_20101022_04/data/Carbon_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
INOVA-500 "ulnmr500"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 30154.5 Hz
320 repetitions
OBSERVE C13, 125.6346841 MHz
DECOUPLE H1, 499.6429206 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 131072
Total time 38 min, 29 sec

^{13}C NMR Spectrum of

2-2-2h

Sample: ZJ-3-74-2
Sample ID: ZhuangJ_20101023_04
File: ZhuangJ_20101023_04/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMR5-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7535764 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

2-2-2i

Sample: ZJ-3-74-2
Sample ID: ZhuangJ_20101023_05
File: ZhuangJ_20101023_05/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
64 repetitions
OBSERVE C13, 100.5181167 MHz
DECOUPLE H1, 399.7555752 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 9 min, 49 sec

¹³C NMR Spectrum of
2-2-2i

zj-3-113-2

Sample Name:

zj-3-113-2

Data Collected on:

ulnmr400-vnmrs400

Archive directory:

/home/walkup/vnmrsys/data/ZhuangJ

Sample directory:

zj-3-113-2_20111022_01

FidFile: PROTON_001

Pulse Sequence: PROTON (s2pul)

Solvent: cdcl3

Data collected on: Oct 22 2011

Temp. 25.0 C / 298.1 K

Sample #44, Operator: ZhuangJ

Relax. delay 1.000 sec

Pulse 45.0 degrees

Acq. time 2.556 sec

Width 6410.3 Hz

8 repetitions

OBSERVE H1, 399.7275307 MHz

DATA PROCESSING

FT size 32768

Total time 0 min 28 sec

^1H NMR Spectrum of

zj-3-115-2

Sample Name:
zj-3-115-2
Data Collected on:
ulnmr400-vnmrs400
Archive directory:
/home/walkup/vnmrsys/data/ZhuangJ
Sample directory:
zj-3-115-2_20111108_02
FidFile: CARBON_001

Pulse Sequence: CARBON (s2pul)
Solvent: cdcl3
Data collected on: Nov 8 2011

Temp. 25.0 C / 298.1 K
Sample #42, Operator: ZhuangJ

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.285 sec
Width 25510.2 Hz
20000 repetitions
OBSERVE C13, 100.5094869 MHz
DECOUPLE H1, 399.7212548 MHz
Power 42 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 12 hr, 41 min

^{13}C NMR Spectrum of

zj-3-113-2

Sample Name:
zj-3-113-2
Data Collected on:
ulnmr400-vnmrs400
Archive directory:
/home/walkup/vnmrsys/data/ZhuangJ
Sample directory:
zj-3-113-2_20111022_01
FidFile: FLUORINE_001

Pulse Sequence: FLUORINE (s2pul)
Solvent: cdcl3
Data collected on: Oct 22 2011

Temp. 25.0 C / 298.1 K
Sample #44, Operator: ZhuangJ

Relax. delay 1.000 sec
Pulse 35.0 degrees
Acq. time 0.603 sec
Width 108.7 kHz
16 repetitions
OBSERVE F19, 376.1196667 MHz
DATA PROCESSING
FT size 131072
Total time 0 min 26 sec

¹⁹F NMR Spectrum of

Zj-3-110-2-1

Sample Name:
Zj-3-110-2-1
Data Collected on:
ulnmr400-vnmrs400
Archive directory:
/home/walkup/vnmrsys/data/ZhuangJ
Sample directory:
Zj-3-110-2-1_20110707_01
FidFile: PROTON_001

Pulse Sequence: PROTON (s2pul)
Solvent: cdcl3
Data collected on: Jul 7 2011

Temp. 25.0 C / 298.1 K
Sample #37, Operator: ZhuangJ

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.556 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7353714 MHz
DATA PROCESSING
FT size 32768
Total time 0 min 28 sec

¹H NMR Spectrum of

derived from

ZJ-3-110-2-1

Sample Name:

ZJ-3-110-2-1

Data Collected on:

ulnmr400-vnmrs400

Archive directory:

/home/walkup/vnmrsys/data/ZhuangJ

Sample directory:

ZJ-3-110-2-1_20110708_01

FidFile: CARBON_001

Pulse Sequence: CARBON (s2pul)

Solvent: cdcl3

Data collected on: Jul 8 2011

Temp. 25.0 C / 298.1 K

Sample #42, Operator: ZhuangJ

Relax. delay 1.000 sec

Pulse 45.0 degrees

Acq. time 1.285 sec

Width 25510.2 Hz

20000 repetitions

OBSERVE C13, 100.5135391 MHz

DECOUPLE H1, 399.7373701 MHz

Power 41 dB

continuously on

WALTZ-16 modulated

DATA PROCESSING

Line broadening 0.5 Hz

FT size 65536

Total time 12 hr, 41 min

^{13}C NMR Spectrum of

derived from

Zj-3-110-2-1

Sample Name:

Zj-3-110-2-1

Data Collected on:

ulnmr400-vnmrs400

Archive directory:

/home/walkup/vnmrsys/data/ZhuangJ

Sample directory:

Zj-3-110-2-1_20110707_01

FidFile: FLUORINE_001

Pulse Sequence: FLUORINE (s2pul)

Solvent: cdcl3

Data collected on: Jul 7 2011

Temp. 25.0 C / 298.1 K

Sample #37, Operator: ZhuangJ

Relax. delay 1.000 sec

Pulse 35.0 degrees

Acq. time 0.734 sec

Width 89285.7 Hz

16 repetitions

OBSERVE F19, 376.1270443 MHz

DATA PROCESSING

FT size 131072

Total time 0 min 28 sec

^{19}F NMR Spectrum of

derived from

Sample: zj-3-113-1
Sample ID: ZhuangJ_20111017_01
File: ZhuangJ_20111017_01/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
INOVA-500 "ulnmr500"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 7994.4 Hz
8 repetitions
OBSERVE H1, 499.6404224 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 31 sec

¹H NMR Spectrum of

derived from

3-2-1i

209

Pulse Sequence: CARBON (s2pul)
Solvent: cdcl3
Data collected on: Oct 17 2011

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.285 sec
Width 25510.2 Hz
20000 repetitions
OBSERVE C13, 100.5115675 MHz
DECOUPLE H1, 399.7295294 MHz
Power 42 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 12 hr, 41 min

Sample: zj-3-113-1
Sample ID: ZhuangJ_20111017_01
File: ZhuangJ_20111017_01/data/Fluorine_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Fluorine_01
INNOVA-500 "ulnmr500"

¹⁹F NMR Spectrum of

Relax. delay 1.000 sec
Pulse 30.0 degrees
Acq. time 1.000 sec
Width 103.5 kHz
16 repetitions
OBSERVE F19, 470.1317140 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 262144
Total time 0 min, 36 sec

derived from

3-2-1i

ROESY Spectrum of

Sample: zj-3-41-2
 Sample ID: ZhuangJ_20100610_01
 File: /home/walkup/vnmrsys/data/auto_2010.06.08/ZhuangJ_20100610_01/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
 Temp. 25.0 C / 298.1 K
 Operator: ZhuangJ
 File: Proton_01
 VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
 Pulse 45.0 degrees
 Acq. time 2.049 sec
 Width 6410.3 Hz
 8 repetitions
 OBSERVE H1, 399.7698366 MHz
 DATA PROCESSING
 Resol. enhancement -0.0 Hz
 FT size 65536
 Total time 0 min, 30 sec

¹H NMR Spectrum of

3-3-2-2c

Sample: xj-3-41-2
Sample ID: ZhuangJ_20100610_02
File: /home/walkup/vnmrsys/data/auto_2010.06.08/ZhuangJ_20100610_02/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
2000 repetitions
OBSERVE C13, 100.5222010 MHz
DECOUPLE H1, 399.7718181 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 1 hr, 16 min, 41 sec

sample: zj-3-50-1
sample ID: ZhuangJ_20100730_03
file: /home/walkup/vnmrsys/data/auto_2010.07.19/zhuangJ_20100730_03/data/proton_01.fid

Pulse Sequence: s2pul

solvent: cdcl3
temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: proton_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7698350 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Sample: zj-3-53-2
Sample ID: ZhuangJ_20100807_05
File: /home/walkup/vnmrSYS/data/auto_2010_08.02/ZhuangJ_20100807_05/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7698193 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

Sample: xj-3-50-1
Sample ID: ZhuangJ_20100730_04
File: /home/walkup/vnmrSYS/data/auto_2010.07.19/ZhuangJ_20100730_04/data/Carbon_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
1000 repetitions
OBSERVE C13, 100.5222010 MHz
DECOUPLE H1, 399.7718181 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 38 min, 21 sec

Sample: zj-3-53-2
Sample ID: ZhuangJ_20100807_05
File: /home/walkup/vnmrSYS/data/auto_2010.08.02/ZhuangJ_20100807_05/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7698193 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

Sample: zj-3-53-2
Sample ID: ZhuangJ_20100807_07
File: /home/walkup/vnmrsys/data/auto_2010.08.02/ZhuangJ_20100807_07/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
5000 repetitions
OBSERVE C13, 100.5222010 MHz
DECOUPLE H1, 399.7718181 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 3 hr, 11 min, 43 sec

¹³C NMR Spectrum of

3-3-2-2g

Sample: xj-3-53-2
Sample ID: ZhuangJ_20100807_06
File: /home/walkup/vnmrsys/data/auto_2010.08.02/ZhuangJ_20100807_06/data/Fluorine_01.fid

Pulse Sequence: s2pul

Solvent: cdc13
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Fluorine_01
VNMRS-400 "ulnar400"

Relax. delay 1.000 sec
Pulse 30.0 degrees
Acq. time 1.000 sec
Width 78125.0 Hz
16 repetitions
OBSERVE F19, 376.1594577 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 262144
Total time 0 min, 36 sec

Sample: xj-3-54-2
Sample ID: ZhuangJ_20100810_02
File: /home/walkup/vnmrsys/data/auto_2010.08.09/ZhuangJ_20100810_02/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Proton_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 6410.3 Hz
8 repetitions
OBSERVE H1, 399.7603207 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

3-3-2-2h

Sample: xj-3-54-2
Sample ID: ZhuangJ_20100810_04
File: /home/walkup/vnmrsys/data/auto_2010.08.09/ZhuangJ_20100810_04/data/Carbon_01.fid

Pulse Sequence: s2pul
Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMR-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
192 repetitions
OBSERVE C13, 100.5198126 MHz
DECOUPLE H1, 399.7623195 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 9 min, 49 sec

¹³C NMR Spectrum of

3-3-2-2h

Sample: zj-3-54-2
Sample ID: ZhuangJ_20100810_03
File: /home/walkup/vnmrsys/data/auto_2010.08.09/ZhuangJ_20100810_03/data/Fluorine_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Fluorine_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 30.0 degrees
Acq. time 1.000 sec
Width 78125.0 Hz
16 repetitions
OBSERVE F19, 376.1505201 MHz
DATA PROCESSING
Line broadening 0.9 Hz
FT size 262144
Total time 0 min, 36 sec

¹⁹F NMR Spectrum of

3-3-2-2h

-105.120
-105.128
-105.134
-105.141
-105.155

-40 -60 -80 -100 -120 -140 -160 -180 -200 -220 ppm

Sample: xj-3-52-2
 Sample ID: ZhuangJ_20100805_01
 File: /home/walkup/vnmrSYS/data/auto_2010.08.02/ZhuangJ_20100805_01/data/Proton_01.fid

Pulse Sequence: s2pul
 Solvent: cdcl3
 Temp. 25.0 C / 298.1 K
 Operator: ZhuangJ
 File: Proton_01
 VNMRs-400 "ulnmr400"

Relax. delay 1.000 sec
 Pulse 45.0 degrees
 Acq. time 2.049 sec
 Width 6410.3 Hz
 8 repetitions
 OBSERVE H1, 399.7698193 MHz
 DATA PROCESSING
 Resol. enhancement -0.0 Hz
 FT size 65536
 Total time 0 min, 30 sec

¹H NMR Spectrum of

3-3-2-2i

Sample: xj-3-52-2
Sample ID: ZhuangJ_20100805_02
File: /home/walkup/vnmrsys/data/auto_2010.08.02/ZhuangJ_20100805_02/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Operator: ZhuangJ
File: Carbon_01
VNMRS-400 "ulnmr400"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 24509.8 Hz
64 repetitions
OBSERVE C13, 100.5222010 MHz
DECOUPLE H1, 399.7718181 MHz
Power 40 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 65536
Total time 9 min, 49 sec

SPA14-3-4
Zhuang Jin
15 mg in cdcl3

Sample: SPA14-3-4
Sample ID: hammond_SPA14-3-4_20110613_02
File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.06.13_02/hammond_SPA14-3-4_20110613_02/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #2, Operator: hammond
File: Proton_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 5605.4 Hz
8 repetitions
OBSERVE H1, 699.8134495 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Physangulidine A (1)

SPA14-3-4
Zhuang Jin
15 mg in cdcl3

Sample: SPA14-3-4
Sample ID: hammond_SPA14-3-4_20110613_02
File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.06.13_02/hammond_SPA14-3-4_20110613_02/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #2, Operator: hammond
File: Carbon_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 43103.4 Hz
2000 repetitions
OBSERVE C13, 175.9682318 MHz
DECOUPLE H1, 699.8169486 MHz
Power 36 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 131072
Total time 1 hr, 16 min, 41 sec

¹³C NMR Spectrum of

Physangulidine A (1)

SPA14-3-4
Zhuang Jin
15 mg in cdcl3

Sample: SPA14-3-4
Sample ID: hammond SPA14-3-4_20110613_02
File: mnt/ulnmr700/wakeup/vnmrsys/data/auto_2011.06.13_02/hammond SPA14-3-4_20110613_02/data/Ghsqcad_01.fid

Pulse Sequence: ghsqcad

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #2, Operator: hammond
File: Ghsqcad_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Mixing 0.500 sec
Acq. time 0.230 sec
Width 5605.4 Hz
2D Width 29917.7 Hz
2 repetitions
2 x 256 increments
OBSERVE H1, 699.8134488 MHz
DECOUPLE C13, 175.9814294 MHz
Power 43 dB
on during acquisition
off during delay
W40_HCN_CP modulated
DATA PROCESSING
Gauss apodization 0.106 sec
F1 DATA PROCESSING
Gauss apodization 0.008 sec
FT size 4096 x 2048
Total time 22 min, 6 sec

HSQC Spectrum of

Physangulidine A (1)

2
3
4
5
6
7

```
Sample: SPA14-3-2
Sample ID: hammond SPA14-3-2_20110613_01
File: /mnt/ulnvr700/walkup/vnmrsws/data/auto_2011.06.13_02/hammond SPA14-3-2_20110613_01/data/Proton_01.fid
```

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #3, Operator: hammond
File: Proton_01
VNMR5-400 "ulnmr-ds"

```
Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 11261.3 Hz
8 repetitions
OBSERVE H1, 699.8134495 MHz
DATA PROCESSING
  Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec
```

¹H NMR Spectrum of

```

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 43103.4 Hz
10000 repetitions
OBSERVE C13, 175.9682318 MHz
DECOUPLE H1, 699.8169486 MHz
Power 36 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 0.5 Hz
FT size 131072
Total time 6 hr, 23 min, 26 sec

```

Physangulidine B (2)

234

F2

061302

/data/ai

3

4

7

7

6

5

4

3

2

1

F1 (ppm)

SPA14-3-2 (Zhuang Jin)

< 3mg in cdcl3

Sample: SPA14-3-2

Sample ID: hammond_SPA14-3-2_20110808_01

File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.08.08/hammond_SPA14-3-2_20110808_01/data/Ghmhc_01.fid

Pulse Sequence: gHMBC

SPA14-3-2 (Zhuang Jin)
< 3mg in cdcl3

Sample: SPA14-3-2
Sample ID: hammond_SPA14-3-2_20110809_01

File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.08.08/hammond_SPA14-3-2_20110809_01/data/Roesy_01.fid

Pulse Sequence: ROESY

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Sample #5, Operator: hammond
File: Roesy_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Mixing 0.400 sec
Acq. time 0.177 sec
Width 5787.0 Hz
2D Width 5787.0 Hz
32 repetitions
2 x 256 increments
OBSERVE H1, 699.8134510 MHz
DATA PROCESSING
Gauss apodization 0.082 sec
F1 DATA PROCESSING
Gauss apodization 0.082 sec
FT size 4096 x 4096
Total time 7 hr, 18 min, 36 sec

Physangulidine B (2)

ROESY Spectrum of

SPA14-3-3
Zhuang Jin
4.1 mg in cdcl3

Sample: SPA14-3-3
Sample ID: hammond SPA14-3-3_20110613_01
File: mnt/ulnmr700/walkup/vnmrsvs/data/auto_2011.06.13_02/hammond SPA14-3-3_20110613_01/data/Proton_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #1, Operator: hammond
File: Proton_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 2.049 sec
Width 11261.3 Hz
8 repetitions
OBSERVE H1, 699.8134935 MHz
DATA PROCESSING
Resol. enhancement -0.0 Hz
FT size 65536
Total time 0 min, 30 sec

¹H NMR Spectrum of

Physangulidine C (3)

SPA14-3-3
Zhuang Jin
4.1 mg in cdcl3

Sample: SPA14-3-3
Sample ID: hammond_SPA14-3-3_20110613_02
File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.06.13_02/hammond_SPA14-3-3_20110613_02/data/Carbon_01.fid

Pulse Sequence: s2pul

Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #1, Operator: hammond
Files Carbon_01
VNMR-400 "ulnmr-da" 574

Relax. delay 1.000 sec
Pulse 45.0 degrees
Acq. time 1.300 sec
Width 43103.4 Hz
10000 repetitions
OBSERVE C13, 175.9682318 MHz
DECOUPLE H1, 699.8169486 MHz
Power 36 dB
continuously on
WALTZ-16 modulated
DATA PROCESSING
Line broadening 1.0 Hz
FT size 131072
Total time 6 hr, 23 min, 26 sec

¹³C NMR Spectrum of

SPA14-3-3
Zhuang Jin
4.1 mg in cdcl3

Sample: SPA14-3-3
Sample ID: hammond_SPA14-3-3_20110613_02
File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.06.13_02/hammond_SPA14-3-3_20110613_02/data/Ghsqcad_01.fid

Pulse Sequence: gHSQCAD
Solvent: cdcl3
Temp. 22.0 C / 295.1 K
Sample #1, Operator: hammond
File: Ghsqcad_01
VNMR5-400 "ulnmr-ds"

Relax. delay 1.000 sec
Mixing 0.500 sec
Acq. time 0.230 sec
Width 5605.4 Hz
2D Width 29917.7 Hz
2 repetitions
2 x 256 increments
OBSERVE H1, 699.8134922 MHz
DECOUPLE C13, 175.9814294 MHz
Power 43 dB
on during acquisition
off during delay
W40_HCN_CP modulated
DATA PROCESSING
Gauss apodization 0.106 sec
F1 DATA PROCESSING
Gauss apodization 0.008 sec
FT size 4096 x 2048
Total time 22 min, 6 sec

HSQC Spectrum of

Physangulidine C (3)

SPA14-3-3 (Zhuang Jin)
- 6m in cdcl3

Sample: SPA14-3-3
Sample ID: hammond_SPA14-3-3_20110808_01
File: mnt/ulnmr700/walkup/vnmrsys/data/auto_2011.08.08/hammond_SPA14-3-3_20110808_01/data/Ghmhc_01.fid

Pulse Sequence: gHMBC

HMBC Spectrum of

Physangulidine C (3)

SPA14-3-3 (Zhuang Jin)
6 mg in cdcl3

Sample: SPA14-3-3
Sample ID: hammond_SPA14-3-3_20110809_01
File: mnt/ulnmr700/walkup/vnmrsys/data/hammond_SPA14-3-3_20110809_01/data/Roesy_01.fid

Pulse Sequence: ROESY

Solvent: cdcl3
Temp. 25.0 C / 298.1 K
Sample #6, Operator: hammond
File: Roesy_01
VNMR-400 "ulnmr-ds"

Relax. delay 1.000 sec
Mixing 0.400 sec
Acq. time 0.183 sec
Width 5605.4 Hz
2D Width 5605.4 Hz
16 repetitions
2 x 256 increments
OBSERVE H1, 699.8134459 MHz
DATA PROCESSING
Gauss apodization 0.084 sec
F1 DATA PROCESSING
Gauss apodization 0.084 sec
FT size 4096 x 4096
Total time 3 hr, 40 min, 37 sec

ROESY Spectrum of

Physangulidine C (3)

CURRICULUM VITAE

Name: Zhuang Jin

Home Address:

788 Raymond Kent Ct Apt 3, Louisville, KY 40217

Lab Address:

Chemistry Building, Room 311, Department of Chemistry, University of Louisville, KY 40292

Phone: 502-852-5978 E-mail: jinzhuangzhao@163.com

Personal Information: Born in Inner Mongolia, PR China in Dec 9, 1978. Married to Ms. Lanlan Bao (currently third year graduate student in Department of Chemistry, University of Louisville). Daughter: Solonga Julia Jin, born in Jan 6, 2011, Louisville, KY.

Education:

- 1996 – 2000 BS, Inner Mongolia University GPA:3.30
- 2001 – 2004 MS, Inner Mongolia University GPA: 3.70
- 2007 –present University of Louisville, graduate student GPA: 3.69

Professional experience:

- 2004 – 2007: Lecturer, Inner Mongolia Agricultural University.
- 2007-2012: Graduate research assistant, Graduate teaching assistant of general chemistry (Chem 103 and 201), organic chemistry (Chem 341 and 342) and organic chemistry lab (Chem 343 and 344), University of Louisville.

Honors and awards:

- **Awards Received in the United States:**
- 2010 IMD3 Student Travel Award, University of Louisville
- 2010 David Carew Student Travel Award, from The American Society of Pharmacognosy

- 2008 Graduate Students Lab Safety Awards, University of Louisville
- **Awards Received in China:**
- 2004 Excellent Master Thesis, Inner Mongolia University
- 2002 Guanghua Fellowship, Inner Mongolia University
- 1999 Excellent Students Awards, Inner Mongolia University
- 1998 Guanghua Fellowship, Inner Mongolia University
- 1997 Excellent Students Awards, Inner Mongolia University

Manuscript in Preparation or Submitted

- (2) "Replacement of BF_4^- by PF_6^- , Makes Selectfluor Greener" (in preparation).
- (1) "Stereoselective Synthesis of Fluoroalkyl (*E*)- α,β -Unsaturated Ketones Using a Combination of Gold and Selectfluor" (submitted).

Publications:

- (7) "Physangulidines A, B and C, Three New Antiproliferative Withanolides from *Physalis angulata* L." **Z. Jin**, M. S. Mashuta, N. J. Stolowich, A. J. Vaisberg, N. Stivers, P. J. Bates, W. H. Lewis, G. B. Hammond. *Org. Lett.* **2012**, *14*, 1230-1233.
- (6) "Gold-Catalyzed Intramolecular Oxygen-Transfer Reaction of 2-Alkynyl-1,5-Diketones or 2-Alkynyl-5-ketoesters. Scope, Expansion and Mechanistic Investigation on A Novel [4+2] Cycloaddition" L.-P. Liu, D. Molhotra, **Z. Jin**, R. S. Paton, K. N. Houk, and G. B. Hammond. *Chem. Eur. J.* **2011**, *17*, 10690-10699.
- (5) "Cytotoxic and Antiinfective Phenolic Compounds Isolated from *Mikania decora* (Compositae) and *Crematosperma microcarpum* (Annonaceae)" J. C. Aponte, **Z. Jin**, A. J. Vaisberg, D. Castillo, E. Málaga, W. H. Lewis, C. Sarasara, M. Sauvain, R. H. Gilman, and G. B. Hammond. *Planta Medica.* **2011**, *77*, 1597-1599.
- (4) "Copper Mediated Oxidation of Amides into Imides by Selectfluor" **Z. Jin**, B. Xu, and G. B. Hammond *Tetrahedron Lett.* **2011**, *52*, 1956-1959.
- (3) "Synthetic Evolutions in the Nucleophilic Addition to Alkynes" B. Xu, W. Wang, L. Liu, J. Han, **Z. Jin** and G. B. Hammond *J. Organomet. Chem.* **2011**, *696*, 269-276.
- (2) "Green Synthesis of Vicinal Dithioethers and Alkenyl Thioethers from the Reaction of Alkynes and Thiols in Water" **Z. Jin**, B. Xu and G. B. Hammond, *Eur. J. Org. Chem.* **2010**, 168-173.
- (1) "Antihyperlipidemic Compounds from the Fruit of *Piper Longum* L." **Z. Jin**, G. Borjihan, R. Zhao, Z. Sun, G. B. Hammond and T. Uryu, *Phytother. Res.* **2009**, *23*, 1194-1196.

Patents:

- (2) "Physangulidines A, B and C: Three New Antiproliferative Withanolides from *Physalis angulata* L." patent application is currently underway.
- (1) "Preparation of piperamides and their uses in medicines and health food for reducing blood fat" G. Borjihan, S. Na, **Z. Jin**, R. Zhao, J. Gao, and L. Song, (2005), CN 1634029 A 20050706.

Presentations in National and Regional Meetings:

- (5) "Isolation of bioactive metabolites from targeted peruvian amazonian plants" G. B. Hammond, P. J. Bates, **Z. Jin**, J. C. Aponte, N. S. Stivers, and A. J. Vaisberg. Poster presented at 13th Annual Symposium of the Institute for Molecular Diversity & Drug Design (IMD3), University of Louisville. Mar 8, 2011.
- (4) "Isolation of bioactive metabolites from targeted peruvian amazonian plants" G. B. Hammond, P. J. Bates, **Z. Jin**, J. C. Aponte, N. S. Stivers, and A. J. Vaisberg. Poster presented at the Conference on Innovative Minds in Prostate Cancer Today (IMPACT), Hilton Orlando, Florida, March 9-12, 2011. **Poster:** P4-5.
- (3) "Antileishmanial, trypanocidal, antituberculosis and cytotoxic activities of targeted plant species from the Peruvian rain forest" **Z. Jin**, J. C. Aponte, M. Sauvain, R. H. Gilman, and G. B. Hammond. Oral presentation at the 240th ACS National Meeting, Boston, MA, August 22 – 26, 2010. **Abstract:** AGFD 202.
- (2) "Hydroxyl group directed copper/Selectfluor-activated C-H bond amination" **Z. Jin**, J. C. Aponte, B. Xu, and G. B. Hammond. Oral presentation at the 240th ACS National Meeting, Boston, MA, August 22 – 26, 2010. **Abstract:** ORGN 1.
- (1) "Antileishmanial, trypanocidal, antituberculosis and cytotoxic activities of targeted plant species from the Peruvian rain forest" **Z. Jin**, J. C. Aponte, A. J. Vaisberg, R. H. Gilman, and G. B. Hammond. Oral presentation at the 2010 Joint Annual Meeting of the American Society of Pharmacognosy and the Phytochemical Society of North America, St. Petersburg Beach, Florida, July 10-14, 2010. **Abstract:** O-33.

Membership in Professional Organizations:

1. American Chemical Society, Student Member since 2010.