

University of Louisville

ThinkIR: The University of Louisville's Institutional Repository

Faculty Scholarship

11-1-2022

Advancing the Use of Fecal Sludge for Timelier and Better-Quality Epidemiological Data in Low- and Middle-Income Countries for Pandemic Prevention

Petros Chigwechokha

Renée Street

Rochelle H. Holm

University of Louisville, rochelle.holm@louisville.edu

Follow this and additional works at: <https://ir.library.louisville.edu/faculty>



Part of the [Environmental Health and Protection Commons](#)

Original Publication Information

Petros Chigwechokha, Renée Street, and Rochelle H. Holm. Environmental Science & Technology DOI: 10.1021/acs.est.2c07788

ThinkIR Citation

Chigwechokha, Petros; Street, Renée; and Holm, Rochelle H., "Advancing the Use of Fecal Sludge for Timelier and Better-Quality Epidemiological Data in Low- and Middle-Income Countries for Pandemic Prevention" (2022). *Faculty Scholarship*. 912.

<https://ir.library.louisville.edu/faculty/912>

This Article is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Faculty Scholarship by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. For more information, please contact thinkir@louisville.edu.

Advancing the Use of Fecal Sludge for Timelier and Better-Quality Epidemiological Data in Low- and Middle-Income Countries for Pandemic Prevention

Petros Chigwechokha, Renée Street, and Rochelle H. Holm*



Cite This: <https://doi.org/10.1021/acs.est.2c07788>



Read Online

ACCESS |

Metrics & More

Article Recommendations

KEYWORDS: COVID-19, fecal sludge, low- and middle-income countries, pathogen, wastewater-based epidemiology

The availability and accessibility of water, sanitation, and hygiene (WASH) services are key in preventing disease transmission.¹ Monitoring non-infectious severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) RNA in wastewater can serve as an early indicator of changes in coronavirus disease (COVID-19) cases in a particular catchment area.² Fecal sludge is excreta held on site (unsewered) in sanitation systems, such as at households, in shared outhouses, and in septic tanks. These on-site sanitation systems, a positive step toward improved household WASH services, are used by 43% of the global population, mainly comprising low- and middle-income countries (LMICs).^{1,3} There is an urgent need to expand decentralized (non-sewered system) surveillance as LMICs are currently being overlooked in pandemic prevention as it is currently focused on SARS-CoV-2 wastewater from centralized (sewered) systems.

Wastewater surveillance of centralized systems presents a vastly distinct set of challenges compared with fecal sludge surveillance. For example, sewers also contain gray water and sometimes stormwater, and wastewater is commonly sampled as a composite volume over a 24 h period, whereas fecal sludge is most commonly grab sampled. Broadening the focus of non-sewered sanitation surveillance (NSSS) should accommodate fecal sludge analysis to support globally relevant public health strategies.

Even before the COVID-19 pandemic, fecal sludge was used as a valuable source of information for understanding community health.⁴ Unfortunately, interest in fecal sludge research has been largely limited to a small group of researchers funded by a few core donors. The pandemic has renewed the focus of both science and policy. Surveillance of pathogens in fecal sludge may need to be adapted and could include *Vibrio cholerae* or poliovirus; however, this should be determined by assessing localized needs. The potential of the public health impact in LMICs can be amplified by using fecal sludge data in addition to clinical data in multipathogen disease surveillance and continuous local monitoring for early warnings of a pandemic.

Research partnerships are required to progress from individual clinical patient samples (which has been happening

at a limited scale in healthcare settings) to pooled community samples in terms of fecal sludge. The importance of research partnerships is not new,⁵ but possibly more complex in the Southern African sanitation field with five decades of human waste research.⁶ With no formal global network, international nonprofit institutions can fill this role by bringing together academic research partners and public health officials for interdisciplinary and collaborative research and increased awareness of NSSS. Such partnerships can enable these settings to innovatively adapt academic or healthcare laboratory operations while maintaining high-quality data and also promote peer-reviewed literature originating from resource-limited laboratory settings.

Timelier and better-quality global epidemiological data sets are required for better pandemic preparedness in the future. There is a need in LMICs for the different, but unique, value of NSSS compared to that of wastewater surveillance, built-in capacity at local laboratories that can include fecal sludge as part of continuous multipathogen disease surveillance, and long-term global research collaborations focused on creating systems with data-driven approaches. Challenges will remain, including importation and lengthy delays in supplies and equipment (such as polymerase chain reaction primers and probes that are not manufactured in many LMICs), repurposing existing LMIC laboratory spaces for fecal sludge samples, and limited LMIC access to biosafety level 3 and 4 laboratories for sample processing, thereby limiting pathogen target lists. We recommend that LMICs navigate through these barriers to ensure that fecal sludge surveillance benefits public health, generates local policy-relevant information for future pandemic prevention and control, and holistically supports global WASH services.

Special Issue: Data Science for Advancing Environmental Science, Engineering, and Technology

AUTHOR INFORMATION

Corresponding Author

Rochelle H. Holm – Christina Lee Brown Envirome Institute, School of Medicine, University of Louisville, Louisville, Kentucky 40202, United States; orcid.org/0000-0001-8849-1390; Email: rochelle.holm@louisville.edu

Authors

Petros Chigwechokha – Department of Biological Sciences, Malawi University of Science and Technology, Limbe, Malawi

Renée Street – Environment & Health Research Unit, South African Medical Research Council, Cape Town 7505, South Africa; Environmental Health Department, Faculty of Health Sciences, University of Johannesburg, Johannesburg 2028, South Africa

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.est.2c07788>

Notes

The authors declare no competing financial interest.

Biography



Rochelle H. Holm is an associate professor with the Christina Lee Brown Envirome Institute at the University of Louisville. She received her doctoral degree in Environmental Science from Washington State University in the United States and has lived in Malawi for the past decade. She has broad interests in sanitation systems, water quality, and public health and believes in bridging the gaps across the boundaries of research and innovative practice by collaborating with communities, officials, academics, and analysts. Her current research includes SARS-CoV-2 wastewater surveillance and multipathogen surveillance for pandemic prevention in low-resource settings. Dr. Holm held a former faculty appointment at the Mzuzu University.

ACKNOWLEDGMENTS

This work was supported by a grant from the Rockefeller Foundation, grants from the James Graham Brown Foundation and Owsley Brown II Family Foundation, and a Jon Rieger Seed Grant. The funders had no role in the study design, data collection, and analysis; decision to publish; or manuscript preparation.

REFERENCES

(1) World Health Organization (WHO) and the United Nations Children's Fund (UNICEF). Progress on household drinking water, sanitation and hygiene 2000–2020: Five years into the SDGs. 2021. <https://data.unicef.org/resources/progress-on-household-drinking-water-sanitation-and-hygiene-2000-2020/> (accessed 2022-10-21).

(2) Johnson, R.; Sharma, J. R.; Ramharack, P.; et al. Tracking the circulating SARS-CoV-2 variant of concern in South Africa using wastewater-based epidemiology. *Sci. Rep.-UK* **2022**, *12*, 1182.

(3) Strande, L.; Ronteltap, M.; Brdjanovic, D. *Faecal Sludge Management: Systems Approach for Implementation and Operation*; IWA Publishing: London, 2014. <https://iwaponline.com/ebooks/book/384/Faecal-Sludge-ManagementSystems-Approach-for> (accessed 2022-10-21).

(4) Capone, D.; Chigwechokha, P.; de los Reyes, F. L., III; et al. Impact of Sampling Depth on Pathogen Detection in Pit Latrines. *PLoS Negl. Trop. Dis.* **2021**, *15* (3), No. e0009176.

(5) Perry, M.; Sharp, J.; Aanyu, K.; Robinson, J.; Duclos, V.; Ferdous, R. Research Partnerships Across International Contexts: A Practice of Unity or Plurality? *Development in Practice* **2022**, *32*, 635–646.

(6) Muller, M. Have Five Decades of Development Engineering Research Improved Sanitation in Southern Africa? *J. Int. Dev.* **2020**, *32* (1), 96–111.