

	Frequency (%)
<b>Mother age (n=100)</b>	
< 25 years old	25%
25- 34 years old	60%
35- 44 years old	15%
<b>Accommodation ownership (n=100)</b>	
Rental	99%
Personal	1%
<b>Maternal employment during pregnancy (n=100)</b>	
Unemployed	98%
Full time job	2%
<b>Husband employment (n=100)</b>	
Unemployed	4%
Part time job	11%
Full time job	85%
<b>Maternal education (n=100)</b>	
Illiterate	21%
Primary School	68%
Secondary School	8%
Higher	3%
<b>Number of births (n= 100)</b>	
Secundiparous	30%
Tertiparous	13%
Multiparous ( $\geq 4$ )	57%
<b>Inter-pregnancy interval (with previous pregnancy) (n=100)</b>	
< 6 months	8%
6-12 months	10%
> 1 year	82%
<b>BMI Categories (n=74)</b>	
Underweight (<18.5 Kg/m <sup>2</sup> )	5.40%
Normal weight (18.5- 25 Kg/m <sup>2</sup> )	55.40%
Overweight (25-30 Kg/m <sup>2</sup> )	20.30%
Obese ( $\geq 30$ Kg/m <sup>2</sup> )	18.90%
<b>Classification of total gestational weight gain based on recommendations (n=74)</b>	
Not within recommended range based on pre-pregnancy BMI	75.68% (56/74)
Within recommended range based on pre-pregnancy BMI	24.32% (18/74)
Gained more than the required	20.27% (15/74)
Gained less than the required	55.41% (41/74)
<b>CUTOFF Newborn birth weight (n=100)</b>	
Less than 2500 gram	6%
$\geq 2500$ gram	94%
<b>CUTOFF Hemoglobin level (n=100)</b>	
Less than 11 g/dl	27%
$\geq 11$ g/dl	73%